Digitized by Google

Digitized by Google

570

Phys
464

Digitized by Google

Digitized by Google

~~1182~~

~~Hist. nat~~

~~570.~~

~~Phyt. No. 345.~~

~~753~~

Digitized by Google

PRÆLUDIA BOTANICA

ROBERTI MORISON

SCOTI

ABERDONENSIS.

Digitized by Google

Bayerische
Staatsbibliothek
München

Digitized by Google

HORTUS REGIUS BLESENSIS AUCTUS,

CUM

Notulis durationis & Charactiſmis Plantarum tam additarum, quam non ſcriptarum;

ITEM

Plantarum in eodem horto Regio *Bleſenſi* aucto contentarum, nemini hucuſque ſcriptarum, brevis & ſuccincta delineatio.

Quibus acceſſere

Obſervationes Generaliores (Plantarum in eodem horto Regio *Bleſenſi* aucto contentarum) rei Herbariæ ſtudioſis valdè neceſſariæ, & cognitu perutiles.

PRÆLUDIORUM BOTANICORUM

Pars prior.

AUTHORE

ROBERTO MORISON Medico & Profeſſore Botanico, *ſereniſſimo* CAROLO *ſecundo, magnæ* Britanniæ, *&c. REGI.*

Doctis (in arte) ſcribo:
Indoctos docebo.

LONDINI.

Typis Tho. Roycroft, *Impenſis* Jacobi Alleſtry *ad Inſigne Roſæ Coronatæ in vico vulgò dicto* Duckiane, *Anno* 1669.

Digitized by Google

LABORE.

Digitized by Google

SERENISSIMO
ET
POTENTISSIMO
CAROLO II.
Magnæ Britanniæ, Franciæ, &
Hyberniæ,
REGI.

Sereniſſime REX:

CUM *Hortum Regium* Bleſenſem *auctum*, *pro tertia vice typis committere decreveram. Statim ſubiit animum meum, aliquid de novo addere,*

præter nomenclaturam Authorum, in plantis recitandis: quod Reipublicæ Botanicæ *ejusque emolumento publico inservire queat. Quare ex longa experientia edoctus: durationem Stirpium omnium ibidem contentarum, per notulas* ☉ *scilicet &* ☿, *appositè satis ad eandem indigitandam quadrantes; alios docere æquum judicavi. Plantas præterea* 260. *a nullo Authore descriptas exhibeo, & Charactismo* ♋, *serenissimi principis* Gastonis Franciæ (*Majestatis vestræ*) *Avunculi easdem honoravi. Quia ejus liberalitate & impensis fuerunt acquisitæ. Easque hic breviter & succincte delineavi; ut prodessent omnibus & singulis, quibus volupe est Artis* Botanicæ *studium. Porro observationes generaliores (Plantarum in horto eodem Regio contentarum) rei*

Her-

Digitized by Google

Herbariæ ſtudioſis valde neceſſarias & cognitu perutiles edo. Plantas inſuper 360, *in eodem prædicto horto Regio auctо contentas, plus minuſve, Aſteriſco affectas offero. Quo augmento accrevit idem hortus Regius* Bleſenſis *ab Anno* 1655, *uſque ad Annum* 1660 *quo Anno, extinctus eſt ſereniſſimus, Majeſtatis veſtræ, Avunculus, omnium* Botanicorum *Patronus & Mœcenas vere Regius: verſatiſſimus & oculatiſſimus, in Arte* Botanica *Princeps. Cui ego per decennium, in indagandis & in hortum Regium invehendis plantis, operam navavi iiſdem ex omnibus* Galliæ *tractibus collectis; præter meridionales Alpinos & Pyrenæos Montes: ad quos nos mittere (ſi Dii voluiſſent) decreverat. Quin & methodum meam novam a*

natura datam, a me ſolummodo (citra jactantiam) obſervatam; a nullo niſi meipſo, in hunc uſque diem detectam, quamvis mundi incunabilis ſit coæva; Deſignatoribus, Chalcographis & (id genus) aliis hominibus ingenioſis commiſiſſet, publicique juris feciſſet. Sed proh! dolor atque decus Botanicorum *inevitabili fato, morte prævertus, tam Charus Mœcenas; qui nullis ſumptibus pepercit, ut tam indigenas, in* Gallia *ſponte venientes, quam aliunde, ex omnibus mundi plagis; omnis generis haberet plantas. Nunquam ſatis ob hoc, omnibus* Botanicis *tam præſentibus, quam adventuris & mihi præcipue laudandus. Ego ut viam præſternerem methodo novæ, a me ſolummodo (ut ſupra dictum) detectæ, concinnaturus*

cinnaturus hiſtoriam generalem, omnium Stirpium, tam Arborum, fruticum quam ſuffruticum, per Tabulas cognationis ſeu affinitatis, amicorum quorundam (imo doctorum) Botanicorum precibus, pluries tam hic quam alibi terrarum, verbis coram, literis è longinquo ſolicitatus. Ut has meas primitias (tanquam præludia Botanica, *ad methodum prædictam) in lucem darem. Oratum me ſæpe fateor, tandem me exoratum habuere. Atque ingenii mei, in Arte* Botanica *ſpecimen exhibere volui. Quin & Patronum (Majeſtate veſtra Auguſta) aptiorem eligere non potui, cui collectionem hanc Plantarum, voverem aut dicarem: quicquid eſt, eſt novum. Cui inquam aptius voverem aut inſcriberem, quam M. V.*

quum

quum eundem mihi obtines locum, quem defunctus vester Avunculus. Alumnus sum vester, in vestro solo, antiquo scilicet vestro Regno Scotorum, *natus & educatus. Ad idem Officium vobis reddendum vocatus, eandemque quam apud Avunculum M. V. sustineo hic Provinciam. Ideo M. V. in defuncti Avunculi vestri locum vicemque sufficere, par erat. Quare primitias has nostras, præludiorumve* Botanicorum *horum nostrorum, partem hanc primam accipe (serenissime REX) fronte serena, meque benigne intuere. Dum sublimiora quæ meditor, id est methodum novam, nulli adhuc auditam, per Tabulas cognationis disponentem, omnes vegetabilium classes: Modo faveat Augusta vestra Majestas, nostris conatibus, & laboribus æstate, lucubra-*

lucubrationibus hyeme, per aliquot annos exantlatis: partim ſub Auſpiciis Avunculi veſtri in Gallia, Gaſtonis Franciæ; *partim ſub Auſpiciis veſtris hic. Quod ſi annuere hoc mihi digneris: Polliceor* Britanniam *veſtram, cum methodo exactiſſima (quæ eſt naturæ ipſius) impoſterum, in re* Botanica *gloriari poſſe: quemadmodum,* Italia, Gallia, Germania, *ſuperiori ſæculo, ſine methodo, in Scientia* Botanica, *gloriatæ ſunt. Penes me eandem nunc habeo, ſed unde deſignatorum Chalcographorumque labores compenſentur, non mihi ſuppetit. Tandem ut concludam, quod beatæ memoriæ, ſereniſſimi Principis* Gaſtonis, *Patroni & Domini (quondam) mei digniſſimi vovi; jam tandem partim perſolvi: Cætera (favente Numine) perſolvam. Has noſtras primitias,*

primitias, benigno ſi M.V. intueatur oculo; animum nobis addet, ad majora & ſublimiora prompte edenda.

Sereniſſimæ veſtræ

MAJESTATIS,

Fideliſſimus & devotiſſimus

Servus Domeſticus

MEDICUS

&

Profeſſor *Botanicus*

ROBERTUS MORISON.

PRÆFATIO

AD

Lectorem Botanicum.

CAtalogum omnium Stirpium, in horto Regio Bleſenſi *vegetantium, ſub interitum Mecœnatis ipſius, exhibiturus. Certiorem te redderem (amice lector) neminem in quacunque re fœlicitèr verſari: niſi inclinatione naturæ, fatove vel divinâ ordinatione ad eam trahatur. Quæ omnia in Celſitudine ejus Regiâ,* Aurelianenſi *Duce defuncto, in re* Botanicâ *concurrere videbantur. Nam ab ineunte ætate, (pueritiâ ſcilicet) incepit illi hic ardor* Botanicus, *quique cum adoleſcentiâ accrevit; ætate mediâ, ſtabilitæ & conformata fuit. Ut paucis multa comprehendam, principem oculatiorem, poſt* Salomonem, *in re* Botanicâ *orbis nunquam tulit. Quapropter non potiore jure, naturali propen-*

propensione, Medicinam affectantes; Apollo Æsculapius, Machaon, *atque* Podalyrius, *in Divorum numerum à Veteribus relati: quam serenissimus hic Princeps*, Botanicorum *Divus, posteris denominandus. Observabis (amice lector) plantas omnes hac notulâ* ☉ *affectas, vivaces seu perennes denotari. Plantas vivaces seu perennes appello, quarum radix (postquam deflorverint, semenque produxerint plantæ) novos quotannis emittit Asparagos caulesve, quin & ex eâdem radice, de novo singulis annis reviviscunt plantæ perennes. Arbores & frutices nulla indigitantur notâ, quippe sponte sese declarant vivaces. Plantas autem quibus passim affigitur hæc notula* ♀ *intellige (amice lector) annuas, per annuas, non solùm intelliguntur eæ plantæ, quæ intra anni spatium, semen suum perficiunt, sed & quæ secundo, aut perrarò tertio anno, semine perfecto, exsiccantur pereuntque nulla restibili radice; ex quâ de novo, stolones ut perennes emittant. Has duas notulas* ☉, *&* ♀ *ad plantarum durationem indigitandam elegi: quippè ut aurum, in multos annos ignem facilè perfert, sine perditione; ita plantæ perennes, plurimos Solis*

accessus

acceſſus & receſſus, paritèr facillimè tollerant, ſeſe renovantes de novo quotannis. Mercurii omnium metallorum magìs fluidi & ad ignem poſiti, citiſſimè evanidi notulam hanc ☿ annuis affixi; quippè annuæ intra annum (ut plurimum) pereunt. Nota inſupèr plantas in Catalogo hoc contentas, quibus adjicitur Characliſmus ♃ eſſe non ſcriptas apud Authores, ſed à nobis primò exhibitas. Quoniam ſereniſſimi principis Gaſtonis Franciæ, *imperio, juſſu & expenſis, acquiſitæ & examinatæ fuerunt; ſeu indigenæ in* Gallia, *à nobis collectæ, ſive è longinquo ad nos delatæ eaſdem ipſius Principis Characliſmo honoravimus. Nota plantas Aſteriſco in margine notatas, fuiſſe recuperatas ab Anno* 1655. *uſque ad Annum* 1660. *quo extinctus eſt Mecœnas ipſe ſereniſſimus dux* Aurelianenſis. *Sic quinque illis Annis* 360 *plùs minúſve plantis, noviſſimum hunc à me auctum percipies Catalogum. Deniquè intuendo, ictu oculi percipere potes, quæ Stirpes ſunt vivaces, quæ annuæ, quæque à Celſitudine ejus Regiâ, primò exhibitæ: ipſi, ejuſque memoriæ eas acceptas refer. Cum enim, omnium Stirpium, tam in agris ſponte naſcentium, quam in hortis*

tis Celſitudinis ejus, Regiæ educatarum (ex frequenti omnium ſeminum inſpectione & ad invicèm collatione) durationem, longâ, nempè decennali experientiâ didicimus: hanc liberè Botanicæ *rei ſtudioſis exhibemus. Nec hoc à quovis ante nos factum. Ideò omnibus & ſingulis* Botanicis, *opus gratum ac maximè deſideratum eſſe concludimus. Porrò accipe plantarum* 260 *in eodem horto Regio* Bleſenſi *auctо contentarum; brevitèr & ſuccinctè à nobis factam delineationem. Quin & inſupèr adjecimus (quorundam amicorum precibus) obſervationes generaliores,* Botanicis *valdè neceſſarias & cognitu perutiles; atque ex hiſce ſuprà citatis, conſtruere præludiorum* Botanicorum *noſtrorum partem primam, operæpretium duximus.* Vale.

(Amice Lector)

Tuus humillimus

& ad omnia paratus, ſervus

Rob. Moriſon M. & P.B.R.

SERENISSIMO
ET
CELSISSIMO PRINCIPI,
DOMINO
JACOBO
Eboracenſi & Albanenſi, &c.
DUCI.

Sereniſſime PRINCEPS:

EA *Medicinæ parte* (*quam* Botanicen, *ſeu vegetabilium Scientiam appellamus*) *nullam jucundiorem aut delectabiliorem, unquam judicavit ſereniſſimus veſter Avunculus,*

zunculus, Gaſton. Franciæ, *Dux* Aurelianenſis *in Arte, aut (ſi mavis) Scientia* Botanica, *verſatiſſimus Princeps. Ille enim ab adoleſcentia ad ultimum ſpiritus halitum*; *Scientiam hanc* Botanicam, *rite coluit amavitque. Quin & affirmare audacter audeo*; (*noli credere me Celſitudinis veſtræ Regiæ auribus dare*) *poſt* Salomonem, *in re* Botanica, *oculatiorem ipſo principem*; *orbem nunquam tuliſſe. Hortum ſuum Regium* Bleſenſem; *omnium hortorum, qui unquam in* Europa *extitere, plantis omnis generis refertiſſimum*: *moriens reliquit. Atque vivus & valens, lapſus ſeu hallucinationes* Caſpari Bauhini, Coryphæi *omnium* Botanicorum, *ante ſe ſcribentium*; *obſervare in votis habuit. Nobiſque in mandatis, per aliquot annos,*

Bleſis

Blesis *commorantibus dedit. Quod faciendum in nos suscepimus; non minore ardore, quam si ipse in vivis fuisset. Et jam tandem, partim sub ipsius Auspiciis ante obitum, partim sub serenissimi Domini, Domini nostri, Regis Auspiciis: ut Beatæ memoriæ Principis & Domini mei quondam, (nunquam satis mihi deplorandi) votum persolveremus, eadem* Caspari Bauhini, *in Pinace hallucinationes, tam in digerendis quam denominandis plantis in lucem edimus. Item alterius seu secundi, omnium* Botanicorum Coryphæi, Johannis Bauhini, Caspari *Fratris: animadversiones pariter, in tres Tomos, historiæ suæ universalis; Typis commisimus. Tenuem hunc nostrum laborem, Celsitudini vestræ Regiæ, inscribere & dicare: (quippe quod votum serenis-*

simi Principis Avunculi vestri fuerit) non tenuem puto esse gloriam. Dum duorum Fratrum Caspari & Johannis Bauhini, *hallucinationes seu lapsus observo. Eo magis aliquid, Celsitudini vestræ Regiæ (pro Patrocinio vestro) Encomii advenire putarem: quod ambo, omnium* Botanicorum, *quotquot unquam extitere; (absque controversia) vere* Coryphæi *fuere. Uterque docte, concinne & valde laborioſe scripsit.* Casparus Bauhinus, *Pinacem suum, opus fuisse quadraginta annorum asserit.* Johannes Bauhinus *in compilandis tribus Tomis, historiæ suæ universalis; quinquaginta annos, sese insumpsisse affirmat. Accipe igitur (serenissime PRINCEPS) benigne, hos nostros labores per aliquot annos exantlatos. Et favore tuo eos intueri dignare:*

Et

Et ad majora, & ab omnibus hujus Artis maximopere deſiderata, audenda, nos incitabis, inque æternum tibi devincies.

Sereniſſime Princeps

Veſtrum devotiſſimum

&

Obſequentiſſimum Clientem.

ROB. MORISON.

PRÆFATIO
AD
Lectorem Botanicum.

QVorundam Botanicorum *votis ſatisfacere cupiens, præludia* Botanica *typis committere, animum induxi tandem meum. Cum jam præludiorum meorum* Botanicorum, *pars prior, penitùs prelo exceſſerit. Hallucinationes* Caſpari Bauhini *in Pinace, tàm in digerendis, quàm denominandis plantis; à me obſervatas, habes. Item animadverſiones in tres tomos, hiſtoriæ univerſalis* Johannis Bauhini *offero. Eaſque publici juris facio; ut penes quos eſt ejuſdem* Johannis Bauhini, *plantarum hiſtoria univerſalis, tribus tomis digeſta: ad manum paritèr haberent, ipſius errata. Noli quæſo (amice lector) me glorioſum aut inſolentem exiſtimare ſcriptorem: quod non ſolùm hujus ſæculi, ſed omnium (quotquot adhuc extitêre)* Botanicorum; *Coryphæos duos,* Caſparum & Johannem Bauhinum *Fratres, corrigo. Ex frequenti enim ipſorum*

ipſorum voluminum inſpectione, & ex longâ & diutinâ eorundem examinatione. Ipſos, ſummi judicii, indefeſſi laboris, & incomparabilis doctrinæ (*in Scientiâ* Botanicâ) *homines fuiſſe obſervo: ipſoſque ſeu fato, ſeu naturali propenſione; maximum in promovendis ſtudiis* Botanicorum, *habuiſſe deſiderium; paritèr declaro. Erráſſe tamen multotiès utrumque, nemo* (*in re* Botanicâ, *mediorcitèr verſatus*) *inficias ibit; hos meos tractatulos legendo. Labi nihilominùs, humanum eſſe confiteor; homo ſum ipſe, humani à me nihil alienum puto; in hiſce meis tractatulis, hallucinatum me eſſe non dubito. Quapropter hallucinationes meas, ab amico lectore, notari deſidero. Quin & paritèr obſervabit lector, me hæc concináſſe, tàm ut voto meo prædicto ſatisfieret, quàm ut publicis inſervirem commodis. Et quam ego de Fratribus* Bauhinis Coryphæis; *eandem de me exiſtimationem habeat lector, velim.* Vale,

(Amice Lector)

Tuus humillimus

ſervus & ad omnia paratus,

Rob. Moriſon M. & P.B.R.

In Hortum Regium BLESENSEM.

HINC, nulli biferi miranda Roſaria peſti.
Nec mala heſperidum, vigili ſervata Dracone.
Si Paradiſiacis quicquam (ſine crimine) campis;
Conferri poſſit. Blæſis *mirabile ſpecta!*
Magnifici Gaſtonis *opus: qui terra capaci;*
Quicquid alit fœcunda ſinu, plantaſque tenellas;
Robur & Arboreum, modico conſevit in Horto.
Berſabidem perhibent, moviſſe à Graminis herba,
Ad Libani *Cedros. Coluit propriiſque locavit*
Sedibus hic princeps: ſiccis montana reponens
Udis ima locis. Ut vultu cuncta venirent
Nativo, facilique florent tibi nota labore.

poſuit JACOBUS METELANUS Scotus

Ad Lectorem BOTANICUM.

DEſine mirari claris Babylonis *in Hortis.*
Pendula, coctilibus, ſemina fulta pilis.
Nec ſeſe Alcinoi *jactent Pomaria, plures*
Plantas Bleſenſis *nobilis Hortus alit,*
Quas decus Hectoridum variis diſtinxit arenis
Gaſtonus, *populo Pharmaca ſana ſuo:*
Principibus cædes multis, Laus maxima habetur.
Cura erat huic hominum, principe digna, ſalus.
Regia progenies, mens Regia, Regia virtus,
Regia vita, obitus Regius ipſe fuit.
Gaſtoni *Auſpiciis magni, collegerat herbas*
Quotquot in Europa, *quotquot uhique virent,*
Primus & Aſtragalos meminit, Scutellariaſque
Moriſonus, *Patriæ ſplendida fama ſuæ.*

Moriſoni nomen cunctis celebrabitur Hortis:
Dum ſua cum foliis ſemina planta feret.

Canebat CLAUDIUS CRITCHTONIUS Scotus.

In Horti Regii BLESENSIS, ejuſque Authoris Encomium: ſcribit Authoris amicus.

QUÆ Dives Natura mihi miracula jactat,
Theſauros pandens Ambitioſa novos?
Mirandum Gaſtonis *opus! dabit Area, quicquid*
Orbe datum eſt; ingens in nuce mundus adeſt.
Quicquid ubique viget plantarum, hæc horrea ſervant,
Hic ſtupet ornatus flora vel ipſa ſuos,
Area ſparſa ſuis ſtellis, velut Æthera, luce,
Invidioſa polis, inferiore micat;
Nec Moriſone *minor Tua gloria; Quælibet herba*
Naturæ paret legibus, atque Tuis,
Sceptra tenens, clauſo plantarum in carcere regnas
Surgit & Auſpiciis herba caditque Tuis.
Quaſque prius terris, nunc chartis inſeris herbas
Et datur in Libris Vita ſecunda Tuis.
Quæque ſuis foliis poterat tam ſæpe perire
Non poterit foliis planta perire Tuis.
Quid quod & hæc tantum vibras luſoria tela
Ridet & hæc tanquam ludicra majus opus.
Non aliter Liber hic quam magni Anteambulo Regis
Sive ſuum veluti Phoſphorus ante diem
Illa edas cunctis optata volumina votis
Illud, quo majus non habet Orbis, opus.
Illic innumeras Tu digeris ordine plantas
Illic in proprios herba redacta locos
Illa Tibi fient monumenta perennia; vitam
Quam dederas herbis & dabit herba Tibi.

ALEXANDER MORISON *A. M.*

Explicatio nominum Authorum, in Horto Regio *Bleſenſi* auctorum Citatorum.

AD. & Adv. Lob. Lobelius in Adverſariis.

Ad. Lob. part. pr. & part. alt. Lobelius in Adverſariorum parte prima, & parte altera.

Ama. Amat. Amatus Luſitanus.

Apul. Apuleius.

Bapt. Ferr. & Ferrar. Joh. Baptiſta Ferratyus.

Bell. Bellon. Petrus Bellonius.

Bauh. Pin. B Pin. Caſparus Bauhinus in Pinace.

Bauh. Prod. B. Prod. & Prodr. Caſparus Bauhinus in Prodromo.

Bauh. & B. Phytop. Caſparus Bauhinus in Phytopinace.

B. Pin. app. Bauhinus in Pinacis appendice.

C. B. Pin. Caſparus Bauhinus in Pinace.

Caſp. Bauh. in Pin. C. B. P. aliquando C. B. Caſparus Bauhinus in Pinace.

C. B. Prod. descr. Casparus Bauhinus in Prodromo, descriptione.

Cæs. & Cæsalp. Andreas Cæsalpinus.

Cam. & Camer. Camerarius in Epitome Matthioli & horto suo.

Cast. Castor Durantes.

Clus. hist. vel Clus. Carolus Clusius in historia sua rariorum.

Clus. desc. Clusius in historia rariorum descriptione.

Clus. cur. post. Carolus Clusius in curis posterioribus.

Clus. ex. & exot. Carolus Clusius in historia exoticarum.

Clus. hist. app. 1. & app. alt. Carolus Clusius in appendice prima & altera ad historiam.

Col. & Colum. Fabius Columna in historia de minùs cognitis.

Col. Phytob. Fabius Columna in Phytobasano.

Cord. & Cord. hist. Cordus in historia sua Plantarum.

Corn. & Cornuti, Jacobus Cornutus de plantis suis Canadensibus.

Dalech. & Dal. Dalechampius. Dal. & Dalechamp. Dalechampii hist. Lugd. est historia Lugdunensis citans Dalechampium.

Dioſc. Dioscorides.

Dod.

Dod. Dodonæus in historia pemptadum, editione Latina in folio.

Dod. Gall. ed. Dod. ut. & utr. Dodonæus in editione Gallica, & utraque tam Gallica (à Carolo Clusio edita) quam sua Latina.

Dod. desc. Dodonæus in folio Latinè, descriptione.

Desc. & descr. post citationem Authoris, denotant plantam ab ipso Authore citato scriptam, non figuratam: ut Lob. Dod. Clus. &c. desc. significat ab ipsis esse descriptam.

Eyst. & Eystet. & hort Eyst. hortus Eystettensis à Besslero editus.

Fuch. Fuchius in historia sua Latina, in folio.

Ger. Johan. Gerardus, emaculatus à Jansono.

Ges. Gesn. Gesnerus in corrallario, historiæ Cordi, addito: & in Libello de collectione Stirpium

Hist. Lugd. & Lugdun. historia Lugdunensis Latinè edita in folio.

Hist. & histor. Mex. & Mexican. historia Mexicana in folio.

Hist. ſ. histor. Brasil. historia Brasiliana in folio.

Hort. Farnes. & Farn. hortus Farnesianus.

Hort. Eyst. & Eystet. hortus Eystettensis.

Icon. Lob. Lobelius in Iconibus.

Imperat. ferantes Imperatus in ſua hiſtoria naturali.

Joh. Bauh. & J. B. Johannes Bauhinus. in univerſali ſua hiſtoria, tribus Tomis digeſta.

Lob. Icon. obſ & Ad & Adver. Lobelius in Iconibus, forma oblonga, obſervationibus & Adverſariis Stirpium: in folio.

Lug. & Lugdun. hiſtor. hiſt Lugd. & Lugdun. hiſtoria Lugdunenſis, Lugduni excuſa, lingua Latina: in folio.

Matth. & Matt. ed. Bauh. Mattheoli editionis Bauhini citatio.

Maran. & Maranthæ, Bartholomæus Maranta.

Muton. & Mutoni. Mutonus ab Authoribus citatus.

Obſ. & Icon. Lob. Lobelius in Obſervationibus & Iconibus ſuis.

Obſ. Lob. & ad. Lob. Lobelius in Obſervationibus & Adverſariis ſuis.

Park. Johannes Parkinſon, in ſuo Theatro Botanico.

Park. in appad. Theat. Joh. Parkinſon in appendice ad Theatrum ſuum Botanicum.

Park. parad. idem Joh Parkinſon in ſuo Paradiſo terreſtri.

Paſſ.

Paſſ. Paſſæi. Paſſæus in Iconibus ſuis in forma oblonga.

Plin. Plinius.

Pon. & Ponæ, Johannes Pona in editione ſua Latina.

Pon. & Ponæ deſc. Mont. Bald. Johannes Pona in Montis Baldi deſcriptione.

Proſp. Alp & Alpin. de plant. & plan. Ægypt. Proſper Alpinus de plantis Ægypti.

Proſp. Alp. exot. Proſper Alpinus de exoticis.

Rob. Steph. Præd. Ruſt. Robertus Stephanus in Prædio Ruſtico ſuo.

Ruel. & Ruell Johannes Ruellius.

Stapel. & Stapelii, Johannes Boda Stapelius, noviſſimus in Theophraſtum Commentator.

Swert. Swertius in florilegio ſuo.

Tab. & Tabern. Theodorus Tabernemontanus in Iconibus ſuis in forma oblonga.

Thal. & Thalii, Johannes Thalius in ſylva ſua Harcynica.

Theoph. & Theophr. Theophraſtus.

Trag. & Tragi, Hieronymus Tragus in hiſtoria.

Veſl. & Veſling. Obſ. in Proſp. Alp. Veſlingii Obſervationes in Proſperum Alpinum.

Valet. hortus valet, Acupictorum in uſum factus.

Deſc.

Desc. & descript. intellige Authoris citati plantam scriptam, non figuratam.

Col. & Colum. in hallucinationibus Caspari Bauhini *à nobis primo exhibitis, Columnam designat. I. & II. Pl. primam & secundam demonstrat (post paginam designatam) plantam prædictæ Columnæ.*

Quicunque plura, de Authoribus desiderat; consulat Casparum Bauhinum, *sub introitum Pinacis sui, ibi prolixiùs docet. Qui fuêre Authores omnes, qui (ante suum sæculum) de re* Botanica *scripsêre: & quo volumine & quandò ipsorum opera, in lucem prodiêre.*

Nota insuper, quando nomina Theophrasti, Dioscoridis, Plinii, Mattheoli Tragi, Thalii, Lobelii, Dodonæi, Clusii Fuchsii, Dalechampii, Ponæ, *& cæterorum (id genus) Authorum aliorum; ad longum, ab aliis citantur; scribuntur ut suprâ. Et citantes ipsos curtè in* Botanographorum *monumentis, in medium adferuntur ut passim à nobis, & aliis antè nos scribentibus facilitatum, observare licet.*

HOR-

HORTUS REGIUS BLESENSIS AUCTUS.

A

ABIES Hiſt. Lugd. alba ſive fœmina B. pin.

Abrotonum mas Dod. mas anguſtifolium majus B. pin. ⊙

Abrotonum mas incanum Col. mas anguſtifolium incanum B. pin. ⊙

* Abbrotanum mas anguſtifolium maximum C. B. P. magnum camphoratum quibuſdam incenſaria cam. ⊙

Abrotonum odoratum humile Dod. mas anguſtifolium minus B. pin. ⊙

Abrotonum campeſtre B. pin. Artemiſia tenuifolia Dod. ⊙

Abrotonum fœmina vulgare Cluſ. fœmina foliis teretibus B. pin. ⊙

Abrotonum fœmina Narbonenſe Cluſ. fœmina flore majore B. pin. ⊙

Abrotonum fœmina viride B. pin. fœm. ſextum Cluſ. deſcr. ⊙

Abſinthium vulgare Fuchſ. latifolium Dod. Ponticum, ſeu Romanum officinarum, ſeu Dioſcoridis B. pin. ⊙

Abſinthium inſipidum, Abſinthio vulgari ſimile B. pin. ♀

Abſinthium maritimum Lavendulæ folio B. pin. latifolium marinum hort. Eyſt. ⊙

Abſinthium marinum Latifolium Park. ⊙

Abſinthium Seriphium vulgare, perperam Dioſcoridis Icon. Lob. Seriphium Belgicum B. pin. ⊙

Abſinthium Seriphium Gallicum Matth. ed. & B. pin. Narbonenſe Dod. deſcr. ⊙

Abſinthium tenuifolium Dod. Ponticum Matth. ed. B. Ponticum tenuifolium incanum B. pin. ⊙

Abſinthium Auſtriacum tenuifolium Cluſ. Ponticum, Tridentinum herbariorum Obſ. Lob. Ponticum tenuifolium Auſtriacum B. pin. ⊙

Abſinthium

Abſinthium Alpinum umbelliferum Cluſ. Alpinum umbelliferum latifolium B. pin. ☉

Acacia Americana Robini, Cornuti.

Acacia Ægyptiaca hiſt. Lugd. foliis ſcorpioidis leguminoſæ B. pin.

Acanthium Dod. Spina alba tomentoſa latifolia ſylveſtris B. pin. ♀

Acanthium Illyricum Obſ. Lob. ♀

Acanthium Luſitanicum. ♃ ♀

* Acanthium Illyrico accedens capitulis minoribus. ♃ ♀

Acanthus ſativus Dod. ſativus vel mollis Virgilii B. pin. ☉

Acanthus ſylveſt. Dod. aculeatus B. pin. ☉

Acarna flore luteo patulo B. pin. Carlina ſylveſtris minor Dod. ♀

Acarna flore purpurorubente patulo B. pin. Acanthioides parva Apula Col. ♀

Acarna major caule folioſo B. pin. Chamæleon Salmanticenſis Cluſ. ♀

Acer major Dod. montanum candidū B. pin.
Acer montanum flavum .ſ. criſpum B. pin.
Opulus montanus hiſt. Lugd. deſcr.
Acer minor Dod. campeſtre & minus B. pin.
Acer Monſpeſſulanum hiſt. Lugd. trifolia B. pin.

* Acetoſa criſpa I. B. ☉
Acetoſa pratenſis B. pin. Oxalis Dod. ☉
Acetoſa montana maxima B. pin. Oxalis major Broccenbergenſis Thal. ſyl. Harcyn. ☉
Acetoſa veſicaria peregrina hort. Eyſt. Americana foliis longiſſimis pediculis donatis B. pin. & prodr. ♀
Acetoſa rotundifolia hortenſis B. pin. Oxalis rotundifolia Dod. ☉
Acetoſa Cretica ſemine aculeato B. prodr. & pin. ♀
Acetoſa arvenſis lanceolata B. pin. Oxalis minima Dod. ☉

Aconitum hyemale Ponticum hort. Eyſt. Unifolium luteum bulboſum B. pin. ☉
Aconitum lycoctonon luteum majus Dod. Lycoctonon luteum B. pin. ☉

Aconitum

Aconitum lycoctonon luteum minus Dod. deſcr. ⊙

Aconitum lycoctonon flore Delphinii Dod. cœruleum glabrum flore conſolidæ regalis B. pin. ⊙

Aconitum cœruleum .ſ. Napellus primus B. pin. Napellus Dod. ⊙

Aconitum violaceum .ſ. Napellus ſecundus B. pin. Lycoctonon quartum Tauricum Cluſ. ⊙

Aconitum lycoctonon quintum Neubergenſe Cluſ. purpureum .ſ. Napellus tertius B. pin. ⊙

Aconitum lycoctonon cœruleum parvum Dod .cœruleum minus.ſ. Napellus minor B. pin. ⊙

Aconitum ſpica florum pyramidali. ♃ ⊙

Aconitum novum Judenbergenſe Cluſ. ⊙

Aconitum ſalutiferum .ſ. Anthora B. pin. Anthora Dod. ⊙

Aconitum racemoſum B. pin. Chriſtophoriana Dod. ⊙

Aconitum baccis niveis Cornuti. ⊙

Aconitum baccis rubris Cornuti. ⊙

Acorus verus .ſ. Calamus aromaticus officinarum

narum B.pin. Acorum legitimum Cluſ. ☉

Adiantum foliis Coriandri B. pin. Adiantum ſ. Capillus Veneris verus Ad. Lob. ☉
Adiantum Americanum Cornuti. Fruticoſum Braſilianum B. prodr. ☉
Adiantum nigrum Plinij Lob. Dryopteris nigra Dod. ☉

Adonis Matth. ed. B. ſylveſtris flore phœnicéo B. pin. ♀
Adonium luteum Tab. ♀

Æthiopis Dod. foliis ſinuoſis B. pin. ♀

Ageratum Matthioli hiſt. Lugd. foliis ſerratis B. pin. ☉
Ageratum floribus candidis Obſ. Lob. ☉
Ageratum purpureum hiſt. Lugdun. ſerratum Alpinum B. pin. ☉
Ageratum villoſum purpureum. ♃ ☉

* Agrimonoides Col. ☉
Alaternus prima Cluſ. Philyca elatior B.pin.
Alaternus foliis ex albo variegatis. ♄
Alaternus foliis ex luteo variegatis. ♄

Alcea

Alcea vulgaris Dod. vulgaris major B. pin. ☉

Alcea folio rotundo laciniato B. pin. Malva .ſ. Alcea montana Col. ☉

Alcea fruticoſa cannabino folio Cluſ. cannabina B. pin. ☉

Alcea Syriaca hort. Eyſt. arboreſcens Syriaca B. pin. Althæa frutex ſecunda Cluſ. ☉

Alcea Syriaca flore candido Hort. Eyſt. ☉

Alcea Veneta hort. Eyſt. veſicaria Dod. B. pin. ♀

* Alcea veſicaria capitis bonæ ſpei ♃ ♄

Alcea villoſa Dalechampij hiſt. Lugd. hirſuta B. pin. ♀

Alcea Indica parvo flore B. pin. Ægyptia Cluſ. Bamia Alexandrina Cæſalp. ♀

Alcea Ægyptiaca villoſa B. pin. Bamia moſchata Veſlingij Obſ. in proſp. Alp. ♀

Alchimilla Dod. vulgaris Cluſ. B. pin. ☉

Alchimilla minor. ♃ ☉

Alchimilla minima montana Col. Saxifraga .ſ. Perchepier Anglorum Icon. Lob. Chærophillo

rophyllo nonnihil ſimilis B. pin. ♀

Alliaria Dod. B. pin. ♀

Allium Dod. ſativum B. pin. ☉
Allium ſphæriceo capite, folio latiore B. pin. Ampelopraſum Dod. Scorodopraſum primum Cluſ. ☉
Allium Urſinum Fuchſ. ſylveſtre latifolium B. pin. ☉
Allium anguinum Matthioli hiſtor. Lugdun. montanum latifolium maculatum Bauh. pin. Victorialis longa Cluſ. ☉
Allium ſylveſtre tenuifolium Dod. campeſtre juncifolium capitatum purpuraſcens majus B. pin. ☉
* Allium inſipidum juncifolium Gethoïdes ſylveſtre Col. ☉
Allium montanum bicorne flore obſoletiore B. pin. Moly lurido colore Swertij. ☉
Allium juncifolium bicorne luteum B. prodr. & pin. ☉
Allium montanum capite rotundo B. pin. Allium ſeu Moly montanum quintum Cluſ. ☉
Allium montanum foliis Narciſſi majus B. pin.

pin. Allium .ſ. Moly Narciſſinis foliis primum Cluſ. ☉

Alnus Dod. vulgaris Cluſ. rotundifolia glutinoſa yiridis B. pin.
Alnus nigra baccifera hiſt. Lugd. B. pin. Frangula Dod.
Alni effigie lanato folio major B. pin. Aria Theophraſti effigie Alni Ad. & Icon. Lob.
Alni effigie lanato folio minor B. pin. Amelanchier Gallorum Obſ. Lob.

Aloe Dod. vulgaris B. pin. ☉
Aloe ex America Dod. folio in oblongum aculeum abeunte B. pin. ☉

Alſine heredulæ folio B. pin. hederacea Tab. ♀
Alſine foliis Veronicæ Tab. Veronicæ foliis, floſculis cauliculis adhærentibus B. pin ♀
Alſine foliis Triſſaginis Icon. Lob. chamædryfolia floſculis pediculis oblongis inſidentibus B. pin. ♀
Alſine triphyllos cœrulea B. pin. Elatine triphyllos hiſt. Lugd. ♀
Alſine ſcandens baccifera B. pin. Cucubalus Plinii hiſt. Lugd. ☉

Alſine altiſſima nemorum B. pin. major Dod. ♀
Alſine media B. pin. minor Dod. ♀
Alſine minor multicaulis B. pin. minima Dod. ♀
Alſine petræa minima Tab. minor Lini capitulis B. pin. ♀
Alſine affinis Androſace dicta major B. pin. Androſace altera Matthioli Cluſ. ♀
Alſine ſpergula dicta major B. pin. Sagine ſpergula Obſ. Lob. ♀
Alſine ſpergulæ facie .ſ. Spergula minor ſubcœruleo floſculo B. prodr. ♀
* Alſine ſpergula flo. violacio major C. B. P. ſpergula marina Dal. ☉
* Alſine ſpergula ſemine foliaceo nigro. ♃ ♀
* Alſine ſpergula, ſemine foliaceo ſubfuſco. ♃ ♀
Alſine fontana Tab. aquatica media B. pin. ♀
Alſine littoralis foliis portulacæ B. pin. Alſines Pelagicum genus Cluſ. ☉
Alſine nodoſa Germanica B. prodr. ♀
Alſine tenuifolia muſcoſa B. pin. muſcoſa hiſt. Lugd. ♀

Alſine

Alſine inaperto flore. Noctiflora minima ☌ ☿

Alſine hirſuta magno flore B. pin. Auricula muris quibuſdam Dod. Gall. ed. ☿

* Alſine hirſuta viſcoſa altera C. B. P. ☿ Folia hujus anguſtiora & ſaturatius virentia ſunt.

Althæa, Ibiſcus Dod. vulgaris Cluſ. Althæa Dioſcoridis & Plinij B. pin. ☉

Althæa arborea Olbiæ Lob. fruteſcens folio acuto, parvo flore B. pin. ☉

Althæa Theophraſti flore luteo B. pin. Abutilon Avicenæ hort. Eyſt. ☿

* Althæa flo. luteo minimo, ſemine roſtrato. ☌ ☿

Alyſſon Galeni Dod. verticillatum foliis profundè inciſis B. pin. ☉

Amaranthus maximus B. pin. Quinüa, ſ. Blitum majus Peruanum Cluſ. Idem ſemine rubro. ♃

Amaranthus coccineus elegans maximus Park. ☿

Amaranthus maximus ſemine nigro. ☌ ☿

Amaranthus

Amaranthus ſpicis viridibus. ☌ ♀
Amaranthus criſtatus Camerarii hort. Eyſt. paniculâ incurvâ B. pin. ♀
Amaranthus purpureus Dod. paniculâ conglomeratâ B. pin. ♀
Amaranthus ſpicatus hort. Eyſt. ſimplici paniculâ B. pin. ♀
Amaranthus tricolor hort. Eyſt. Icon. Lob. folio variegato B. pin. ♀
* Amaranthus ſpica erecta viridi. ☌ ♀
* Amaranthus ſpica erecta obſoletè rubra ☌ ♀
* Amaranthus ſpica erecta rubra, ſemine albo. ☌ ♀
* Amaranthus auriculatus luteus Chriſtatus. ☌ ♀
* Amaranthus auriculatus luteus ſpicatus ☌ ♀
* Amaranthus auriculatus Phœniceus ſpicatus. ☌ ♀
* Amaranthus auriculatus Phœniceus criſtatus. ☌ ♀

Ambroſia campeſtris repens B. pin. Coronopus repens Lob. hiſt. Lugd. ♀

Ammi vulgare Dod. majus B. pin. ♀
Ammi Matth. ed. B. ♀ A-

Amygdalus Dod. ſativa B. pin.
Amygdalus pumila ♃ ☉
* Amygdalus putamine fragili. ♃

Anagallis phœniceo flore B. pin. mas phœnicea Icon. Lob. ♀
Anagallis flore cœruleo B. pin. fœmina cœrulea Icon. Lob. ♀
Anagallis tenuifolia Monnelli. Cluſ. app. alt. ☉
Anagallis lutea nemorum B. pin. flavo flore Cluſ. ☉
Anagallis omnium minima. ♃ ♀
Anagallis aquatica major folio oblongo B. pin. Berula major Tab. ☉
Anagallis aquatica minor folio ſubrotundo B. pin. aquatica .ſ. Beccabunga Germanorum Ad. Lob. ☉
Anagallis aquatica folio rotundo non crenato B. pin. aquatica tertia Icon. Lob. ☉
Anagallis aquatica anguſtifolia B. pin. aquatica quarta Icon. Lob. Eadem flore albo. ☉

Anagyris Dod. fœtida B. pin.
Anagyris non fœtida major vel Alpina B. pin. Laburnum Dod. Gall. ed.

Anchuſa

Anchuſa puniceis floribus B. pin. minor .ſ. Alcibiadion, vel Onochiles Lob. ⊙
Anchuſa lignoſior anguſtifolia Ad Lob. anguſtifolia B. pin. ⊙

Androſæmum Dod. maximum fruteſcens B. pin. ⊙
Androſæmum Campoclarenſe Col. Aſcyrum .ſ. Hypericum bifolium glabrum non perforatum B. pin. ⊙
Androſæmum alterum Apulum Col. perfoliatum & perforatum B. pin. ⊙
Androſæmum alterum hirſutum Col. hirſutum B. pin. ⊙
Androſæmum fœtidum .ſ. Tragodes hort. Eyſt. ⊙

Anemone tenuifolia flore ſimplici dilutè purpureo. ⊙
Anemone tenuifolia flore ſimplici igneo. ⊙
Anemone tenuifolia flore ſimplici violaceo. ⊙
Anemone tenuifolia flore ſimplici albo, circa caulem purpuraſcente. ⊙
Anemone tenuifolia flore ſimplici ex albo rubeſcente. ⊙

Anemone tenuifolia flore ſimplici ſubalbido. ⊙

Anemone tenuifolia flore ſimplici dilutè violaceo. ⊙

Anemone flore pleno coccineo foliolis interius ianthinis, *Amaranthe.* ⊙

Anemone flore pleno foliis & foliolis coccineis. *Double rouge.* ⊙

Anemone flore pleno ex viridi rubeſcente, foliolis interius viridibus. ⊙

Anemone ex albo & igneo variegata. *Orlate Romaine.* ⊙

Anemone Orlate magis albeſcens. *Orlate d' Angleterre.* ⊙

Anemone flore pleno flammeo. ⊙

Anemone flore pleno dilutè violaceo. ⊙

Anemone flore pleno exiguo albo. ⊙

Anemone flore pleno hyacinthino. *Illuſtre de Leon.* ⊙

Anemone florum mali perſicæ quoad colorem æmula. ⊙

Anemone multiplex flore Ranunculi. ⊙

Anemone tenuifolia hermaphrodita. ⊙

Anemone tenuifolia ſcala dicta. ⊙

Anemone tenuifolia Aurelianenſis flore pleno eleganter variegato. ⊙

Anemone tenuifolia flore pleno coccineo Paſſæi Icon. tenuifolia multiplex purpurea B. pin. *Cramoiſine.* ☉

Anemone latifolia pavo minus polyphylla B. pin. hortenſis latifolia pavo major ſecunda Cluſ. ☉

Anemone tertia Matth. ed. B. Sylveſtris prima Cluſ. Sylveſtris alba major B. pin. ☉

Anethum Dod. hortenſe B. pin. ♀

Angelica major Dod. ſativa B. pin. ♀

Angelica ſylveſtris Dod. Sylveſtris major B. pin ♀

Angelica ſylveſtris minor .ſ. erratica B. pin. Podagraria Germanica .ſ. Belgica Obſ. & Icon. Lob. ☉

Angelica montana maxima. Archangelica Dod. Cluſ. ♀

Angelica lucida Canadenſis Cornuti. ♀

Aniſum Dod. ♀

Anonis ſpinoſa flore purpureo B. pin. Anonis Fuchſ. Dod. Eadem flore albo quæ Anonis flore albo Tab. ☉ Eadem non ſpinoſa purpurea. ☉ A-

Digitized by Google

Anonis floribus luteis exiguis. Ononis lutea ſylveſtris minima Col. ☉

Anonis non ſpinoſa flore luteo variegato B. pin. Ononis flore luteo variegato non ſpinoſa hort. Eyſt. ☉

* Anonis fruteſcens verna ſiliqua hirſuta pendula. ♃ ☉

* Anonis ſiliqua hirſuta pendula flo. luteo, viſcoſa minor flo. pendulo B. prod. ♀

* Anonis ſiliqua hirſuta pendula, flo. purpureo; minima hirſuta Cherleri J. B. ♀

Anonis mitis annua purpurea. ♃ ♀

Anthillis prior lentifolia maritima Peplios effigie Ad. & Icon. Lob. Portlandiæ lentifolia Penæ hiſt. Lugd. ☉

Anthyllis marina incana alſinefolia Narbonenſium Lob. maritima alſinefolia B. pin. ☉

Antirrhinum majus rotundiore folio B. pin. Cynocephalus Plinij hiſt. Lugd. ♀

Antirrhinum Dod. majus alterum folio longiori B. pin. ♀ Idem quod

Antirrhinum flore albo hort. Eyſt. ♀

Antirrhinum flaveſcens Dod. deſcr. ♀

Antirrhinum ex dilutè albo purpur aſcens Dod. deſcr. ♀
Antirrhinum arvenſe majus B. pin. primum Matth. ed. B. Idem flore albo. ♀

Aparine Dod. vulgaris B. pin. ♀
Aparine ſemine lævi, lævis Park. deſc. ♀
Aparine fructu verrucoſo. ſemine coriandri ſaccharati Park. deſc. ♀

Apios Americana Cornuti. ⊙

Apium hortenſe Dod. hortenſe .ſ. Petroſelinum vulgò B. in pin. ♀
Apium vel Petroſelinum criſpum B. pin. Petroſelinum criſpum Cam. hort. med. ♀
Apium hortenſe maximum B. prodr. ♀
Apium montanum folio tenuiore B. pin. Petroſelinum Dalechampii hiſt. Lugd. ♀
Apium montanum folio ampliore B. pin. Oreoſelinum Cluſ. ⊙
Apium ſylveſtre .ſ. Thyſſelium Dod. ſylveſtre lacteo ſucco turgens B. pin. ⊙
Apium paluſtre, & Apium officinarum B. pin. Eleoſelinum Dod. ♀
Apium Macedonicum B. pin. Petroſelinum Mace-

Macedonicum Ad. Lob. ♀

* Apium peregrinum i. Cluſii peregrinum fol. ſubrotundis C. B. P. ⊙

Apocynum folio ſubrotundo B. pin. primum latifolium Cluſ. ⊙

Apocynum folio oblongo B. pin. Periploca ſerpens anguſtiore folio Obſ. & Icon. Lob. ♀

Apocynum majus Syriacum rectum Cornuti. ⊙

Apocynum rectum minus Canadenſe Cornuti. ⊙

* Apocynum tertium latifolium Cluſ. hiſt; Periploca prior Dod. ⊙

* Apocynum Canadenſe foliis Androſemi. ♃ ⊙

* Apocynum Canadenſe anguſtifolium flo. aureo. ♃ ⊙

* Apocynum Canadenſe maximum flore minimo herbaceo. ♃ ⊙

Aquifolium Matth. ed. B. Ilex aculeata baccifera folio ſinuato B. pin.

Aquilegia Fuchſ. ſylveſtris B. pin. ⊙

Eadem quæ

Aquilegia flore albo ſimplici hort. Eyſt. ☉

Aquilina rubra Icon. Lob. ☉

Aquilegia pumila præcox Canadenſis Cornuti. ♀

Aquilegia hortenſis multiplex flore magno B. pin. Aquileia multiplici flore cœruleo Dod. ☉

Aquilegia flore pleno albo hort. Eyſt. ☉

Aquilegia flore pleno incarnato hort. Eyſt. ☉

Aquilegia flore pleno verſicolori. ♃ ☉

Aquilegia flore roſeo multiplici B. pin. ſtellata flore purpureo hort. Eyſt. ☉

Aquilegia ſtellata flore violaceo hort. Eyſt. ☉

Aquilegia ſtellata flore albo. ☉

Arbutus Dod. folio ſerrato B. pin.

Argemone capitulo torulis canulato Obſ. Lob. capitulo breviori B. pin. ♀

Argemone capitulo longiore Obſ. Lob. B. pin. ♀

* Argemone capitulo Longiori glabro. ♃ ♀

Ari-

Ariſarum anguſtifolium Dod. anguſtifolium Dioſcoridis fortè B. pin. ⊙

Ariſtolochia rotunda Dod. rotunda flore ex purpurâ nigro B. pin. ⊙
Ariſtolochia longa Dod. longa vera B. pin. ⊙
Ariſtolochia clematitis recta B. pin. Clematitis vulgaris Cluſ. ⊙
Ariſtolochia, Piſtolochia dicta B. pin. Piſtolochia Cluſ. ⊙
Ariſtolochia, Piſtolochia Cretica B. pin. Piſtolochia altera ſemper virens Cluſ. app. prim. ⊙

Artemiſia Matth. ed. B. vulgaris major B. pin. ⊙
* Artemiſia variegatis foliis. ♃ ⊙

Arum venis albis B. pin. ⊙
Arum venis albis lituris nigris ſignatum. ⊙
Arum Fuchſ. vulgare non maculatum B. pin. ⊙
Arum maculatum maculis nigris B. pin. maculatum Tab. ⊙
Arum maculatum maculis candidis B. pin. Caſt. deſcr. ⊙

Arum indicum amplo flore. ♃ ⊙
Arum Ægyptium minus Veſlingii Obſ. in proſp. Alp. ⊙
Arum maximum Ægyptiacum, quod vulgò Colocaſia B. pin. Colocaſſia Ægyptia florida Veſlingii Obſ. in proſp. Alp. ⊙

Arundo vallatoria .ſ. phragmites Obſ. & Icon. Lob. vulgaris .ſ. phragmites Dioſcoridis B. pin. ⊙
Arundo donax .ſ. Cypria Dod. ſativa B. pin. ⊙
Arundo Indica variegata .ſ. Laconica Theophraſti Cornuti ⊙
Arundo Indica florida, Cannacorus quorundam. Obſ. & Icon. Lob. Indica latifolia B. pin. Canna Indica rubra hort. Eyſt. ♀
Arundo Indica latifolia flore luteo punctato B. pin. Canna Indica lutea punctata hort. Eyſt. ♀

Aſarum Dod. B. pin. ⊙
Aſarum Canadenſe Cornuti. ⊙

Aſclepias Fuchſ. albo flore B. pin. ⊙
Aſclepias anguſtifolia flore flaveſcente. ⊙

Aſclepias nigro flore B. pin. altera flore nigro Obſ. Lob. ☉

Aſcyrum Matth. ed. B. Dod. Lob. ☉

Aſparagus hortenſis Dod. ſativa B. pin. ☉
Aſparagus marinus Cluſ. maritimus craſſiore folio B. pin. ☉
Aſparagus foliis acutis B. pin. Corruda prior Cluſ. ☉

Aſperula odorata flore albo Dod. Aſperula .ſ. Rubeola montana odora B. pin. ☉
Aſperula cœrulea Dod. cœrulea arvenſis B. pin. ♀
* Aſperula verticillata Latifolia C. B. P. cruciata muralis Col. ♀

Aſphodelus albus ramoſus mas B. pin. major albus hort. Eyſt. ☉
Aſphodelus albus non ramoſus B. pin. Haſtula regia .ſ. Aſphodelus Matt. ed. B. ☉
Aſphodelus luteus Dod. luteus & flore & radice B. pin ☉
Aſphodelus foliis fiſtuloſis B. pin. minor Cluſ. Phalangium Cretæ Salonenſis Ad. Lob. ♀

Aſplenium Dod. Ceterach officinarum B. pin. ☉

Aſter luteus foliolis ad florem rigidis B. pin. Aſter primus Cluſ. ♀
Aſter cernuus Col. atticus foliis circa florem mollibus B. pin. ♀
Aſter montanus luteus ſalicis glabro folio B. pin. Auſtriacus ſecundus Cluſ. ☉
Aſter conyzoides odoratus luteus hort. Eyſt. ☉
Aſter conyzoides Monſpel. ♃ ♀
Aſter atticus ſupinus Dod. luteus ſupinus B. pin. ♀
Aſter luteus alatus Cornuti. ☉
Aſter atticus Dod. atticus cœruleus vulgaris B. pin. ☉
Aſter præcox flore cœruleo majore. ♃ ☉
Aſter elatior flore cœruleo minore. ♃ ☉
Aſter montanus cœruleus magno flore foliis oblongis B. pin. Alpinus cœruleo flore .ſ. ſeptimus Cluſ. ☉
Aſter minor Narbonenſium Tripolii flore, Linariæ folio Lob. Tripolii flore B. pin. ☉
Aſter flore Tripolii latifolius. ♃ ☉

Aſteriſcus

Aſteriſcus latifolius autumnalis Cornuti. ☉
Aſter ſerotinus. Canadenſis anguſtifolius flo. cœruleo. ♃ ☉
Aſter flore Bellidis ſylveſtris minoris. ♃ ☉
Aſter Canadenſis annuus. ♃ ♀ vide Senecio. ♀

Aſtragalus Bæticus Cluſ. Bæticus lanuginoſus radice ampliſſima. B. pin. ☉
Aſtragalus Monſpelienſis Cluſ. ♀
Aſtragalus ſylvaticus Thal. ſylvaticus foliis oblongis glabris B. pin. ☉
* Aſtragalus Monſpelienſium J. B. ☉

Atractylis Dod. lutea B. pin. ♀

Atriplex ſativa alba Icon. Lob. hortenſis alba ſ. pallidè virens B. pin. ♀
Atriplex hortenſis rubra B. pin. rubra Tab. ♀
Atriplex ſylveſtris vulgatior ſinuata Lob. ſylveſtris folio ſinuato candicante B. pin. ♀
Atriplex ſylveſtris latifolia B. pin. Pes anſerinus Dod. ♀
Atriplex ſylveſtris latifolia acutiore folio

B. pin. Chenopodium, Pes anſerinus ſecundus Tab. ☿

Atriplex ſylveſtris Mori fructu major, ſpinachia *B*aptiſt. ferarii. ☿

Atriplex ſylveſtris fragifera media, ſyl. baccifera Cluſ. hiſt. ☿

Atriplex ſylveſtris fragifera minor lappulas habens B. prodr. ☿

Atriplex fœtida B. pin. vulvaria Tab. ☿

Atriplex ſylveſtris Polygoni, aut helxines foliis Obſ. & Icon. Lob. anguſto oblongo folio B. pin. ☿

Atriplex folio haſtato .ſ. deltoïde. ♃ ☿

Atriplex marina Dod. maritima laciniata B. pin. ☿

* Atriplex procumbens folio ſinuato craſſo. ♃ ☿

* Atriplex halimi folio C. B. P. halimoedes Lob. Obſ. ☿

* Atriplex maritima ſinuato folio C. B. prod. ☿

Avena Dod. vulgaris .ſ. alba B. pin. ☿

Avena nigra B. pin. ☿

Azedarach Dod.

B

BALSAMINA cucumerina punicea Obſ. & Icon. Lob. rotundifolia repens .ſ. mas B. pin. ☿

Balſamina fœmina perſicifolia Obſ. & Icon. Lob. fœmina B. pin. ☿

* Balſamina lutea .ſ. Noli me tangere C. B. P. Impatiens herba Dod. ☿

Barba capræ Fuchſ. floribus oblongis B. pin. ⊙

Barba capræ floribus compactis B. pin. ulmaria Cluſ. ⊙

Barba Jovis hiſt. Lugd. B. pin.

Bellis major Dod. ſylveſtris caule folioſo major B. pin. ⊙

Bellis Alpina major rigido folio B. prodr. ♃ ⊙

* Bellis latioribus ſerratis foliis, radice repente. ♃ ⊙

Bellis ramoſa umbellifera Cornuti. ☿

Bellis media hiſt. Lugd. ſylveſtris media caule

caule carens B. pin. ☉
Bellis minor ſylveſtris Fuchſii hiſt. Ludg. Sylveſtris minor B. pin. ☉
* Bellis pratenſis duplici florum ſerie. ♃ ☉
Bellis hortenſis multiplici flore Obſ. Lob. ☉
Bellis minor flore multiplici albo hort. Eyſt. ☉
* Bellis hortenſis flo. herbaceo. ♃ ☉
* Bellis hortenſis flo. prolifero. ♃ ☉
Bellis cœrulea caule folioſo B. pin. cœrulea .ſ. Globularia Monſpeliaca Icon. Lob. ☉
Bellis cœrulea montana fruteſcens B. pin. Scabioſa decima .ſ. repens Cluſ. ☉
Bellis lutea foliis profundè inciſis major B. pin. Chryſanthemum ſegetum Obſ. & Icon. Lob. ♀
* Bellis tanaceti fol. glabris. ☉ Vide tanacetum inodorum.
* Bellis tanaceti fol. hirſutis. ☉

Berberis dumetorum B. pin. vulgaris Cluſ.

* Beta marina ſpontanea Lob. Obſ. ♀
Beta alba Matth. ed B. alba vel palleſcens quæ cicla officinarum B. pin. ♀

Beta

Beta rubra vulgatior Obſ. & Icon. Lob. rubra vulgaris B. pin. ♀
Beta rubra radice Rapæ B. pin. rubra Romana Dod. ♀
Beta rubra radice Rapæ coſtâ & foliis latioribus. ♀
Beta ſicula coſtâ amplâ aureâ. ♃ ♀
Beta Cretica ſemine aculeato B. prodr. ♀

Betonica Dod. purpurea B. pin. ☉
Betonica flore albo Matth. ed. B. alba B. pin. ☉
Betonica major Danica Park. ☉
* Betonicæ folio alopecuros quorundam J. B. ☉

Betula Dod. B. pin.

Biſtorta Dod. major radice minus intorta B. pin. ☉
Biſtorta major radice magis intorta B. pin. Serpentaria mas, ſive Biſtorta Fuchſ. ☉
* Biſtorta minor noſtras Park. ☉

Blattaria Fuchſ. Matth. ed. B. lutea folio longo laciniato B. pin. ♀
Blattaria alba, Verbaſculum Tab. alba B. pin. ♀

* Blattaria ſolio verbaſci incano, amplo luteo flore. ♃ ☿
* Blattaria perennis flo. cœruleo. ♃ ☉
* Blattaria perennis flo. carneo. ♃ ☉
* Blattaria perennis flo. Gilvo. ♃ ☉

Blitum album Matth. ed. B. album majus B. pin. ☿
Blitum minus Dod. album minus B. pin. ☿
Blitum polyſpermon Tab. B. pin. ☿
Blitum rubens Dod. rubrum majus B. pin. ☿
Blitum rubrum ſupinum Icon. Lob. rubrum minus B. pin. ☿

Botrys Dod. ambroſioïdes vulgaris B. pin. ☿
Botrys ambroſiïdes Mexiocana B. pin. app. deſcr. ☿
Botrys chamædryoïdes B. pin. Chamædrys laciniatis foliis Obſ. & Icon. Lob. ☿

Braſſica capitata albida Dod. capitata alba B. pin. ☿
Braſſica Sabauda Hyberna Icon. Lob. alba capite oblongo non penitùs clauſo B. pin. ☿
Braſſica capitata polycephalos hiſt. Lugd.

ex

ex capitibus pluribus conglobata B. pin. ♀

Braſſica rubra capitata Dod. .B pin. ♀

Braſſica gongylodes Matth. ed. b. ejuſdem pin. ♀

Braſſica dicta, Napobraſſica B. pin. ♀

Braſſica cauliflora Dod. B. pin. ♀

Braſſica vulgaris ſativa Dod. alba vel viridis B. pin. ♀

Braſſica rubra B. pin. Cumana .ſ. rubra Dod. Gall. ed. ♀

Braſſica criſpa Matth. ed. B. alba criſpa B. pin. ♀

Braſſica aſparagodes Dalechampii hiſt. Lugd. aſparagodes criſpa *B.* pin. ♀

Braſſica fimbriata Obſ. & Icon. Lob. *B.* pin. ♀

Braſſica Anglica minima Ad. Lob. fimbriata pumila *B.* pin. ♀

Braſſica Seliniſia latifolia Tab. Lato Apii folio *B.* pin. ♀

Braſſica ſelinoides Dod. anguſto Apii folio *B.* pin. ♀

Braſſica ſylveſtris Dod. arvenſis *B.* pin. ♀

Braſſica marina ſylveſtris monoſpermos Penæ hiſt. Lugd. maritima *B.* pin. ⊙

Braſſica

Braſſica ſylveſtris perfoliata Dod. Campeſtris perfoliata flore albo B. pin. ☿

Braſſica ſylveſtris foliis circa radicem cichoraceis B. pin. Sinapi album hiſt. Lugd. ⊙

Braſſica ſylveſtris foliis integris, & hiſpidis B. pin. Turritis Icon. Lob. ⊙

* Braſſica ſyl. ramoſa tota pene Glabra B. pin. Turritis maj. plateau Cluſ. ⊙

Brunella major folio non diſſecto b. pin. Brunella Dod. Eadem flore albo. ⊙

Brunella folio laciniato B. pin. Symphytum petræum Obſ. & Icon. Lob. Eadem flore albo, & roſeo. ⊙

Brunella hyſſopyfolia B. phytop. ⊙

Brunella cœrulea magno flore hort. Eyſt. B. pin. ⊙

* Brunella Alpina folio glabro, anguſto integro. ♃ ⊙

Bryonia alba Dod. aſpera .ſ. alba baccis rubris B. pin. ⊙

Bryonia ſylveſtris, Sigillum Mariæ Ad. Lob. lævis .ſ. nigra racemoſa B. pin. ⊙

Bugloſſum latifolium, Borrago B. pin. Borrago

Borrago Dod. Idem flore albo. ♀

Bugloſſum ſempervirens Ad. & Icon. Lob. latifolium ſempervirens B. pin. ☉

Bugloſſum ſempervirens minus. Symphytum pumilum repens Borraginis facie Obſ. & Icon. Lob. ☉

Bugloſſum anguſtifolium majus B. pin. anguſtifolium Obſ. & Icon. Lob. ♀

Bugloſſum minus ſativum Dodonæi hiſt. Lugd. anguſtifolium minus. B. pin. ♀

Bugloſſum ſylveſtre minus B. pin. minus ſylveſtre Dodonæi hiſt. Lugd. ♀

Bugloſſum ſylveſtre caulibus procumbentibus B. pin. Alyſſum Germanicum echioïdes Obſ. & Icon. Lob. ♀

* Bugloſſum Luſitanicum bullatis foliis. ♃ ♀

* Bugloſſum procumbens annuum pullo minimo flo. ♃ ♀

* Bugloſſum luteum minimum nobis, Echioïdes lutea Apula minima campeſtris. Col. ♀

Bulbocaſtanum majus folio Apii B. pin. Bulbocaſtanum Dod. ☉

Bulbus eriophorus Cluſ. eriophorus Orientalis B. pin. ☉

Buphtalmum lanuginoſum foliis millefolii B. pin. Chryſanthemum valentinum Cluſ. ☉

Bupleurum anguſtifolium Dod. folio ſubrotundo .ſ. vulgatiſſimum B. pin. ☉
Bupleuron anguſtiſſimo folio B. pin. minimum Col. ♀

Burſa paſtoria Dod. paſtoris major folio ſinuato B. pin.
Burſa paſtoria minor Dod. ♀
Burſa paſtoris major loculo oblongo B. pin. Draba minima muralis diſcoïdes Col. ♀
Burſa paſtoris minor loculo oblongo B. pin. Paronychia alſine folia Obſ. & Icon. Lob. ♀
Burſæ paſtoris ſimilis ſiliquoſa major B. pin. Aizoon telephium Dalechampii hiſt. Lugd. ♀

Buxus Fuchſ. arboreſcens B. pin.
Buxus humilis Dod. deſcr. foliis rotundioribus B. pin.
Buxus foliis aureis. ♃

C

*CACALIA tomentosa B. prod. ⊙
* Cacalia glabro folio Clus. ⊙
* Cacalia foliis crassis hirsutis B. pin. Ghini Lugd. ⊙

* Cachris forulaceis foliis sem. fungoso lævi. ♃ ⊙
* Cachris Peucedani folio semine fungoso sulcato lævi. ♃ ⊙
* Cachris Peucedani folio semine fungoso sulcato aspero. ♃ ⊙

Cæpa capitata Matth. ed. B. vulgaris B. pin.
Cæpa sectilis Matth. ed. B. ⊙
Cæpa Ascalonica Matth. ed. B. sterilis B. pin. ⊙

* Calceolus Mariæ luteus C. B. Pin. desc. ⊙
* Calceolus Mariæ albus C. B. P. desc. ⊙

Calamintha montana vulgaris Obſ. & Icon. Lob. vulgaris, vel officinarum Germaniæ B. pin. ☉

Calamintha montana præſtantior hort. Eyſt. magno flore B. pin. ☉

Calamintha vulgaris exiguo flore B. prodr. ☉

Calamintha arvenſis verticillata B. pin. aquatica Matthioli hiſt. Lugd. ☉

* Calamintha altera gravi odore Pulegii Lob. adv. odore Pulegii ſ. nepeta C. B. P. ☉

Caltha vulgaris B. pin. Calendula Dod.
Eadem flore citrino. ♀

Caltha polyanthos major B. pin. Calendula lutea flore pleno hort. Eyſt.
Eadem quæ
Calendula ſativa polyanthos melina Tab. ♀

Caltha arvenſis B. pin. Calendula arvenſis Tab. ♀

Caltha paluſtris Dod flore ſimplici B. pin. ☉

Caltha paluſtris flore pleno Cluſ. ☉

Campanula minor rotundifolia Obſ. & Icon. Lob. minor rotundifolia vulgaris B.

Digitized by Google

B. pin. Eadem flore albo. ☉
Campanula minor rotundifolia Alpina B. pin Cluſ. deſcr. ☉
Campanula maxima foliis latiſſimis B. pin. ☉
Campanula hortenſis Rapunculi radice B. pin. Trachelium in maritimis Ad. Lob. ☉
Campanula hortenſis folio & flore oblongo B. pin. Viola mariana Dod. Eadem flore ſubalbido. ♀
Campanula vulgatior foliis Urticæ B. pin. Cervicaria major Dod. Eadem duplici florum ſerie, & flore albo. ☉
Campanula foliis Echii B. pin. Cervicaria major tenuifolia Thal. Echium montanum Dal. hiſt. Lugd. ♀
Campanula pratenſis flore conglomerato B. pin. Trachelium minus hiſt. Lugd. Eadem flore albo. ☉

Champhorata hirſuta B. pin. Monſpeliaca Icon. Lob. ☉
Camphoratæ congener B. pin. Anthyllis altera Italorum Obſ. & Icon. Lob. ♀

Cannabis mas hiſt. Lugd. ſativa B. pin. ♀
Cannabis fœmina hiſt. Lugd. ♀

Cannabis lutea ſterilis Contareni proſp. Alp. exot. ☉

Capparis retuſo folio Obſ. Lob. Spinoſa fructu minore folio rotundo B. pin. ☉
Capparis fabago Dod. portulacæ folio B. pin. ☉

Carduus pratenſis Tragi, Acanthii facie Ad. Lob. ☉
Carduus bulboſus Monſpel. Ad. & Icon. Lob. pratenſis Aſphodeli radice latifolius B. pin. ☉
Carduus vinearum repens folio Sonchi B. pin. Ceanothos Theophraſti Col.
Idem flore albo. ☉
Carduus mollis foliis Helenii B. pin. mollis Camerarii Cluſ. ☉
Carduus albis maculis notatus vulgaris B. pin. Leucographus Dod. ♀
* Carduus Mariæ hirſutus cap. minore, ♃ ♀
* Carduus Mariæ non maculatus. ♃ ♀
Carduus albis maculis notatus exoticus B. pin. lacteus Syriacus Cam.
Idem flore albo. ♀

Carduus

Carduus albis maculis notatus exoticus minor Gataclites J. B. ♀

Carduus ſphærocephalus Dod. ſphærocephalus latifolius vulgaris B. pin. ☉

Carduus ſphærocephalus cœruleus minor B. pin. Crocodilium Monſpelienſium Dal. hiſt. Lugd. ☉

Carduus ſphærocephalus acutus Dod. Sphærocephalus capitulo longis ſpinis armato B. pin. ☉

* Carduus ſpærocephalus annuus Bœticus minor. ♃ ♀

Carduus eriocephalus Dod. capite rotundo tomentoſo B. pin. ♀

Carduus aſininus capite maj. nutante. Onopyxus tertius Dalech. hiſt. Lugd. ♀

Carduus aſininus capite maj. nutante flore albo. Acanthium ſylveſtre flore albo hort. Eyſt. ♀

Carduus aſininus capitulis parvis. Sylveſtris primus Dod. ♀

Carduus ſpinoſiſſimus vulgaris, Polyacantha Theophraſti Obſ. & Icon. Lob. Spinoſiſſimus anguſtifolius vulgaris B. pin. ☉

Carduus Polyacanthæ æmulus capite nutante. ♃ ♀

Carduus lanceolatus Tab. lanceatus latifolius B. pin. Idem flore albo. ♀
Carduus ſtellatus Dod. ſtellatus foliis papaveris erratici B. pin. Idem flore albo ♀
Carduus ſtellatus luteus foliis cyani B. pin. Spina ſolſtitialis Dod. ♀
Carduus ſolſtitialis capitulis minus ſpinoſis. Spina ſolſtitialis mitior Apula Col. ♀
* Carduus Leucoii foliis, ſubtus incanis. ♃ ♀

Carlina herbariorum Obſ. Lob. acaulos magno flore B. pin. ☉
Carlina cauleſcens magno flore B. pin. elatior, Chamæleon albus vulgaris Cluſ. ☉
Carlina acaulos minore purpureo flore B. pin. Carduus acaulis Septentrionalium Obſ. & Icon. Lob. ☉

Carpinus Matth. ed. B. Dod.

Carum Dod. Cuminum pratenſe, Carvi officinarum B. pin. ♀

Caryophyllata Dod. vulgaris B. pin. ☉
Caryophyllata vulgaris majore flore B. pin. Sylveſtris Fuchſ. ☉ Cary-

Caryophyllata montana prima Cluſ. aquatica nutante flore B. pin. ⊙
Caryophyllata montana minor flore nutante. ⊙
Caryophyllata major rotundifolia Obſ. & Icon. Lob. alpina lutea B. pin. ⊙
Caryophyllata alpina omnium minima Pon. deſcr. mont. Bald. alpina minor B. pin. ⊙

Caryophyllus maximus ruber B. pin. maximus plenus flore rubro hort. Eyſt. ⊙
Caryophyllus multiplex flore albo hort. Eyſt. ⊙
Caryophyllus multiplex maximus variegatus hort. Eyſt. ⊙
Caryophyllus hortenſis ſimplex flore majore B. pin. Caryophylleus flos ſimplex Dod. ⊙
Caryophyllus arboreſcens Creticus B. prodr. Sylveſtris arboreus Proſp. Alp. de exot. ⊙
Caryophyllus ſimplex flore minore pallidè rubente B. pin. Sylveſtris tertius flore pallido ſuaverubente Icon. Lob. ⊙
Caryophyllus ſimplex ſupinus latifolius B. pin. minimus pulchellus ſupinus maculis aureis, argenteiſve aſperſus Icon. Lob. ⊙

Caryophyllus

Caryophyllus barbatus flore multiplici B. pin. armerius pleno flore Cluſ. Idem flore albo. ☉

Caryophyllus barbatus hortenſis latifolius B. pin. Viola barbata latifolia hiſt. Lugd. Idem qui

Flos armerius ruber hort. Eyſt. ☉

Flos armerius albus ejuſdem.

Flos armerius variegatus ejuſdem.

Caryophyllus barbatus ſylveſtris B. pin. Viola barbata anguſtifolia hiſt. Lugd. ♃

Caryophyllus ſylveſtris prolifer B. pin. armeria prolifera Icon. Lob. ♃

Caryophyllus montanus primus Tab. ſylveſtris vulgaris latifolius B. pin. ☉

Caryophyllus flore tenuiſſimè diſſecto B. pin. Caryophylleus minor Dod. ☉

Idem qui

Caryophyllus plumarius albus Tab.

Caryophyllus omnium humillimus. ♃ ☉

Caryophyllus pratenſis flore laciniato ſimplici .ſ. flos cuculi B. pin. Odontitis Plinii ſimplici flore Cluſ. ☉

Idem flore albo.

Caryophyllus pratenſis flore pleno B. pin. Odontitis pleno flore Cluſ. ☉

Caryophyllus

Caryophyllus ſylveſtris alter flore laciniato odoratiſſimo B. pin. Sylveſtris ſextus Cluſ. ☉

Caryophyllus holoſtius Icon. Lob. arvenſis hirſutus flore majore B. pin. ☉

Caryophyllus holoſtius tomentoſus anguſtifolius B. prodr. ☉

Caryophyllus arvenſis umbellatus folio glabro B. pin. Alſine verna Dal. hiſt. Lugd. ♀

Caryophyllus arvenſis glaber flore majore B. pin. Gramen leucanthemum Dod. ☉

Caryophyllus arvenſis glaber flore minore B. pin. Gramen floridum minus Tab. ☉

Caryophyllus holoſtius Alpinus gramineus B. prodr. ☉

Caryophyllus montanus major flore globoſo B. pin. Statice hiſt. Lugd. ☉

Caryophyllus montanus minor flore globoſo B. pin. Gramen polyanthemum minus Dod. ☉

Caryophyllus minimus flore globoſo. ♃ ☉

Caryophyllus minimus muralis B. pin. Tunica minima hiſt. Lugd. ♀

Caſtanea Dod.
Caſtanea equina Cluſ. folio multifido B pin.

Caucalis maxima ſphondylii aculeato ſemine B. pin. major Cluſ. vide Tordylium. ☿
Caucalis arvenſis echinata latifolia B. pin. echinophora quarta major πλατύφυλλον purpurea Col. Eadem flore albo. ☿
Caucalis arvenſis echinata magno flore B. pin. albo flore Cluſ. ☿
Caucalis arvenſis echinata parvo flore & fructu B. pin. echinophora tertia λεπτόφυλλον purpurea Col. ☿
Caucalis Monſpeliaca echinato magno fructu B. pin. Lappa boaria Plinii hiſt. Lugd. ☿
Caucalis nodoſo echinato ſemine B. prodr. ☿
Caucalis folio Peucedani B. pin. Anglica Tab. ⊙

Cedrus folio Cupreſſi major fructu flaveſcente B. pin. Oxycedrus Lycia Dod.
* Cedrus folio cupreſſi minor C. B. P.

Centaurium majus folio in lacinias plures diviſo B. pin. magnum Dod. ⊙

Centauri-

Digitized by Google

Centaurium foliis Cinaræ Cornuti. ☉
* Centaurium Rha capitatum fol. Enulæ Lugd. ☉
Centaurium majus luteum Cornuti. Alpinum luteum B. prodr. ☉
Centaurium minus Dod B. pin. Idem quod Centaurium minus album Tab. ♀
Centaurium luteum perfoliatum B. pin. parvum flavo flore Cluſ. ♀
Centaurium luteum puſillum B. pin. luteum novum Col. ♀
Centaurium minimum purp. ♃ ♀

Cepæa Cluſ. ♀

Ceraſus fructu acidulo Dod. deſcr. Ceraſus auſtera Matth. Ed. B.
Ceraſa craſſa carne dura B. pin. Duracina hiſt. Lugd. deſcr.
Ceraſus Dod. major, ac ſylveſtris fructu ſubdulci nigro colore inficiente B. pin.
Ceraſus hortenſis flore roſeo B. pin. multiflora Tab.
Ceraſus folio laurino B. pin. Lauroceraſus Cluſ.
Ceraſus pumila B. pin. Chamæceraſus Dod.

Ceraſus racemoſa ſylveſtris fructu non edu-
li B. pin. Ceraſus avium racemoſa Icon.
Lob.
Ceraſo affinis B. pin. Mahaleb Matthioli
hiſt. Lugd.

Cerinthe quorundam major verſicolore
flore Cluſ. Cerinthe, ſive Cynogloſſum
montanum majus B. pin. ☿
Cerinthe major quorundam flore ex rubro
purpuraſcente Cluſ. deſcr. flore ex ru-
bro purpuraſcente B. pin. ☿
Cerinthe quorundam major flavo flore
Cluſ. flore flavo aſperior B. pin. ☿

Chærophyllum ſativum B. pin. Cherefoli-
um Dod. ☿
Chærophyllum ſylveſtre B. pin. Anthriſcus
Plinii hiſt. Lugd. ☿

Chamæceraſus alpina fructu rubro gemino
duobus punctis notato B. pin. Xyloſte-
um alterum Dod.
Chamæceraſus dumetorum fructu gemino
rubro B. pin. Xyloſteum Dod.

Chamæciſtus

Chamæciſtus vulgaris flore luteo B. pin. primus Cluſ. ☉

Chamæciſtus foliis minoribus, & incanis B. pin. quintus Cluſ. quoad figuram, quartus deſcr. ☉

Chamæciſtus foliis Myrti minoris incanis B. pin. tertius Cluſ. ☉

Chamæciſtus anguſtifolia B. pin. ſextus Cluſ. ☉

* Chamæciſtus Halimi folio. ☉ ♃

* Chamæciſtus parvo flo. luteo J. B. an Serpilli folio nigricante et hirſuto flo. aureo odorato eidem. ☉

Chamæciſtus annuus. ♃ ☿

Chamædrys repens minor Dod. B. pin. Eadem flore albo. ☉

Chamædrys alpina ciſti flore B. pin. montana fruteſcens durior Ad. Lob. ☉

Chamædrys alpina minima hirſuta B. prodr. ☉

Chamædrys ſpuria major anguſtifolia B. pin. Veronica ſupina facie Teucrii pratenſis Icon. Lob. ☉

Chamædrys ſpuria minor rotundifolia B. pin. Teucrium officinarum Obſ. Lob. ☉

Chamægeniſta

Chamægeniſta foliis Geniſtæ vulgaris B. pin. prima Cluſ. ⊙
Chamægeniſta ſagittalis B. pin. ſecunda Cluſ. ⊙

Chamæiris variegata B. pin. cœruleo, & violaceo colore hort. Eyſt. ⊙
Chamæiris minor flore purpureo B. pin. Iris minor Dod. ⊙
Chamæiris flore pallidè luteo B. pin. ⊙
Chamæiris flore obſoleto pallido B. pin. latifolia minor ſexta Cluſ. ⊙

* Chamæleon niger cortuſi Dod. niger Lob. Obſ. ⊙

Chamæmelum vulgare Dod. vulgare leucanthemum Dioſcoridis B. pin. ⊙
Chamæmelum nobile .ſ. odoratius B. pin. odoratum Dod. ⊙
Chamæmelum nobile flore multiplici B. pin. Romanum flore multiplici hort. Eyſt. ⊙
Chamæmelum nudum capitulo aphyllo B. pin. Anthemis chryſanthemos Penæ & Lobellii hiſt. Lugd. ⊙
Chamæmelum inodorum B. pin. Cotula non

non fœtida Dod. Gall. ed. Idem flore pleno. ☉

Chamæmelum fœtidum B. pin. Cotula fœtida Dodonæi hiſt. Lugd. ♀

Chamæmeſpilus Geſneri Cluſ. Epimelis hiſt. Lugd.

Chamæpitys moſchata foliis ſerratis B. pin. Spuria prior, ſive Anthyllis altera Dod. ☉

Chamæpitys lutea vulgaris .ſ. folio trifido B. pin. Ajuga .ſ. Chamæpitys mas Dioſcoridis Obſ. & Icon. Lob. ♀

Chamæſyce Dod. B. pin. ♀

Chamelæa tricoccos hiſt. Lugd. B. pin. ☉

Chelidonium majus vulgare B. pin. majus Dod. ☉

Chelidonium majus foliis quernis B. pin. majus laciniato flore Cluſ. ☉

Chelidonium majus foliis tenuiſſimè diſſectis. ♃ ☉

Chelidonium maximum Canadenſe ἄκαυλον Cornuti deſcr. ☉

Chelidonium minus hiſt. Lugd. Cheledonia rotundifolia minor B, pin. ☉
* Chelidonium minus flo. pleno Park. ☉

Chondrilla juncea viſcoſa arvenſis B. pin. viminea viſcoſa Penæ hiſt. Lugd. ☉
Chondrilla viſcoſa humilis B. prodr. ♀
Chondrilla verrucaria foliis intubaceis viridibus B. pin. Cichorium verrucatum, Zacintha Cluſ. ♀
Chondrilla cœrulea altera cichorii ſylveſtris foliis B. pin. cœrulea cichorii ſylveſtris laciniosis foliis Obſ. Lob. ☉
Chondrilla foliis laciniatis ſerratis purpuraſcente flore B. pin. rara purpurea Lobelii hiſt. Lugd. ♀
Chondrilla cœrulea cyani capitulo B. pin. Seſamoïdes parvum Matth. ed. B. ☉

Chryſanthemum flore partim candido partim luteo B. pin. Creticum mixtum hort. Eyſt. ♀
Chryſanthemum majus folio profundiùs laciniato magno flore B. pin. luteum hort. Eyſt. ♀
Chryſanthemum alpinum incanum foliis lacini-

laciniatis B. pin. alpinum primum Cluſ. ☉
Chryſanthemum tenuifolium Hiſpanicum. ♃ ☉
Chryſanthemum alpinum foliis Abrotani multifidis B. pin. Achillea montana Penæ hiſt. Lugd. ☉
* Chryſanthemum Myconi Lugd. Bellis lutea fol. ſubrotundis C. B. P. ♀
* Chryſanthemum tenuifolium Eyſt. Bupthalmum Matthioli Lob. ♀

Cicer arietinum Dod. ſativum B. pin. Idem flore & fructu rubro B. pin. deſcr. ♀
* Cicer Arietinum fructu nigro. ♀
Cicer ſylveſtre herbariorum Obſ. Lob. ſylveſtre foliis oblongis hiſpidis majus B. pin. ♀

Cichorium latioris folii Dod. ſativum B. pin. ♀
Cichorium ſylveſtre Matth. ed. B. ſylveſtre ſ. officinarum B. pin. Idem flore albo & purpureo. ♀
Cichorium montanum anguſtifolium hirſutie aſperum B. pin. Hieracium intybaceum alperum Tab. ☉

Cicuta Dod. major B. pin. ♀
Cicuta minor Petroſelino ſimilis B. pin. Cicutaria fatua Ad. & Icon. Lob. ♀
* Cicutaria latifolia fœtidiſſima Ger. Seſeli Peleponenſe Lugd. ⊙
Cicutaria paluſtris tenuifolia B. pin. Phellandryum Dod. ⊙

Cinara hortenſis foliis non aculeatis B. pin. non aculeata Dodonæi hiſt. Lugd. ♀
Cinara hortenſis non aculeata capite ſubrubente. ♀
Cinara hortenſis aculeata B. pin. Scolymus Dioſcoridis, Cactos Theophraſti Cluſ. ♀
Cinara maxima Anglica Icon. Lob. Scolymus ſ. Cinara maxima hort. Eyſt. ♀
Cinara ſpinoſa cujus pediculi eſitantur B. pin. Carduus aculeatus Matth. ed B. ♀

Ciriium maximum Aſphodeli radice B. pin. maximum montanum incano folio bulboſa radice quintum Cluſ. deſcr. ⊙
Cirſium majus ſingulari capitulo magno, vel incanum variè diſſectum B. pin, Anglicum Obſ. & Icon. Lob. ⊙

Ciſtus

Ciſtus mas folio oblongo incano B.pin. mas primus Cluſ. ☉

Ciſtus fœmina folio ſalviæ B. pin. fœmina Cluſ. ☉

Ciſtus folio ſalicis B.pin. humilis annuus folio ſalicis humilis Obſ. Lob. ☿

Ciſtus annuus flore guttato Ponæ deſcr. mont. Bald. flore pallido punicante macula inſignito B. pin. ☿

Idem flore non maculoſo.

Ciſtus ledon foliis laurinis B. pin. Ciſti ledi primi altera ſpecies Cluſ. deſcr. ☉

Ciſtus ledon quintum Cluſ. ledon Narbonenſe Tab. ☉

* Ciſtus mas folio rotundo hirſut. C.B.P. folio rotundiore Matth. Lugd. ☉

* Ciſtus Roſmarini folio. ☉

Clematitis ſylveſtris latifolia B. pin. Atragene Theophraſti Cluſ. ☉

Clematis .ſ. flammula repens B. pin. Flammula Dod. ☉

Clematis .ſ. flammula recta B. pin. Flammula altera ſurrecta Obſ. Lob. ☉

Clematitis cœrulea erecta B. pin. cœrulea pannonica hort. Eyſt. ☉

Clematitis cœrulea vel purpurea repens B. pin. peregrina purpurea ſimplici flore hort. Eyſt. ☉

Clematitis cœrulea flore pleno B. pin. peregrina purpurea flore pleno hort. Eyſt. ☉

* Clematis peregrina flore parvo rubro. ☉

* Clematis peregrina flore parvo albo. ☉

Clematitis trifolia flore roſeo clavato B. pin. Granadilla, Flos paſſionis hiſt. Mex. ☉

Clematis, ſeu flos paſſionis flore viridi. Murucuiæ ſpecies foliis hederæ ſcandentis Georgii Marcgravii. In hiſt. Braſ. ☉

Clematis tetraphyllos Americana. ♃ ☉

Clematis daphnoïdes minor B. pin. Clematis daphnoïdes Dod. ☉

Clematis daphnoïdes flore purpureo hort. Eyſt. ☉

Clematis daphnoïdes major Dod. B. pin. ☉

Clinopodium Matth. Cluſ. Origano ſimile B. pin. ☉

Clinopodium vulgare Cluſ. arvenſe Ocimi facie B. pin. ♀

* Clinopodium Auſtriacum Ponæ. ☉

Cneorum Matthioli Cluſ. Thymelææ affinis facie externa B. pin. ☉. Cne-

Digitized by Google

Cneoron album folio Oleæ argenteo molli B. pin. Dorycnium Plateau Cluſ. ☉

Cnicus vulgaris Cluſ. ſativus, ſ. Carthamum officinarum B. pin. ♀

Cnicus cœruleus aſperior B. pin. alter Cluſ. cœruleo flore. ☉

Cnicus ſylveſtris hirſutior, ſ. Carduus benedictus B. pin. Carduus benedictus Dod. ♀

Cochlearia Dod. folio ſubrotundo B. pin. ♀

* Cochlearia Armorica flore parvo ſubpurpureo. ♂ ☉

* Cochlearia Britannica Lob. Dod. folio ſinuato C. B. P. ☉

Colchicum commune B. pin. vulgare flore purpuraſcente hort. Eyſt. vulgare album ejuſdem. ☉

Colchicum commune folio lineis albis picto. ♂ ☉

Colchicum polyanthes lato hellebori albi folio B. pin. Byzantinum πλατύφυλλον Cluſ. ☉

Colchicum candidum multiflorum B. pin. Pannonicum πολύανθες albo flore Cluſ. ☉

Colchicum variegatum Cornuti. ☉

Colchicum pleno flore Cluſ. B. pin. ☉
Colchicum luteum majus B. pin. Narciſſus autumnalis major Cluſ. ☉
* Colchicum foliorum oris luteis. ♃ ☉

Colocynthis pyriformis .ſ. pepo amarus B. pin. pyriformis Obſ. Lob. Pſeudocolocynthis pyri formâ hort. Eyſt. ♀

Colutea veſicaria Matthioli hiſt. Lugd. veſicaria B. pin. ☉
Colutea ſiliquoſa .ſ. ſcorpioïdes major B. pin. Scorpioïdes prima elatior Cluſ. ☉
* Colutea ſcorpioïdes maritima glauco folio B. in prod. ☉
* Colutea humilis Buxi foliis nobis, Anonimos flore coluteæ Cluſ. Pſeudochamabuxus Eyſt. ☉

Conſolida media pratenſis cœrulea B. pin. Bugula Dod. Eadem flore albo & carneo ☉
Conſolida media foliis hirſutis. Eadem flore albo & carneo. ☉

Conſolida regalis hortenſis flore minore B. pin.

pin. Delphinium elatius ſimplici flore Cluſ. Eadem flore albo & carneo. ♀

Conſolida regalis multiplicato violaceo flore hort. Eyſt. ♀

Conſolida regalis multiplicato incarnato flore hort. Eyſt. ♀

Conſolida regalis flore pleno albo hort. Eyſt. ♀

Conſolida regalis flore pleno rubro hort. Eyſt. ♀

Conſolida regalis flore multiplici cinericeo hort. Eyſt. ♀

Conſolida regalis flore pleno variegato. ♂ ♀

Conſolida regalis arvenſis B. pin. Flos regius ſylveſtris Dod. ♀

Conſolida regalis latifolia parvo flore B. prodr. ♀

* Conſolida regalis hortenſis flore minore verſicolori. ♂ ♀

Convolvulus major albus B. pin. Smilax lævis major Dod. ⊙

Convolvulus minor arvenſis B. pin. Smilax lævis minor Dod. ⊙

* Convolvulus virginianus nobis Battatas cammotes

cammotes Hiſpanorum Cluſ. ſiſarum Peruvianum Ger. Tab. ☉

Convolvulus minor ſemine triangulo B. pin. Helxine ciſſampelos altera Atriplicis effigie Lob. ♀

Convolvulus cœruleus hederaceo anguloſo folio B. pin. aſureus, ſ. cœruleus Obſ. Lob. ♀

Convolvulus purpureus folio ſubrotundo B. pin. Indicus flore violaceo hort. Eyſt. ♀

Convolvulus peregrinus cœruleus folio oblongo B. pin. & prodr. ♀

Convolvulus minimus ſpicæfolius Obſ. Lob. Linariæ folio B. pin. ☉

Convolvulus minor argenteus radice repente. ♃ ☉

Convolvulus hederaceus perennis. ♃ ☉

Conyza major vera Ad. Lob. mas Theophraſti, major Dioſcoridis B. pin. ☉

Conyza minor vera Ad. Lob. fœmina Theophraſti, minor Dioſcoridis B. pin. ♀

Conyza major vulgaris B. pin. Baccharis Monſpelienſium Obſ. & Icon. Lob. ☉

Conyza pannonica lanuginoſa B. pin. tertia Auſtriaca Cluſ. ☉

Conyza

Digitized by Google

Conyza cœrulea aeris B. Pin. Erigeron quartum Dod. ☉

Conyza helenitis piloſa Icon. Lob. alpina piloſiſſima B. pin. ☉

Conizis affinis B. pin. Britannica Geſneri & Lugdunenſium Ad. Lob. ☉

Conyza media vulgaris Cluſ. media Aſteris flore luteo, vel tertia Dioſcoridis B. pin. ☉

Conyza minima Dod. major flore globoſo B. pin. ♀

Conyza paluſtris ſerratifolia B. pin. Lingua major Dal. hiſt. Lugd. ☉

Conyza marina hiſt. Lugd. ☉

* Corchoros Plinii Lob. Lugd. melochia .ſ. corchorus Park. ♀

Coriandrum Fuchſ. majus B. pin. ♀

* Coriandrum ſylveſtre fœtidiſſimum C. B. P. ſylveſtre Myconi Lugd. ♀

Coris cœrulea maritima B. pin. Coris quorundam Cluſ. ☉

Cornus mas hiſt. Lugd. hortenſis mas B. pin.

Cor-

Cornus fœmina B. pin. Virga ſanguinea Dod.

Coronopus Matthioli hiſt. Lugd. hortenſis B. pin. ♀

Coronopus maritima major B. pin. Plantago marina Obſ. & Icon. Lob. ⊙

Corylus Cluſ. ſativa fructu albo minore .ſ. vulgaris B. pin.

Corylus ſativa fructu oblongo rubente B. pin. domeſtica fructu oblongo rubro Cluſ. deſcr.

Corylus ſylveſtris B. pin. Avellana nux ſylveſtris Fuchſ.

Cotinus Matth. ed. B. Coggygria Cluſ. ⊙

Cotonaſter folio oblongo ſerrato B. pin. Cotonaſter forte Geſneri Cluſ. ⊙

Cotyledon Dod. major B. pin. ⊙

Cotyledon radice ſerpente. ♃ ⊙

Cotyledon media foliis oblongis ſerratis B. pin. minus hort. Eyſt. Palmaria Tab. ⊙

Cotyledon minor foliis ſubrotundis ſerratis B. prodr. ⊙ Coty-

Digitized by Google

Cotyledon ſtellata B. pin. Aizoon peregrinum Cam. ♀

Cratægus Theophraſti hiſt. Lugd. Sorbus torminalis Dod.

Crithmum ſ. fœniculum maritimum minus B. pin. marinum Dod. ⊙
Crithmum chryſanthemum Dod. maritimum flore Aſteris attici B. pin. ⊙
Crithmum ſpinoſum Dod. maritimum ſpinoſum B. pin. ⊙

Crocus ſativus Matthioli hiſt. Ludg. ſativus B. pin. ⊙
* Crocus albus ſativus C. B. P. deſcr. ⊙
Crocus vernus latifolius flore purpureo verſicolore Cluſ. vernus latifolius purpureus variegatus B. pin. ⊙
Crocus vernus latifolius flore purpureo magno B. pin. Cluſ. deſcr. ⊙
Crocus vernus latifolius albo flore Cluſ. ⊙
Crocus flore argenteo lineis violaceis ſtriato Icon. Paſſæi. ⊙
Crocus vernus latifolius flavo flore Cluſ. vernus latifolius flavus B. pin. ⊙

Crocus

Crocus vernus latifolius pallido flore Cluſ. deſcr. latifolius pallidus B. pin. ☉
Crocus vernus latifolius flavo varius B. pin. luteus lineis nigris Swert. ☉

Cruciata Dod. hirſuta B. pin. ☉

Cucumis ſativus hiſt. Lugd. ſativus vulgaris B. pin. ♀
Cucumis anguinus flexuoſuſve Obſ. Lob. flexuoſus B. pin. ♀
Cucumis ſylveſtris Dod. ſylveſtris aſininus dictus B. pin. ♀

Cucurbita lagenaria Obſ. & Icon. Lob. lagenaria flore albo folio molli B. pin. ♀
Cucurbita longior Dod. oblongo flore albo folio molli B. pin. ♀
Cucurbita capitata Tab. B. pin. ♀

Cuminum ſemine longiore B. pin. ſativum Dod. ♀
Cuminum ſylveſtre capitulis globoſis B. pin. ſylveſtre Dod. ♀

Cupreſſus Dod.

Cupreſſus

Cupreſſus fœmina Dod. deſcr.

Cyanus montanus latifolius vel Verbaſculum cyanoïdes B. pin. major Dod. ☉
Cyanus ſegetum B. pin vulgaris Obſ. & Icon. Lob. ☿
Cyanus hortenſis flore ſimplici cœruleo, albo, purpureo. ☿
Cyanus floridus odoratus Turcicus .ſ. Orientalis Park. flore albo & carneo. ☿
Cyanus repens latifolius B. pin. peramarus repens capitulis ſquaroſis Seſamoïdis parvi Ad. Lob. deſcr. ☉

Cyclamen vulgare hort. Eyſt. orbiculato folio inferne purpuraſcente B. pin. Idem flore albo. ☉
Cyclamen vernum hort. Eyſt. folio anguloſo B. pin. ☉
Cyclamen ſerotinum foliis hederaceis hort. Eyſt. hederæ folio B. pin. ☉
Cyclaminum Orientale Cornuti. ☉
* Cyclamen Perſicum odoratum. ☉

Cymbalaria Matthioli hiſt. Lugd. ☉

Cyno-

Cynogloſſum Dod. majus vulgare B. pin. ♀
Cynogloſſum ſylvaticum rubente caule B. pin. Cynogloſſa montana maxima frigidarum regionum Col. ♀
Cynogloſſum Creticum argenteo anguſto folio B. pin. Creticum primum Cluſ. ♀
* Cynogloſſum ſemper virens C. B. Prod. Cynogloſſa altera fol. virente Col. ♀
Cynogloſſum minus B. pin. Lappula ruſticorum, Elatine Tragi hiſt. Lugd. ♀

Cynoſorchis militaris major B. pin. Orchis ſtrateumatica, ſive ſtratiotes major, .ſ. militaris Obſ. & Icon. Lob. ⊙
Cynoſorchis militaris pratenſis humilior B. pin. minor flore incarnato hort. Eyſt. ⊙
Cynoſorchis latifolia ſpicâ compactâ B. pin. Cynoſorchis tertia Dod. ⊙

Cyperus rotundus inodorus Germanicus B. prodr. ⊙
Cyperus longus Dod. odoratus radice longâ .ſ. Cyperus officinarum B. pin. ⊙

Cytiſus Maranthæ Matthioli Obſ. Lob. incanus ſiliquis falcatis B. pin. ⊙

Cityſus

Cytiſus incanus Creticus B. pin. Ebenus Cretica Proſp. Alp. exot. ⊙

Cytiſus ſupinus infernè, & ſiliquis molli hirſutie pubeſcentibus B. pin. ſeptimus hort. Eyſt. ⊙

Cytiſus glabris foliis, ſubrotundis, pediculis breviſſimis B. pin. quintus rotundifolius Tab. ⊙

* Cytiſus incanus Auſtriacus folio oblongo C. B. P. 5 Cluſ. hiſt. ⊙

* Cytiſus glaber nigricans C. B. P. 4 Cluſ. hiſt. ⊙

D.

DAUCUS montanus Apii folio major B. pin. Selinoïdes ſecundus Obſ. & Icon. Lob. ⊙

Daucus Alſaticus B. prodr. ⊙

Daucus pratenſis Dal. hiſt. Lugd. pratenſis millefolii paluſtris folio B. pin. ⊙

Dens caninus purpuraſcente flore Cluſ. latiore rotundioreque folio B. pin. ⊙

Dens leonis Dod. latiore folio B. pin. ☉

Dentaria ἑπτάφυλλος Cluſ. heptaphyllos B. pin. ☉
Dentaria baccifera Cluſ. heptaphyllos baccifera B. pin. ☉
Dentaria τρίφυλλος Cluſ. triphyllos B. pin. ☉
* Dentaria monophyllos, anapodophyllon quibuſdam. ♃ ☉
* Dentariæ affinis, Echii flore, capſula rotunda. ♃ ☉

Digitalis purpurea Dod. purpureo folio aſpero B. pin. ☉ Eadem flore albo.
Digitalis major lutea, vel pallida parvo flore B. pin. quarta Dod. deſcr. ☉
* Digitalis lutea amplo flore Ponæ lutea magno flore C. B. P. ☉
Digitalis latifolia flore ferrugineo. ♃ ☉
Digitalis anguſtifolia flore ferrugineo B. pin. ferruginea Icon. Lob. ☉
Digitalis foliis maximè ſerratis. ♃ ☉
Digitalis foliis ſubrubentibus Lyſimachiæ facie. ♃. ☉

Dipſacus ſativus Dod. B. pin. ♀

Dipſacus

Dipſacus ſylveſtris Dod. ſylveſtris aut virga paſtoris major B. pin. ♀

Dipſacus ſylveſtris capitulo minore, vel virga paſtoris minor B. pin. tertius Dod. ♀

Doronicum latifolium Cluſ. radice ſcorpii B. pin. ☉

Doronicum Plantaginis folio B. pin. minus officinarum Obſ. & Icon. Lob. ☉

Doronicum Plantaginis folio alterum B. pin. Aliſma Matth. ed. B. hort. Eyſt. ☉

Doronicum maximum foliis caulem amplexantibus B. pin. ſeptimum Auſtriacum Cluſ. ☉

Doronicum foliis ſubtus lanuginoſis. ♃ ☉

Doronicum Americanum laciniato folio B. pin. app. Aconitum helianthemum Canadenſe Cornuti. ☉

* Doronicum Americanum minus fol. obſcurius virentibus. ♃ ☉

Draba umbellata, vel Draba major capitulis donata B. pin. prima vulgaris Cluſ. ☉

Draba alba ſiliquoſa repens B. pin. tertia ſucculento folio Cluſ. ☉

Draba lutea ſiliquis ſtrictiſſimis B. pin. lutea Matth. ed. B. ☉

Draba ſiliquoſa floribus violaceis. Forte Leucovium marinum alterum latifolium Obſ. Lob. ♀

Dracunculus hortenſis B. pin. Draco-herba Dod. ☉

Dracunculus pratenſis ſerrato folio B. pin. Ptarmica Fuchſii hiſt. Lugd. ☉

Dracunculus pratenſis flore pleno B. pin. Ptarmica vulgaris flore pleno Cluſ. ☉

Dracunculus polyphyllus B. pin. Dracontium majus Dod. ☉

* Dracunculus .ſ. Serpentaria Triphylla Braſſiliana B. prodr. ☉

E.

ECHIUM ſcorpioïdes arvenſe B. pin. Myoſotis hirſuta reptans Obſ. Lob. ♀

Echium ſcorpioïdes paluſtre B. pin. Myoſotis ſcorpioïdes partim repens partim erecta Icon. Lob. Idem flore albo. ☉

Echium vulgare Cluſ. B. pin. ♀

Echium procumbens floſculis atrorubentibus. ♃ ♀

Elatine

Elatine folio ſubrotundo B. pin. Veronica fœmina Fuchſ. ♀
Elatine folio acuminato in baſi auriculato flore luteo B. pin. altera Dod. ♀

* Elychryſum. ſ. ſtæchas Citrina altera Lob. Icon. Lugd. ⊙
* Elychryſum .ſ. ſtæchas Citrina Cretica amplo flore J. B. Creticum Eyſt. ♀

Ephemerum non lethale Matthioli hiſt. Lugd. ⊙

Epimedium Dod. ⊙

* Equiſetum arvenſe longioribus ſetis C. B. P. Park. ſegetale Ger.
* Equiſetum pratenſe longiſſ. ſetis C. B. P. paluſtre maj. Tab. paluſtre Ger.
* Equiſetum nudum Ger. Junceum Lugd. trag.

Erica vulgaris glabra B. pin. prior Dod. ⊙
Erica floribus ex ſummis caulibus pendulis. Eadem flore albo. ⊙

Erica major ſcoparia foliis deciduis B. pin. coris folio quarta Cluſ. ☉

Erica pumila calyculato Unedonis flore Icon. Lob. humilis cortice cinericeo Arbuti flore B. pin. Eadem flore albo ☉

Erica ſupina maritima Anglica Park. ☉

Eruca ſativa Fuchſ. latifolia alba ſativa Dioſcorid. B. pin. ♆

Eruca latifolia B. prodr. ♆

Eruca ſylveſtris Dod. ſylveſtris major lutea caule aſpero B. pin. ☉

Eruca paluſtris minor Tab. ſylveſtris minor luteo parvoque flore B. pin. ☉

Eruca vinealis parvo flore. An ſylveſtris minor lutea Burſæ paſtoris folio B. prodr. ♆

Eruca lutea latifolia .ſ. Barbarea B. pin. Barbarea Dod. ♆

Eruca Tanaceti foliis. ♃ ☉

* Eruca Monſpeliaca ſiliqua quadrata Echinata C. B. P. ſinapi Echinatum Lugd. ♆

Eruca maritima Italica ſiliquâ haſtæ cuſpidi ſimili B. pin. Cakile Serapionis Obſ. Lob. ♆

* Eruca

* Eruca Maritima Anglica Park. marina Cakile Serapionis Lob. ♀
* Eruca ſ. Barbarea perennis, ſem. craſſiore. ♃

Eryngium marinum Dod. maritimum B. pin. ☉
Eryngium campeſtre Dod. vulgare B. pin. ☉
Eryngium montanum Amethyſtinum B. pin. totum cœruleum hort. Eyſt. ☉
* Eryngium montanum Pannonicum album Cluſ. deſcr. ☉
Eryngium montanum minimum capitulo magno B. pin. Carduncellus montis Lupi Ad. Lob. ☉
Eryngium minimum mitius capitulo magno. ♃ ☉
Eryngium ex plani ſemine degener. ♃ ☉
Eryngium arvenſe foliis Serræ ſimilibus B. pin. quartum Dod. ☉

Eryſimum vulgare B. pin. Irio, ſ. Eryſimum Dod. ♀
Eryſimum latifolium majus glabrum B. pin. Irio Apulus alter lævi folio Erucæ Col. ♀
* Eryſimum anguſtifolium majus C. B. P.

Digitized by Google

Rapiſtrum ſylveſtre Irionis folio Col. ♆
* Eryſimum Polyceration vel corniculatum C. B. P. Irio alter Matth. Lugd. ♆

Euonymus Dod. vulgaris granis rubentibus B. pin.
Euonymus latifolius B. pin. primus .ſ. latifolius Cluſ.

Eupatorium vulgare Matth. ed. B. cannabinum B. pin. ☉
Eupatoria foliis Enulæ Cornuti. ☉
Eupatorium cannabinum chryſanthemum Tab. ♆
Eupatorium cannabinum Americanum flore aureo. ♃ ♆
Eupatorium veterum .ſ. Agrimonia B. pin. Græcorum .ſ. Agrimonia Dod. ☉
Eupatorium odoratum B. pin. alterum odoratum, & aromaticum Col. deſcr. ☉
* Eupatorium odoratum ſantonicum Lob. Adverſ. ☉
* Eupatorium capite nutante C. B. Chryſanthemum J. B. ♆
* Eupatorium Chryſanthemum, canadenſe præaltum, alato caule nobis. Aſter virgineus

gineus luteus alato caule Park.deſcr. ☉

Euphraſia Dod. officinarum B. pin. ♀
Euphraſia minus ramoſa flore ex cœruleo purpuraſcente hort. Eyſt. unicaulis B. pin. deſcr. ♀
* Euphraſia unicaulis ſpicata. Trixago apula unicaulis Col. ♀
Euphraſia lutea viſcata. ♃ ♀
Euphraſia pratenſis rubra B. pin. ſecunda Dodonæi, ſive Cratæogonon alterum quorundam Obſ. Lob. ♀
Eufragia major ſylveſtris purpurea Col. Italica latifolia altera B. pin. ♀
Euphraſia pratenſis lutea B. pin. Sideritis pratenſis lutea hiſt. Lugd. ♀

F.

* FABA major fructu rubro C.B.Pin. ♀
Faba Fuchſ. major recentiorum Obſ. & Icon. Lob. ♀
Faba minor .ſ. equina B. pin. Bona .ſ. phaſelus minor Dod. deſcr. ♀

Faba

Faba ſylveſtris Græcorum Obſ. & Icon. Lob. ſylveſtris fructu rotundo atro B. pin. ♀ Vide vicia.

Fagus Dod.

Ferrum equinum ſiliquâ ſingulari B. pin. equinum Obſ. & Icon. Lob. ♀

* Ferrum equinum Polyceraton Col. Equinum ſiliquâ multiplici C. B. P. ♀

Ferrum equinum Germanicum ſiliquis in ſummitate B. pin. equinum capitatum Col. ☉

Ferrum equinum Gallicum ſiliquis in ſummitate B. pin. Lotus enneaphyllos Dal. hiſt. Lugd. ☉

Ferula galbanifera Obſ. & Icon. Lob. ferulago latiore folio B. pin. ☉

Ficus Dod. communis B. pin. fructu rubro & albo.

Ficus Indica Dod. folio ſpinoſo, fructu majore C. B. Pin. Indica major Park. ☉

* Ficus Indica minor Park. Theat. Indica fol. ſpinoſo fructu minore C. B. P. ☉

Fili-

Filipendula Dod. vulgaris B. pin. ☉
Filix fœmina Dod. ramoſa major pinnulis obtuſis non dentatis B. pin. ☉
Filix floreſcens Tragi hiſt. Lugd. ramoſa non dentata florida B. pin. ☉
Filix mas Dod. non ramoſa dentata B. pin. ☉
Filix mas aculeata major B. prodr. ☉
Filix baccifera Cornuti. ☉
Filix non ramoſa petiolis tenuiſſimis, & tenuiſſimè dentatis B. pin. Filicula petræa prima Tab. ☉
* Filix arborea Tragi. ☉
* Filix paluſtris aculeata. ☉ ♃

Fœniculum Dod. vulgare Germanicum B. pin. ☉
Fœniculum dulce B. pin. Fœniculum .ſ. Marathrum vulgatius dulce Ad. Lob. deſcr. & Icon. ☉

Fœnum Græcum Dod. ſativum B. pin. ☿
Fœnum Græcum ſylveſtre alterum polycration B. pin. Hedyſarum minimum hiſt. Lugd. ☿

Fragaria Fuchſ. vulgaris B. pin. ☉
Fraga candida Dod. deſcr. ☉ Fra-

Digitized by Google

Fragaria fructu parvi Pruni magnitudine B. pin. Fraga fructu magno hort. Eyst. ☉
Fragaria flore duplici foliorum serie. ♃ ☉
Fragaria fructu rubro folioso. ♃ ☉
Fragaria sterilis B. pin. sylvestris minimè vesca .s. sterilis Icon. Lob. ☉
Fragaria sterilis incana. ♃ ☉

Fraxinella Clus. Dictamus albus vulgò .s. Fraxinella B. pin. ☉
Fraxinella niveo flore Clus. descr. ☉

Fraxinus Dod. excelsior B. pin.
Fraxinus florifera. ♃

Fritillaria præcox purpurea variegata B. pin. Meleagris Dod. ☉
Fritillaria pyrenæa Clus. hist. app. flore minore B. pin. ☉
* Fritillaria coronæ imperialis facie, pullo flore Hispanica umbellifera swertii umbellifera Park. Parad.
* Fritillaria præcox alba C. B. P. alba Park. Parad.

Frumentum Indicum Maiz dictum B. pin. Turcicum

Turcicum Dod. Idem granis rubis. ♀
Frumentum Sarracenicum Matth. ed. B. Fegopyrum Dod. ♀
Frumentum Indicum ſpicâ articulatâ, perenne. ♃ ☉

Fumaria Dod. officinarum & Dioſcoridis B. pin. Eadem flore albo. ♀
Fumaria minor tenuifolia P. pin. Capnos tenuifolia Cluſ. ♀
Fumaria ſiliquoſa ſempervirens Cornuti. ☉
Fumaria bulboſa radice non cava minor B. pin. Radix cava minor Dod. Eadem flore albo. ☉
Fumaria bulboſa radice cava major B. pin. Radix cava major Cluſ. Eadem flore albo. ☉
Fumaria tuberoſa inſipida Cornuti. ☉
* Fumaria tenuifolia ſemine compreſſo lini inſtar. ♃ ♀
* Fumaria Claviculis donata C. B. P. Capnos altera Belgica flore albo Cluſ. ♀

G.

GALEGA Dod. vulgaris B. pin. ☉
Galega flore albo hort. Eyst. ☉
Galegæ affinis, Sophera dicta B. pin. Sophera Prosp. Alp. de plant. Ægyp. ♀
* Galegæ affinis, Sesban Prosp. Alpini de plantis Ægyp. ♀

Gallium Dod. luteum B. pin. ☉
Gallium palustre album B. pin. album Tab. ☉
Gallium album tripetalon. ♃ ☉

Genista spinosa major longioribus aculeis B. pin. spinosa Dod. ☉
Genista spinosa brevioribus aculeis, an Genistella minor aspalathioïdes B. prodr. ☉
Genista spinosa minor Germanica B. pin. Genistella Dod. aculeata Obs. & Icon. Lob. ☉
Genistella spinosa Corrudæ similis. Scorpius Theophrasti hist. Lugd. Nepa Theophrasti Ad. Lob. ☉

Genista

Geniſta anguloſa, & ſcoparia B. pin. anguloſa .ſ. vulgaris hiſt. Lugd. ⊙
Geniſta tinctoria Germanica B. pin. tinctoria Dod. ⊙
* Geniſta purgans J. B. ⊙
* Geniſta Acacia trifolia C. B. Acacia Dioſc. Ger. ⊙

Gentiana major lutea B. pin. Gentiana Dod. ⊙
Gentiana Aſclepiadis folio B. pin. ſecunda foliis Aſclepiadis Dod. deſcr. ⊙
Gentianella Alpina anguſtifolia magno flore B. pin. helvetica Icon. Lob. major verna Cluſ. ⊙
Gentiana cruciata B. pin. Cruciata .ſ. Gentiana minor Dod. ⊙
Gentiana paluſtris anguſtifolia B. pin. Pneumonanthe Cord. hiſt. ⊙

Geranium batrachiodes folio Aconiti B. pin. ſecundum βατραχῶδες minus Cluſ. ⊙
Geranium batrachioïdes flore albo hort. Eyſt. ⊙
Geranium batrachiodes flore variegato B. pin. Anglicanum variegatum hort. Eyſt. ⊙

Gera-

Geranium batrachoïdes alterum longius radicatum Icon. Lob. batrachiodes odoratum B. pin. ☉

Geranium ſanguineum maximo flore B. pin. ſeptimum hæmatodes Cluſ. ☉

Geranium batrachiodes, Gratia Dei Germanorum Lob. B. pin. tertium batrachiodes majus Cluſ. ☉

* Geranium batrachiodes minus annuum purpurocæruleum. ♃ ♀

* Geranium batrachioïdes minus perenne verſicolor. ♃ ☉

Geranium montanum fuſcum B. pin. fuſcum hort. Eyſt. ☉

Geranium tuberoſum Dod. tuberoſum majus B. pin. ☉

Geranium nodoſum B. pin. quintum nodoſum Plateau Cluſ. ☉

Geranium lucidum ſaxatile B. pin. ſaxatile hort. Eyſt. ♀

Geranium folio Malvæ rotundo B. pin. Pes columbinus Dod. ♀

* Geranium Melachoïdes flore albo. ♃ ♀

Geranium folio Malvæ rotundo minus B. pin. deſcr. ♀

Geranium columbinum tenuiùs laciniatum B.

B. prodr. Idem flore albo. ♀
Geranium folio Althææ B. pin. malacoïdes Obſ. & Icon. Lob. ♀
Granium cicutæ folio moſchatum B. pin. ſupinum Dod. ♀
Geranium Coriandri folio odorum B. pin. Apulum Coriandri folio alterum odorum Col. ♀
Geranium Cicutæ folio minus & ſupinum B. pin. minus vel arvenſe Tab. ♀
Geranium arvenſe album Tab. ♀
Geranium robertianum Dod. robertianum primum B. pin. ♀
* Geranium Rubertianum flore albo. ♂ ♀
Geranium Creticum annuum hæmatodes Icon. Lob. latifolium longiſſimâ acu B. pin. ♀
Geranium triſte Cornuti. ⊙
Geranium flore verſicolore. ⊙
Geranium fœtidum Park. ⊙

Gingidium umbellâ oblongâ B. pin. Viſnaga Matth. ed. B. ♀

Gladiolus floribus uno verſu diſpoſitis major B. pin. Narbonenſis Obſ. & Icon. Lob. ⊙

Gladiolus flore albo Swert. ☉
Gladiolus floribus uno verſu diſpoſitis minor B. pin. minimus Lob. Ad. part. alt. ☉

Glaux maritima B. pin. exigua maritima Obſ. & Icon. Lob. ♆
* Glaux major flore herbaceo. ♃ ☉

Glycyrrhiza capite echinato B. pin. echinata Dodonæi hiſt. Lugd. ☉
Glycyrrhiza vulgaris Dod. ſiliquoſa vel Germanica B. pin. ☉
Glycyrrhiza ſylveſtris hiſt. Lugd. ſylveſtris floribus luteopalleſcentibus B. pin. ☉

Gnaphalium maritimum B. pin. marinum tomentoſum hiſt. Lugd. ☉
Gnaphalium Americanum Cluſ. latifolium Americanum B. pin. ☉
Gnaphalium vulgare majus B. pin. Filago .ſ. Impia Dod. ♆
Gnaphalium medium B. pin. Filago minor Dod. ♆
Gnaphalium minus repens B. pin. minimum Icon. Lob. ♆
Gnaphalium roſeum B. prodr. ♆

Gnapha-

Digitized by Google

Gnaphalium majus lato oblongo folio B. pin. Gnaphalium Plateau ſecundum Cluſ. ♀

* Gnaphalium 3 Plateau Cluſ. ♀

Gnaphalium Anglicum vel Belgicum folio longiore Icon. Lob. majus anguſto, oblongo folio alterum B. pin. ⊙

Gnaphalium montanum flore rotundiore B. pin. montanum variegatum hort. Eyſt. ⊙

Goſſipium .ſ. Xylon Dod. Goſſipium fruteſcens ſemine albo B. pin. ♀

Gramen caninum arvenſe .ſ. Gramen Dioſcoridis B. pin. caninum medicatum Obſ. Lob. ⊙

Gramen caninum vineale B. prodr. ⊙

Gramen caninum longiùs radicatum majus & minus B. pin. canarium longiùs radicatum latiore paniculâ Ad. Lob. part. alt. ⊙

Gramen caninum ſupinum minus B. pin. caninum ſupinum Icon. Lob. ⊙

Gramen nodoſum avenaceâ paniculâ B. pin. nodoſum hiſt. Lugd. ⊙

Gramen tremulum Hiſpanicum maximum Park. ♀

Digitized by Google

Gramen tremulum majus B.pin. leporinum, & tremulum Tab. ♆

Gramen tremulum minus paniculâ parvâ B. prodr. ♆

Gramen filiceum .ſ. polyanthes hiſt. Lugd. paniculis elegantiſſimis, .ſ. Heragroſtis major B. pin. ♆

Gramen minimum Dalechampii hiſt. Lugd. minimum paniculis elegantiſſimis B.pin. ♆

Gramen pratenſe paniculatum molle B. prodr. lanatum Dal. hiſt. Lugd. ☉

Gramen pratenſe vulgatius hiſt. Lugd. pratenſe paniculatum majus latiore folio, ~~mia~~ Theophraſti B. pin. ☉

Gramen pratenſe paniculatum medium B. pin. pratenſe ſecundum Dod. ☉

Gramen pratenſe paniculatum minus B pin. paniculatum minus album Tab. ☉

Gramen paniculâ multiplici B. prodr. Monſpeſſuli vulgò duriuſculum minus nominatum. ♆

Gramen arvenſe paniculâ criſpâ B. prodr. minimum xerampelinum Narbonenſe Icon. Lob. ♆

Gramen ſegetum paniculâ arundinaceâ B. pin. agrorum venti Spica Icon. Lob. ♆

Gramen

Gramen paniculatum folio variegato B. pin. ſulcatum vel ſtriatum album Icon. Lob. ☉
Gramen paluſtre paniculatum altiſſimum B. pin. aquaticum arundinaceum paniculatum Tab. ☉
Gramen pratenſe criſtatum .ſ. ſpicâ criſtatâ læve B. pin. criſtatum ejuſdem prodr. ☉
Gramen Caryophyllatæ foliis ſpicâ divulsâ B. pin. Caryophylleum Tab. ☉
Gramen typhoïdes maximum ſpicâ longiſſimâ B. prodr. ☉
Gramen typhoïdes ſpicâ multiplici B. pin. typhinum tertium Tab. ☉
Gramen alopecuroïdes ſpicâ aſperâ B. prodr. ♀
Gramen pratenſe tomentoſum paniculâ ſparsâ B. pin. eryophoron Dod. ☉
Gramen ſparteum ſpicatum latifolium B. pin. Spartum noſtras alterum Icon. Lob. ☉
Gramen ſparteum pennatum B. pin. Sparteum Auſtriacum pennatum Cluſ. ☉
Gramen foliolis junceis brevibus minus B. pin. exile durius Icon. Lob. ☉
Gramen junceum folio articulato aquaticum B. pin. aquaticum alterum Icon. Lob. ☉

Gramen juncoïdes junci ſparsâ paniculâ Park. ⊙

Gramen junceum ſpicatum .ſ. triglochin B. pin. Calamagroſtis quarta hiſt. Lugd. ⊙

Gramen cyperoïdes latifolium ſpicâ rufâ .ſ. caule triangulo B. pin. cyperoïdes hiſt. Lugd. ⊙

Gramen cyperoïdes ſpicâ pendulâ breviore B. pin. Pſeudocyperus Dod. ⊙

Gramen cyperoïdes paniculâ ſparsâ majus B. pin. cyperinum majus Tab. ⊙

Gramen cyperoïdes paluſtre majus ſpicâ compactâ B. pin. cyperinum paluſtre Tab. ⊙

Gramen cyperoïdes paluſtre majus ſpicâ divisâ B. pin. cyperoïdes parvum aquaticum Icon. Lob. ⊙

Gramen cyperoïdes maximum foliis & carinâ ſerratis. ⊙

Gramen cyperoïdes maritimum B. pin. maritimum hiſt. Lugd. ⊙

Gramen arundinaceum enode majus montanum B. pin. arundinaceum montanum Tab. ⊙

Gramen nemoroſum caliculis paleaceis B. pin. junceum bufonium Tab. ⊙

Gramen

ed by Google

Gramen hirſutum latifolium majus B. pin. hirſutum nemoroſum Icon. Lob. ♀

Gramen hirſutum anguſtifolium majus B. prodr. ♀

Gramen hirſutum capitulis Pſyllii B. pin. exile hirſutum cyperoïdes Icon. Lob. ♀

Gramen paluſtre aculeatum, vel minus Germanicum B. pin. paluſtre echinatum Icon. Lob. ⊙

Gramen dactylon folio arundinaceo majus B. pin. legitimum Cluſ. ⊙

Gramen dactylon folio latiore B. pin. Iſchæmon Plinii .ſ. dactylon hiſt. Lugd. ⊙

Gramen dactylon ſcoparium B. pin. ſcoparium Iſchæmi paniculis Gallicum Ad. Lob. part. alt. ⊙

Gramen dactylon eſculentum B. pin. Gramen mannæ primum Dod. ♀

Gramen paniceum ſpicâ longis ariſtis armatâ B. pin. Panici effigie gramen ariſtis circumvallatum Icon. Lob. ⊙

Gramen paniceum ſpicâ ſimplici B. pin. Panici effigie gramen ſimplici ſpicâ Icon. Lob. ⊙

Gramen miliaceum Icon. Lob. ſylvaticum paniculâ miliaceâ ſparsâ B. pin. ⊙

Gramen hordeaceum minus & vulgare B. pin. hordeaceum, Holcus Plinii hiſt. Lugd. ♆

Gramen loliaceum ſpicâ longiore B. pin. Lolium Dod. ♆

Gramen loliaceum minus ſpicâ ſimplici B. prodr. ♆

Gramen feſtucarium. Feſtuca graminea glumis hirſutis B. pin. Bromos ſterilis altera Icon. Lob. ♆

Gramen feſtucarium. Feſtuca avenacea ſterilis elatior B. pin. Bromos herba Dod. ♆

Gramen feſtucarium. Feſtuca graminea glumis vacuis B. prodr. ♆

Gramen avenaceum altero alteri innato Park. ♆

Gramen avenaceum ſquammoſâ glumâ Park. ♆

* Gramen Phalaroïdes perenne tuberoſâ radice. ♃ ⊙

* Gramen Feſtucarium capitulis duris, Ægylops Narbonenſis Lob. ♆

* Gramen Miliaceum ſylvaticum. ♆

* Gramen echinatum tribuloïdes Col. ♆

Gramen Parnaſſi Dod. albo ſimplici flore B. pin. ⊙

Grati-

Gratiola Dod. centauroïdes B. pin. ☉

Groſſularia ſimplici acino, vel ſpinoſa ſylveſtris B. pin. Uva criſpa Dod. ☉

* Groſſularia, Uva criſpa, fructu albo & Margaritis ſimili C. B. P. ☉

* Groſſularia Uva criſpa fructu rubro. ☉

Groſſularia rubra hiſt. Lugd. multiplici acino .ſ. non ſpinoſa hortenſis rubra .ſ. Ribes officinarum B. pin. ☉

Groſſularia hortenſis fructu Margaritis ſimili B. pin. Ribes fructu albo hort. Eyſt. ☉

Groſſularia hortenſis majore fructu rubro B. pin. Ribes major fructu rubro hort. Eyſt. ☉

Groſſularia non ſpinoſa fructu nigro B. pin. Ribeſium fructu nigro Dod. ☉

* Groſſularia ribes fructu rubro, folio Uvæ criſpæ. ♃ ☉

H.

HALIMUS latifolius .ſ. fruticoſus B. pin. primus Cluſ. ☉

Halimus .ſ. Portulaca marina B. pin. Portulaca marina, an Crithmum Dioſcoridis Icon. Lob. ☉

Hedera arborea B. pin. corymboſa Obſ. Lob. ☉

Hedera helix Dod. major ſterilis B. pin. ☉

Hedera terreſtris vulgaris B. pin. terreſtris Dod. Eadem flore carneo. ☉

Hedera ſaxatilis magno flore B. pin. Aſarina Lobelii hiſt. Lugd. ☉

Hedera trifolia Canadenſis Cornuti. ☉

Hedera quinquefolia Canadenſis Cornuti. ☉

* Hedera corymboſa fructu luteo, Hedera Dionyſii, & chryſocarpos Lugd. ☉

* Hedera trifolia Canadenſis folio ſinuato, ♃ ☉

Hedyſarum triphyllum Canadenſe Cornuti. ☉

Helenium Dod. vulgare B. pin. ☉

Helenium

Helenium Indicum maximum B. pin. Flos ſolis major hort. Eyſt. ♀

Helenium Indicum ramoſum B. pin. Flos ſolis prolifer hort. Eyſt. Idem ſemine albo, cinericeo, & ſtriato. ♀

Helenium Indicum tuberoſum. Flos ſolis Farneſianus, Aſter pervanus tuberoſus Col. ⊙

Helenium Canadenſe præaltum alato caule. ♃ ⊙

Helenium Canadenſe præaltum Voſacan dictum. ♃ ⊙

Helenium Canadenſe humilius latifolium ♃ ⊙

Helichryſum foliis oblongis Stæchadi citrinæ ſimilibus B. pin. Chryſocome peregrina hort. Eyſt. ⊙

Helichryſum ſ. Stœchas citrina auguſtifolia B. pin. Stœchas citrina Dod. ⊙

Heliotropium majus Cluſ. majus Dioſcoridis B. pin. ♀

Heliotropium minus tricoccum Cluſ. tricoccum B. pin. ♀

Helleborus niger fœtidus B. pin. Helleboraſter maximus flore & ſemine prægnans Obſ. & Icon. Lob. ⊙

Helleborus niger hortenſis flore viridi B. pin. niger adulterinus hortenſis Fuchſ. ⊙

Helleborus niger flore roſeo B. pin. niger legitimus Cluſ. ⊙

Helleborus niger flore roſeo minor Belgicus. ♃ ⊙

Helleborus niger trifolius Ald. hort. Farn. ☿

Helleborus niger Saniculæ folio major B. pin. Aſtrantia nigra Obſ. & Icon. Lob. ⊙

Helleborus Saniculæ folio minor B. prodr. ⊙

Helleborus Hippocratis Tab. niger tenuifolius Buphtalmi flore B. pin. ⊙

Helleborus albus flore ſubviridi B. pin. albus flore herbaceo hort. Eyſt. ⊙

Helleborus albus flore atrorubente B. pin. hort. Eyſt. ⊙

Helleborine latifolia montana B. pin. Epipactis latifolia hort. Eyſt. ⊙

Helleborine anguſtifolia paluſtris, ſ. pratenſis B. pin. Helleborine, ſive Epipactis recentiorum Obſ. & Icon. Lob. ⊙

Helleborine flore albo, vel Damaſonium mon-

Digitized by Google

montanum latifolium B. pin. Epipactis anguſtis foliis hort. Eyſt. ⊙

Helleborine montana anguſtifolia purpuraſcens B. pin. recentiorum ſexta Cluſ. ⊙

Helleborine herbaceo flore. ♃ ⊙

* Helleborine .ſ. calceolus marianus, vide calceolus mariæ.

Heſperis hortenſis B. pin. Viola matronalis flore albo hort. Eyſt. Eadem flore purpureo quæ

Viola matronalis flore purpureo hort. Eyſt. ⊙

Heſperis flore pleno albo. Viola matronalis flore multiplici Cornuti deſcr. ⊙

* Heſperis Syriaca Eyſt. peregrina ſiliquis articulatis C. B. P. ♀

* Heſperis ſylveſtris hirſuta foliis Hieracii B. prodr. ♀

Hieracium majus folio Sonchi, vel Hieracium Sonchites B. pin. majus Fuchſ. ⊙

Hieracium majus foliis Sonchi ſemine curvo B. pin. Creticum pro Endivia lutea miſſum Cam. deſcr. Sonchus aſper laciniatus Creticus B. prodr. ♀

Hiera-

Hieracium arboreſcens paluſtre B. pin. Sonchus lævior Aſtriacus altiſſimus quintus Cluſ. ☉

Hieracium Apulum flore ſuaverubente Col. dentis leonis folio flore ſuaverubente B. pin. ♀

Hieracium odore Apuli ſuaverubentis. ♃ ♀

Hieracium dentis leonis folio monoclonon B. pin. intybaceum Tab. ☉

Hieracium dentis leonis folio hirſutie aſperum minùs laciniatum B. pin. & prodr. ☉

Hieracium dentis leonis folio hirſutie aſperum magis laciniatum B. pin. ☉

Hieracium minus dentis leonis folio oblongo glabro B. pin. Hyoſeris anguſtifolia Tab. ☉

Hieracium minus folio ſubrotundo B. pin. minimum nonum Cluſ. ♀

Hieracium maximum aſperum Chondrillæ folio B. prodr. ♀

Hieracium Chondrillæ folio glabrum B. pin. aphacoïdes Tab. ♀

Hieracium Chondrillæ folio glabro radice ſuccisâ majus B. pin. minus præmorsâ radice Obſ. & Icon. Lob. ☉

Hiera-

Hieracium capitulum inclinans ſemine adunco B. pin. facie Hedypnoïs Icon. Lob. ♀

Hieracium ſiliquâ falcatâ B. pin. Narbonenſe falcatâ ſiliquâ Lobelii hiſt. Lugd. ♀

Hieracium echioïdes capitulis cardui benedicti B. pin. echioïdes, Bugloſſum luteum Tab. ♀

Hieracium octavum folio Hedypnoïdis Cluſ. montanum dentis leonis folio incano, vel laciniatum lanuginoſo folio B. pin. ☉

Hieracium phlomoïdes Tab. Alpinum latifolium hirſutie incanum flore magno B. pin. ☉

Hieracium montanum tomentoſum. An villoſum quintum Cluſ. ☉

Hieracium montanum majus latifolium Tab. montanum latifolium glabrum majus B. pin. ☉

Hieracium murorum folio piloſiſſimo B. pin. Corchorus Dal. hiſt. Lugd. ☉

Hieracium murorum folio non maculato minus piloſo. ♃ ☉

Hieracium fruticoſum latifolium foliis dentatis glabrum B. pin. majus latifolium vel

ſecundum Cluſ. deſcr. fruticeſcens latifolium polyanthos hort. Eyſt. ☉

Hieracium Sabaudum Penæ & Lobelii hiſt. Lugd. fruticoſum anguſtifolium majus B. pin. ☉

Hieracium montanum alterum λεπτομακρόκαυλον Col. ☿

* Hieracium foliis & floribus Dentis leonis bulboſi ſemine curvo. ♃ ☿

* Hieracium anguſtifolium glabrum .ſ. Piloſilla major glabra. ♃ ☉

* Hieracium calice barbato Col. proliferum falcatum C. B. P. ☿

* Hieracium ramoſum magno flore B. in prodr. 14 ☿

* Hieracium Pulmonaria non maculata indeſectis foliis. ♃ ☿

Holoſteum hirſutum albicans minus B. pin. Salmanticenſe minus Cluſ. ☉

Holoſteum ſtrictiſſimo folio majus B. pin. Serpentina Matthioli Dod. ☉

Holoſteum ſtrictiſſimo folio minus B. pin. Serpentina omnium minima Lobelii hiſt. Lugd. ☉

Holoſteum Alſines flore. ♃ ☉

Holoſteo

Holoſteo affinis, Cauda muris B. pin. Cauda murina Dod. ♀

Hordeum polyſtichum hybernum B. pin. polyſtichum Fuchſ. ♀
Hordeum diſtichon Dod. B. pin. ♀

Horminum Creticum Proſp. Alp. de exot. ſativum B. pin. ♀
Horminum Sclarea dictum B. pin. Sclarea Matth. ed B. Obſ. Lob. ☉
Horminum luteum glutinoſum B. pin. Colus Jovis Obſ. & Ad. Lob. ☉
Horminum ſylveſtre latifolium verticillatum B. pin. ſylveſtre tertium Cluſ. ☉
Horminum pratenſe foliis ſerratis B. pin. ſylveſtre Fuchſ. hiſt. Lugd. Idem flore albo ☉
Horminum anguſtifolium laciniatum flore cœruleo. An ſylveſtris quarti, quinta ſpecies Cluſ. ☉
Horminum Lapathi unctuoſi folio. ♃ ☉
* Horminum flore minore folio integro. ♃ ☉
* Horminum flore minore dilute cœruleo folio ſmuato. ♃ ♀

* Horminum flore majore amplo violaceo Lobis punctis aureis notatis. ♃ ♀
* Horminum flore majore Salviæ folio forte Marum Ægyptiacum Prosp. Alp. ♀
* Hormium flore majore sylvestris 5 altera species Clus. ☉

Hyacinthus comosus major purpureus B. pin. comosus spurius hort. Eyst. Idem flore albo & carneo. ☉
Hyacinthus Sanneſius comâ paniculosâ Col. paniculâ cœruleâ B. pin. ☉
Hyacinthus racemosus cœruleus major B. pin. botryoïdes cœruleus hort. Eyst. ☉
Hyacinthus racemosus cœruleus minor latifolius B. pin. botryoïdes cœruleus hort. Eyst. ☉
Hyacinthus botryoïdes floribus albis hort. Eyst. ☉
Hyacinthus racemosus cœruleus minor juncifolius B. pin. racemosus Dod. comosus minor Obs. Lob. ☉
Hyacinthus racemosus moschatus B. pin. Muscari obsoletiore flore Clus. ☉
Hyacinthus racemosus moschatus luteus B. pin. Muscari flavo flore Clus. ☉

Hyacin-

Digitized by Google

Hyacinthus oblongo flore cœruleus major B. pin. Anglicus & Belgicus Obſ. & Icon. Lob. ⊙

Hyacinthus Belgicus candidus Icon. Lob. ⊙

Hyacinthus floribus Campanulæ utrinque diſpoſitis B. pin. Neapolitanus Swert. ⊙

Hyacinthus oblongo cœruleo flore minor B. pin. minor Hiſpanicus Cluſ. cur. poſt. deſcr. Ger. Icon. ⊙

Hyacinthus orientalis albus primus B. pin. Orientalis niveus Cluſ. ⊙

Hyacinthus brumalis orientalis Geminæ ex Lobelio hiſt. Lugd. ⊙

Hyacinthus orientalis maximus B. pin. Orientalis polyanthos Cluſ. ⊙

Hyacinthus orientalis flore duplici hort. Eyſt. B. pin. ⊙

Hyacinthus orientalis flore cœruleo pleno hort. Eyſt. ⊙

Hyacinthus ſtellatus vulgaris Cluſ. ſtellaris trifolius B. pin. ⊙

Hyacinthus ſtellatus albo flore Cluſ. Idem flore carneo. ⊙

Hyacinthus ſtellatus cineracei coloris Cluſ. ſtellaris Sommeri Ger. Stellaris ſpicatus cinereus B. pin. ⊙

Hyacinthus ſtellatus lilifolius Cluſ. Stellaris foliis & radice Lilii B. pin. ☉

Hyacinthus ſtellaris autumnalis minor B. pin. autumnalis minor Dod. ☉

Hyacinthus ſtellatus Periuanus Cluſ. Indicus bulboſus ſtellatus B. pin. Idem flore albo. ☉

Hyacinthus Indicus tuberoſâ radice Cluſ. Indicus tuberoſus flore Hyacinthi orientalis B. pin. ☉

Hyoſcyamus vulgaris vel niger B. pin. niger Dod. ☿

Hyoſcyamus albus Dod. albus major vel tertius Dioſcoridis B. pin. ☿

Hyoſcyamus aureus Proſp. Alp. exot. ☿

Hypecoon legitimum Cluſ. Hypecoum B. pin. ☿

Hypericum Dod. vulgare B. pin. ☉

Hypericum minus ſupinum, vel ſupinum glabrum B. pin. minimum ſupinum Septentrionale Icon. Lob. ☉

Hypericum ſupinum tomentoſum alterum Cluſ. ſupinum Dod. ☉

Hype-

Hypericum fruteſcens Americanum flore albo Park. Pruno ſylveſtri affinis Canadenſis B. pin. app. ☉

Hyſſopus vulgaris Dod. officinarum cœrulea, ſ. ſpicata B. pin. ☉
Hyſſopus rubro flore hort. Eyſt. B. pin. ☉
Hyſſopus flore albo hort. Eyſt. vulgaris alba B. pin. ☉
* Hyſſopus hirſuta & ſtriata. ☉
Hyſſopus latifolia B. prodr. ☉
Hyſſopus anguſtifolia ſpicato flore Cluſ. anguſtifolia ſpicata B. pin. ☉

Hyſſopifolia, ſ. Gratiola minor B. prodr. Gratiola minor Matth. ed. B. ☉

I.

JACEA Auſtriaca villoſo capite Cluſ. latifolia capite hirſuto B. pin. ☉
Jacea Luſitanica ſempervirens. ♂♂ ☉
Jacea nigra Dod. nigra laciniata B. pin. ☉
Jacea lutea capite ſpinoſo B. pin. luteo flore Cluſ. ☉

Jacea cyanoïdes echinato capite B. pin. Stœbe Salmantica ſecunda Cluſ. Cyanoïdes flos Dod. ☉
Jacea latifolia purpurea capitulo ſpinoſo B. prodr. ☉
Jacea capite longis aculeis ſpinoſo B. prodr. ☉ ☉
Jacea montana incana capite Pini B. pin. ♀
Jacea Oleæ folio capitulis ſimplicibus B. pin. Ptarmica Auſtriaca Dod. Cluſ. ♀
Jacea Oleæ folio minore flore B. pin. Ptarmicæ Auſtriacæ ſpecies Cluſ. cur. poſt. ♀
* Jacea Babylonica B. prodr. ☉
* Jacea fruticans Pini folio C. B. Stœbe capitata Roriſmarini folio Ponæ. ☉
* Jacea integri-folia humilis C. B. ☉
* Jacea pumila Adv. Lob. ☉

Jacobæa Dod. vulgaris laciniata B. pin. ☉
Jacobæa foliis amplioribus incanis. ☉
Jacobæa maritima B. pin. Cineraria hiſt. Lugd. ☉
* Jacobæa Crithmi litorei foliis. ♃ ☉
* Jacobæa Senecionis folio. ♃ ♀

Jaſminum Dod. vulgatius flore albo B. pin. ☉ Jaſmi-

Jasminum humilius magno flore B. pin. Gelseminum Catalonicum hort. Eyst. ☉

Jasminum humile luteum B. pin. flore luteo hort. Eyst. ☉

Jasminum luteum vulgo dictum bacciferum B. pin. Polemonium Monspelienſium Lob. ☉

Jasminum flavum odoratum. Gelſeminum Indicum flavum odoratiſſimum Bapt. Ferr. ☉

Jaſminum purpureum prælongis ſiliquis. Gelſeminum hederaceum Indicum Cornuti. Indicum flore phœniceo Bapt. Ferr. ☉

Iberis Cardamantica Obſ. Lob. Iberis latiore folio B. pin. ☉

* Iberis Naſturtii folio Matth. ed. ♄

* Iberis virginiana latifolia ramoſa, ♄

Ilex major glandifera Dod. folio rotundiore molli modiceque ſpinato B. pin.

Ilex minor coccigera Dod. aculeata coccigландifera B. pin.

Imperatoria Matthioli hiſt. Lugd. major B. pin. ☉

Intybus ſativa latifolia, ſ. Endivia vulgaris B. pin. Intybum ſativum Dod. ♀

Intybus ſativa anguſtifolia B. pin. minor Matth. ed. B. ♀

Intybus criſpa B. pin. Intubum criſpum hiſt. Lugd. ♀

Iris noſtras vulgaris Icon. Lob. vulgaris Obſ. ejuſdem vulgaris Germanica, ſ. ſylveſtris B. pin. ⊙

Iris alba florentina B. pin. florentina hort. Eyſt. ⊙

Iris latifolia major, ſ. Suſiana Cluſ. Suſiana flore maximo ex albo nigricante B. pin. ⊙

Iris Dalmatica major hiſt. Lugd. Aſiatica cœrulea polyanthos B. pin. Eadem flore violaceo, & dilutius violaceo. ⊙

Iris exotica Cam. major latifolia 14. Cluſ. latifolia odore Oxyacanthæ B. pin. ⊙

Iris latifolia pannonica colore multiplici B. pin. major latifolia 15. Cluſ. ⊙

Iris paluſtris lutea Tab. Acorus adulterinus B. pin. ⊙

Iris Tripolitana Cluſ. deſcr. media longiſſimis foliis lutea B. pin. ⊙

Iris ſylveſtris Byzantina peramœna Icon. Lob.

Lob. pratenſis anguſtifolia non fœtida altior B. pin. ⊙

Iris anguſtifolia prunum redolens minor B. pin. perpuſilla ſylveſtris anguſtifolia Obſ. Lob. ⊙

Iris bulboſa latifolia .ſ. prima Cluſ. bulboſa latifolia acaulos odora B. pin. Eadem flore albo. ⊙

Iris Perſica variegata præcox Bapt. Ferr. ⊙

Iris bulboſa Anglicana flore cœruleo hort. Eyſt. ⊙

Iris bulboſa Anglicana major flore candido hort. Eyſt. ⊙

Iris bulboſa violaceo flore Cluſ. foliis cepaceis maculatis flore purpureo inodoro B. pin. ⊙

Iris bulboſa latifolia vulgò dicta Iris l' Abbé. ♃ ⊙

Iris alba oris cœruleis Swert. & Rob. in horto Valet. ⊙

Iris bulboſa lutea verſicolor B. pin. ⊙

Iris tuberoſa Belgarum Icon. Lob. tuberoſa folio anguloſo B. pin. ⊙

* Iris tuberoſa matitima C. B. P. ⊙

* Iris bulboſa lutea major. ♃ ⊙

Iſatis ſativa, vel latifolia B. pin. Glaſtum ſativum Obſ. & Icon. Lob. ♀

Jujubæ majores oblongæ B. pin. Ziziphus rutila Cluſ.

Juncus acutus paniculâ ſparsâ B. pin. acutus Dod. ⊙

Juncus maximus .ſ. Scirpus major B. pin. Holoſchœnos Dod. ⊙

Juncus lævis paniculâ ſparsâ major B. pin. lævis Dod. ⊙

Juncus lævis paniculâ non ſparsâ B. pin. lævis glomerato flore Icon. Lob. ⊙

Juncellus omnium minimus Ad. Lob. deſcr. ⊙

Juncus capitulis Equiſeti major B. pin. clavatus Dalechampii hiſt. Lugd. ⊙

Juncus Lithoſpermi ſemine. ♃ ⊙

* Juncus acutus caule triangulo B. in prodr. ⊙

* Juncus acutis Sorghi capitulis. ♃ ⊙

Juniperus Dod. vulgaris fruticoſa B. pin.

K.

KALI geniculatum majus B. pin. Salicornia Dod. ☿

Kali ſpinoſo affinis B. pin. Tragum Matth. ed. B. ☿

* Kali ſempervirens grandius perpetuum B. prodr. deſcr. ☉

* Kali minus Geſn. hort. album Ger. ☿

L.

LACTUCA ſativa B. pin. ſativa folio Scariolæ Obſ. & Icon. Lob. ☿

Lactuca Romana dulcior nigriore, & Scariolæ hortenſis folio, ſemine nigro Ad. Lob. folio obſcurius virente, ſemine nigro B. pin. ☿

Lactuca maculoſa B. pin. fuſci ac veluti purpurei coloris .ſ. ex albido rubicunda Dod. deſcr. ☿

Lactuca capitata Dod. B. pin. ☿

Lactuca Italica laciniata B. prodr. ♆
Lactuca criſpa Dod. B. pin. ♆
Lactuca criſpa ſecunda Tab. ♆
Lactuca ſylveſtris coſtâ ſpinosâ B. pin. ſylveſtris Dod. ♆
Lactuca ſylveſtris odore viroſo B. pin. agreſtis Ad. & Icon. Lob. ♆
Lactuca montana purpurocœrulea major B. pin. Sonchus montanus purpureus πλεσπίταλον Col. ⊙

Lamium purpureum fœtidum folio ſubrotundo B. pin. Urtica iners altera Dod. Idem flore albo. ♆
Lamium purpureum, vel album non fœtens folio oblongo B. pin. Urtica iners .ſ. Lamium primum Dod. Idem flore luteo. ⊙
Lamium folio oblongo luteum B. pin. luteum Icon. Lob. ⊙
Lamium maculatum, & albâ lineâ notatum B. pin Lamium Plinii Campoclarenſe & montanum Col. ⊙
Lamium foliis caulem ambientibus majus B. pin. Ballote criſpa major hiſt Lugd. ♆
Lamium maximum ſylvaticum B. pin. Urtica

tica herculea Tragi hiſt. Lugd. Galeopſis legitima Cluſ. ☉

Lamium maximum ſylvaticum alterum B. pin. Galeopſis maxima pannonica Cluſ. ☉

Lamium montanum Meliſſæ folio B. pin. Meliſſophyllum verum Fuchſ. Idem flore albo quod

Lamium pannonicum primum albo flore Cluſ. ☉

Lamium peregrinum .ſ. Scutellaria B. pin. Lamium aſtragaloides Cornuti. Caſſida Col. ☉

* Lamium Parietariæ facie. ♃ ☉

Lapathum ſativum Dod. utr. ed. Hippolapathum hortenſe hiſt. Lugd. ☉

Lapathum hortenſe .ſ. Spinacia ſemine ſpinoſo B. pin. Spinacia hiſt. Lugd. ♀

Lapathum hortenſe ſ. ſpinacia ſemine ſpinoſo B. pin. Spinachia ſemine non ſpinoſo Matth. ed. B. ♀

Lapathum folio acuto plano B. pin. ſylveſtre .ſ. Oxylapathum Dod. ☉

Lapathum folio acuto criſpo B. pin. acutum criſpum Tab. ☉

Lapathum ſanguineum Obſ. Lob. folio acuto rubente B. pin. ♀ La-

Lapathum acutum minimum Icon. Lob. minimum B. pin. ☉

Lapathum unctuoſum B. pin. Bonus henricus Matth. ed. B. ♀

Lapathum folio ſubrotundo B. pin. folio minus acuto Icon. Lob. fidium inſtar. J. B. ☉

Lapathum rotundifolium Cam. Hippolapathum rotundifolium montanum B. pin. ☉

Lapathum aquaticum folio cubitali B. pin. Hydrolapathum majus Obſ. Lob. ☉

Lappa major, Arcium Dioſcoridis B. pin. Bardana, ſive Lappa major Dod. ♀

Lappa minor, Xanthium Dioſcoridis B. pin. Xanthium Fuchſ. ♀

Larix Dod.

Laſerpitium Gallicum B. pin. è regione Maſſiliæ repertum Ad. Lob. ☉

Laſerpitium alterum. ♃ ☉

* Laſerpitium fol. Lobatis auguſtis. ♃ ☉

* Laſerpitium fol. Lobatis latioribus ſem. criſpo, coſtus amara ſem. criſpo J. B. ☉

Lathy-

Lathyris major B. pin. Cataputia vulgaris hort. Eyst. ♀

Lathyrus sativus flore fructúque albo B. pin. Aracus sive Cicera Dod. Idem flore purpureo. ♀

Lathyrus sylvestris minor B. pin. Catanance Dodonæi hist. Lugd. ♀

Lathyrus latifolius B. pin. Clymenum Matthioli Clus. ☉

Lathyrus latifolius annuus siliquâ hirsutâ. ♃ ☉

Lathyrus sylvestris luteus foliis viciæ B. pin. Legumen terræ glandibus simile Dod. ☉

Lathyrus arvensis repens tuberosus B. pin. Terræ glandes Dod. ☉

Lathyrus flore obsoleto, siliquâ pisi minoris. ♃ ♀

Lathyrus Bæticus flore luteo Park. ♀

* Lathyrus Narbonensis angustifol. major J. B. ☉

* Lathyrus angustifol. Bæticus elegans siliquis Orobi Park. ♀

* Lathyrus πολύλοβος, siliquâ breviori, fol. tenuiori, flore rubro. J. B. ♀

* Lathyrus πολύλοβος, siliquâ pisi minoris flore rubro. ♃ ♀

* Lathyrus λειπόλοβος, sem. hexedro, flore cœruleo. ♃ ♀
* Lathyrus λειπόλοβος, sem. rotundo, flore rubello. ♃ ♀

Lavandula latifolia B. pin. Nardus Italica Matth. ed. B. ☉
Lavandula anguſtifolia B. pin. altera Dod. Eadem flore albo quæ
Lavandula alba Icon. Lob. ☉
Lavandula folio diſſecto B. pin. Lavendula multifido folio Cluſ. ♀

Laureola Dod. ſempervirens flore viridi, quibuſdam Laureola mas B. pin. ☉
Laureola folio deciduo, flore purpureo B. pin. Mezereum Germanicum Obſ. Lob. Idem flore albo B. pin. deſcr. ☉

Laurus vulgaris B. pin. Laurus Matth. ed. B.
Laurus tinus cœruleâ baccâ Icon. Lob. Sylveſtris Corni fœminæ foliis ſubhirſutis B. pin,
* Laurus Indica hort. Ferneſiani.
Laurus Alexandrina fructu pediculo inſidente B. pin. Hippogloſſum Dod. ☉

Laurus

Laurus Alexandrina fructu folio insidente B. pin. Alexandrina vera hist. Lugd. ☉
Laurus Alexandrina fructu longis pediculis caulibus alligato. ♃ ☉

Lenticula palustris quadrifolia B. pin. Lemna Theophrasti hist. Lugd. ♀

Lens vulgaris B. pin. minor Dod. ♀
Lens major Tab. B. pin. ♀

Lentiscus Dod. vulgaris B. pin.

Leontopetalon Dod. ☉

Lepidium Plinii Dod latifolium B. pin. ☉
Lepidium annuum Lobelii hist. Lugd. glastifolium B. pin. ♀
Lepidium, Dentellaria dictum B. pin. Plumbago quorundam Clus. ☉

Leucoium bulbosum vulgare B. pin. bulbosum hexaphyllum Dod. ☉
Leucoium bulbosum trifolium minus B. pin. bulbosum triphyllum Dod. ☉
Leucoium incanum majus B pin. candidum majus Dod. ☉

Leucoium purpureum vel rubrum B. pin. flore rubro ſimplici hort. Eyſt. Idem, quod violaceum Tab. ⊙

Leucoium incanum majus variegatum ſimplici flore B. pin. ⊙

Leucoium incanum majus multiplex B. pin. flore pleno albo, rubro pleno hort. Eyſt. Idem flore pleno violaceo. ⊙

Leucoium incanum majus variegatum pleno flore B. pin. purpureum variegatum flore pleno hort. Eyſt. ⊙

Leucoium maritimum ſinuato folio B. pin. marinum majus Cluſ. ⊙

Leucoium maritimum anguſtifolium B. pin. marinum minus Cluſ. ⊙

Leucoium alyſſoïdes clypeatum majus B. pin. Alyſſon Dioſcoridis Dod. ♀

Leucoium incanum ſiliquis rotundis B. pin. marinum Patavinum Obſ. & Icon. Lob. Eruca peregrina Cluſ. ⊙

Leucoium album odoratiſſimum folio viridi B. prodr. ⊙

Leucoium luteum magno flore B. pin. flos Keiri ſimplex medius hort. Eyſt. ⊙

Leucoium luteum vulgare B. pin. luteum Dod. ⊙

Leu-

Leucoium luteum ſylveſtre anguſtifolium B. pin. ſylveſtre Cluſ. ⊙
Leucoium flore luteo pleno majus hort. Eyſt. B. pin.
Idem flore pleno variegato. ⊙
Leucoium flore luteo pleno minus hort. Eyſt. ⊙
*Leucoium montanum flore pedato Col. ⊙

Libanotis latifolia altera .ſ. vulgatior B. pin. Seſeli Æthiopicum herba Dod. ⊙
Libanotis latifolia Aquilegiæ folio B. prodr. ⊙
Libanotis Ferulæ folio, ſemine anguloſo B. pin. Cachrys verior, Libanotis Galeno Obſ. Lob. ⊙
Libanotis Ferulæ folio & ſemine B. pin. Panaces Aſclepium Dod. ⊙

Liguſticum vulgare, An Libanotis fertilis Theophraſti B. pin. Leviſticum vulgare Dod. ⊙
Liguſticum, quod Seſeli officinarum B. pin. Siler montanum Dod. ⊙
Liguſticum .ſ. Siler montanum anguſtifolium B. prodr. ☉

Liguſtrum Matth. ed. B. Germanicum B. pin.

Liguſtrum foliis laciniatis B. pin. nigrum Proſp. Alp. de exot. Agem Lilag Perſarum Cornuti.

Lilium candidum Dod. album flore erecto & vulgare B. pin.

Lilium album inodorum flore pleno. ⊙

Lilium purpurocroceum majus B. pin. Hemerocallis Matth. ed. B. ⊙

Lilium floribus reflexis montanum B. pin. montanum flore purpuraſcente punctatum hort. Eyſt. ⊙

Lilium floribus reflexis alterum lanugine hirſutum B. pin. flore purpuraſcente non punctatum hort. Eyſt. ⊙

Lilium floribus reflexis albis non punctatis B. pin. montanum minus album non punctatum hort. Eyſt. ⊙

Lilium purpuro-ſanguineum flore reflexo B. pin. Hemerocallis Chalcedonica purpuro-ſanguinea polyanthos Obſ. Lob. ⊙

Lilium flavum anguſtifolium B. pin. montanum flavo flore Lobel. Ad. part 2ª. ⊙

Lilium ſ. Corona imperialis B. pin. Corona imperialis Dod. ⊙ Li-

Lilium .ſ. Corona imperialis flore luteo B. pin. deſcr. & Cluſ. deſcr. ⊙

Lilium .ſ. Corona imperialis duplici coronâ B. pin. Corona imperialis florum claſſe duplici hort. Eyſt. ⊙

* Lilium .ſ. Corona imperialis fol. vario ♃ ⊙

Lilium Perſicum Dod. ⊙

Lilium luteum Aſphodeli radice B. pin. Liliaſphodelus luteo flore Cluſ. ⊙ Anguſtifolius flore luteo.

Lilium rubrum Aſphodeli radice B. pin. Liliaſphodelus puniceus Cluſ. deſcr. ⊙

* Liliaſphodelus latifolius flore luteo. ♃ ⊙

* Lilium .ſ. Martagon Canadenſe. ♃ ⊙

Lilionarciſſus Indicus Bapt. Ferr. ⊙

Lilium convallium Matth. ed B. album B. pin. ⊙

Lilium convallium flore rubente B. pin. flore ſuavérubente Icon. Lob. ⊙

Lilium convallium minus B. pin. Monophyllon Matth. ed. B. ⊙

Limonium Matth. ed. B. maritimum majus B. pin. ⊙

Limonium parvum Bellidis minoris folio B. pin. pumilum Cluſ. deſcr. ☉

Limonium Rauwolfii .ſ. peregrinum Cluſ. cur. poſt. peregrinum foliis Aſplenii B. pin. ♀

Linaria latifolia Dalmatica magno flore B. prodr. ♀

Linaria Bellidis folio B. prodr. purpurea odorata Obſ. & Icon. Lob. ☉

Linaria ſcoparia B. pin. Oſyris Dod. ♀

Linaria vulgaris lutea flore majore B. pin. prima Dod. ☉

Linaria lutea flore minimo B. pin. ♀

Linaria aurea Tragi Obſ. & Icon. Lob. folioſo capitulo luteo major B. pin. ☉

Linaria folioſo capitulo luteo minor B. pin. aurea minor Tab. ☉

Linaria purpurea major odorata B. pin. altera purpurea Dod. ☉

Linaria arvenſis cærulea erecta B. prodr. ♀

Linaria montana floſculis albicantibus B. pin. Anthyllis montana hiſt. Lugd. ☉

Linaria quadrifolia lutea B. pin. tetraphylla lutea Col. ♀

Linaria cærulea repens B. pin. Oſyris purpurocærulea

purocœrulea repens Icon. Lob. Eadem flore albo. ☉

Linaria minima hirſuta flore obſoleto. ♃ ♀

Linaria minor cœrulea recta Park. ♀

* Linaria minor viſcoſa flore luteo. ♃ ♀

* Linaria Armoriaca flore ſaturate violaceo. ♃ ♀

* Linaria major anguſtifolia glabra flore luteo maj. ♃ ♀

Lingua cervina officinarum B. pin. Phyllitis Dod. ☉

Lingua cervina multifido folio B. pin. Phillitis laciniata Dod. ☉

Linum ſativum Dod. B. pin. ♀

Linum ſylveſtre luteum foliis ſubrotundis B. pin. Campanula lutea linifolia Penæ hiſt. Lugd. ☉

Linum maritimum luteum B. pin. marinum luteum Narbonenſe Obſ. & Icon. Lob. ☉

Linum ſylveſtre anguſtifolium floribus dilutè purpuraſcentibus, vel carneis B. pin. ſylveſtre ſextum anguſtifolium Cluſ. deſcr. ☉

Linum ſylveſtre minus B. pin. Chamælinum

ſ. Linum ſylveſtre perpuſillum Obſ. & Icon. Lob. ♀

Linum pratenſe floſculis exiguis B. pin. ſylveſtre catharticum Ger. Linocarpos Thal. ♀

* Linum annuum luteum folio latiore. ♂ ♀

* Linum annuum luteum folio anguſtiore. ♂ ♀

* Linum perenne cœruleum ſativo æmulum. ♂ ⊙

* Linum minimum ſtellatum B. prodr. ♀

Lithoſpermum arundinaceum Dal. hiſt. Lugd. arundinaceum forte Dioſcoridis & Plinii B. pin. ♀

Lithoſpermum majus erectum B. pin. minus Dod. ⊙

Lithoſpermum minus repens latifolium B. pin repens majus Cluſ. ⊙

Lithoſpermum arvenſe radice rubra B pin. nigrum Dal. hiſt. Lugd. ♀

Lithoſpermum Linariæ folio Germanicum B. pin. Linaria altera botryoïdes montana Col. ♀

Lonchitis

Lonchitis âſpera Dod. minor B. pin. ☉

Lotus hortenſis odora B. pin. Trifolium odoratum alterum ſ. Lotus ſativa Dod. ♀

Lotus ſ. Melilotus pentaphyllos minor glabra B. pin. Trifolium corniculatum primum Dod. ☉

Lotus pentaphyllos minor hirſutus ſiliquâ anguſtiſſimâ B. pin. Trifolium corniculatum minus piloſum ejuſdem prodr. deſcr. ☉

Lotus pentaphyllos ſiliquoſus villoſus B. pin. hæmorrhoïdalis major ſ. Trifolium hæmorrhoïdale majus Park. ☉

Lotus ſiliquoſus glaber flore rotundo B. pin. Lybica Dal. hiſt. Lugd. ☉

Lotus fruticoſus incanus ſiliquoſus B. pin. aſperior fruticoſa ſ. Narbonenſis incana Obſ. & Icon. Lob. ☉

Lotus pratenſis ſiliquoſus luteus B. pin. trifolia pratenſis ſiliquoſa Monſpelienſium Icon. Lob. ☉

Lotus ruber ſiliquâ angulosâ B. pin. tetragonolobus Camerarii hort. Eyſt. ♀

Lotus ſiliquis Ornithopodii B. pin. peculiaris ſiliquis Ornithopodii Cam. ♀

Lotus

Lotus pentaphyllos veſicaria B. pin. Trifolium halicacabum Cam. ♆

* Lotus ſiliquis ſingularibus, binis Terniſve J. B. ♆

Loto affinis vulneraria pratenſis B. pin. Anthyllis leguminoſa Belgarum Obſ. & Icon. Lob. ☉

Loto affinis hirſuta flore ſubrubente B. pin. Anthyllis leguminoſa purpuraſcente flore Cluſ. deſcr. ☉

Lotus arbor Obſ. Lob. Lotus fructu Ceraci B. pin.

Lotus latifolia Africana B. pin. Guajacum Patavinum Obſ. & Icon. Lob.

Lupinus ſativus Dod. ſativus flore albo B. pin. ♆

Lupinus ſylveſtris flore cœruleo B. pin. ſylveſtris Dod. ♆

Lupinus flavo flore Cluſ. ſylveſtris flore luteo B. pin. ♆

Lupinus Indicus Cornuti peregrinus major, vel villoſus cœruleus major B. pin. Sativus major hort. Eyſt. ♆

* Lupinus medius cœruleus Park. ♆

Lupulus

Lupulus Dod. Lupulus mas B. pin. ☉

Luteola Obſ. & Icon. Lob. Luteola herba ſalicis folio B. pin. ☿

Lychnis hirſuta flore coccineo major B. pin. Flos Conſtantinopolitanus miniatus hort. Eyſt. Eadem quæ
Flos Conſtantinopolitanus incarnatus hort. Eyſt.
Flos Conſtantinopolitanus albus hort. Eyſt. ☉

Lychnis Chalcedonica flore pleno miniato. ☉

Lychnis coronaria Dod. coronaria flore rubro ſimplici hort. Eyſt. coronaria Dioſcoridis ſativa B. pin. ☉

Lychnis coronaria flore albo ſimplici hort. Eyſt. ☉

Lychnis coronaria flore incarnato punctato hort. Eyſt. ☉

Lychnis coronaria flore rubro pleno hort. Eyſt. ☉

Lychnis ſylveſtris alba multiplex B. pin. Ocimoïdes majus Matthioli hiſt. Lugd. ☉

Lychnis ſylveſtris alba multiplex B. pin.

ſylveſtris flore pleno albo hort. Eyſt. ☉
Lychnis ſylveſtris, ſeu aquatica purpurea ſimplex B. pin. ſylveſtris rubello flore Obſ. Lob. ☉
Lychnis ſegetum major B. pin. Pſeudomelanthium Matth. ed. B. ♀
Lychnis ſegetum rubra foliis Perfoliatæ B. pin. Vaccaria Dod. Eadem flore albo. ♀
Lychnis ſylveſtris quæ Been album vulgò B. pin Behen album, ſ. Polemonium Dod. ☉
* Lychnis, Papaver ſpumeum anguſtifolium glabrum flore purpureo. ♃ ☉
* Lychnis Papaver ſpumeum latifolium vulgare hirſutum. ♃ ☉
Lychnis ſylveſtris latifolia calyculis turgidis ſtriatis B. pin. ſylveſtris ſecunda Cluſ. Eadem flore albo. ♀
Lychnis noctiflora B. pin. Ocimoïdes noctiflora Cam. Ocimaſtrum noctiflorum flore albo hort. Eyſt. ☉
Lychnis maritima repens B. pin. marina Anglica Ad. Et Icon. Lob. ☉
Lychnis montana viſcoſa alba latifolia B. pin. ſylveſtris nona Cluſ. ☉
Lychnis ſylveſtris viſcoſa rubra anguſtifolia

folia B. pin. viſcoſa ſylveſtris flore incarnato hort. Eyſt. ♀

Lychnis ſylveſtris viſcoſa rubra altera B. pin. ſylveſtris tertia Cluſ. ♀

Lychnis viſcoſa purpurea latifolia lævis B. pin. Muſcipula flore rubro hort. Eyſt. Eadem flore albo, quæ Muſcipula flore albo hort. Eyſt. ♀

Lychnis viſcoſa flore muſcoſo B. pin. Seſamoïdes magnum Salmanticum Cluſ. ⊙

Lychnis plumaria Tab. ⊙

Lychnis ſylveſtris lanuginoſa minor B. pin. ſylveſtris hirta minima Obſ. & Icon. Lob. ♀

Lychnis, vel Ocimoïdes repens montanum B. pin. Ocimoïdes repens Polygoni folio Ad. Lob. ⊙

Lychnis Alpina glabra Caryophylli holoſtii flore. ♃ ⊙

Lychnis viſcoſa minor ſaxatilis. ♃ ⊙

Lychnis Alpina pumila folio gramineo .ſ. Muſcus Alpinus Lychnidis flore B. pin. Ocimoïdes muſcoſus Ponæ deſcr. mont. Bald. ⊙

* Lychnis flore muſcoſo capillaceo. ♃ ♀

* Lychnis maritima repens ſtrictiſſimis foliis. ♃ ⊙

* Lych-

* Lychnis minor Alpina glabra foliis longis, an Caryophylli minima ſpecies flore albo J. B. ☉
* Lychnis minor Alpina hirſuta petalis non bifidis latiori fol. ♃ ☉
* Lychnis minor Alpina hirſuta petalis non bifidis juniperinis fol. .ſ. auricula muris pulchro flore, fol. tenuiſſimo J. B. ☉

Lycium facie Pruni ſylveſtris .ſ. Italicum B. pin. Spina infectoria pumila ſecunda Cluſ.

Lycopſis B. pin. altera Anglica Obſ. & Icon. Lob. ♀
Lycopſis Monſpeliaca floribus dilutè purpureis. ♀
Lycopſis Monſpeliaca floribus albis. ♃ ♀

Lyſimachia lutea Fuchſ. lutea major quæ Dioſcoridis B. pin. ☉
Lyſimachia lutea corniculata B. pin. app. ♀
* Lyſimachia corniculata minor lutea Canadenſis. ♃ ♀
* Lyſimachia lutea flore globoſo Park. Ger. ☉

Lyſima-

Digitized by Google

Lyſimachia ſiliquoſa hirſuta magno flore B. pin purpurea Fuchſ. ☉

Lyſimachia ſiliquoſa hirſuta parvo flore B. pin. ſiliquoſa ſecunda Tab. ☉

Lyſimachia ſiliquoſa glabra major B. pin. ſiliquoſa Obſ. & Icon. Lob. ☉

Lyſimachia ſiliquoſa glabra minor B. pin. minor Tab. ☉

Lyſimachia Chamænerion dicta latifolia B. pin. Chamænerion hort. Eyſt. Onagra hiſt. Lugd. ☉

Lyſimachia Chamænerion dicta anguſtifolia B. pin. Pſeudolyſimachium purpureum minus Dod. ☉

Lyſimachia ſpicata purpurea B. pin. purpurea communis major Cluſ. ☉

Lyſimachia purpurea ſpicata lanuginoſa: ♃ ☉

Lyſimachia ſpicata cœrulea B. pin. cœruleo flore Cluſ. ☉

Lyſimachia cœrulea galericulata, vel gratiola cœrulea B. pin. galericulata Obſ. & Icon. Lob. ☉

Lyſimachia galeata minor purpurea. Gratiola latifolia Ger. ☉

M.

MAjorana vulgaris B. pin. Majorana ſ. Marum Dod. ☉

Malva ſylveſtris folio rotundo B. pin. ſylveſtris pumila Dod. ☉

Malva ſylveſtris folio ſinuato B. pin. ſylveſtris procerior Dod. Eadem flore albo. ☉

Malva foliis criſpis B. pin. criſpa Dod. ♀

Malva trimeſtris Cluſ. ♀

Malva folio vario B. prodr. ♃ ♀

Malva roſea folio ſubrotundo B. pin. hortenſis flore ſimplici rubro, incarnato, & albo hort. Eyſt. ☉

Malva roſea folio ſubrotundo flore pleno B. pin. hortenſis flore pleno rubro, atrorubente hort. Eyſt. ☉

Malva roſea multiplex flore albo incarnato hort. Eyſt. ☉

Malva hortenſis flore luteo ſimplici. Eadem flore pleno. ☉

Malva roſea folio ficus B. pin. roſea ſimplex peregrina Tab. ☉

Malva

Malva Hiſpanica. ♃ ♀
* Malva Betonicæ folio. ♃ ♀
* Malva marina arborea noſtras Park. ♀

Malus Dod. hiſt. Lugd.
Malus cydonia fructu oblongo; cydonia fœmina Rob. Steph. præd. ruſt.
Malus cydonia fructu rotundo. Cotonea Dod.
Malus medica B. pin. citria malus Cluſ.
Malus Limonia Dod. Limonia acida B. pin.
Malus aurantia Dod. arantia major B. pin.
Malus arantia humilis B. pin. Poma aurantia nana dicta hort. Eyſt.
Malus Adami B. pin. Aſſyria Dod.
Malus punica Dod. punica ſativa B. pin.
Malus punica flore pleno. Balauſtia hort. Eyſt.
Malus Perſica Dod. Perſica molli carne, & vulgaris & alba B. pin.
Malus Perſica ſucco quaſi ſanguineo B. pin.
Malus Perſica dura carne candida aliquando ex albo ſubrubente B. pin. Perſica duracina alba Tab.
Malus Armenia major hiſt. Lugd. Mala Armeniaca majora B. pin.

Malus Armenia minor hiſt. Lugd. Armeniaca minor B. pin.

Mandragora mas Fuchſ. fructu rotundo B. pin. ⊙

Mandragora fœmina Cluſ. fructu pyri B. pin. ⊙

Marum, Maſtic Gallorum, & Anglorum Ad. Lob. Sampſuchus, .ſ. Marum Maſtichen redolens B. pin. ⊙

* Marum Proſp. Alpini. ⊙

Marrubium Cardiaca dictum B. pin. Cardiaca Dod. ⊙

Marrubium aquatile Dod. paluſtre glabrum B. pin. ⊙

Marrubium nigrum fœtidum, Ballote Dioſcoridis B. pin. Ballote .ſ. Marrubium nigrum Dod. Idem flore albo. ⊙

Marrubium nigrum rotundifolium B. pin. Ocimaſtrum Valentinum Cluſ. ⊙

Marrubium nigrum longifolium B. pin. Herba venti, Parietaria Cordi hiſt. Lugd. ⊙

Marrubium album vulgare B. pin. album Obſ. & Icon. Lob. Marru-

Digitized by Google

Marrubium album villoſum B. prodr. ☉
Marrubium album latifolium peregrinum B. pin. candidum alterum Hiſpanicum Obſ. & Icon. Lob. ☉
* Marrubium aquatile foliis tenuius diſſectis. ♃ ☉
* Marrubium album anguſtifolium peregrinum C. B. P. rectum Dod. Ger. ☉

Matricaria Dod. vulgaris B. pin. ☉
Matricaria flore pleno B. pin. duplici flore Icon. Lob. ☉
Matricaria maritima B. pin. Parthenium maritimum minimum hiſt. Lugd. ☉
* Matricaria nuda nobis, bullatis foliis Park. deſcr. ☉

Melampyrum purpuraſcente coma B. pin. Melampyrum Triticum vaccinum Obſ. & Icon. Lob. ♀
Melampyrum criſtatum flore purpureo. J. B. ♀.
Melampyrum luteum latifolium B. pin. Crateogonon Obſ. & Icon. Lob. ♀
Melampyrum perpuſillum luteum Ad. Lob. luteum minimum B. pin. ♀ Vide Ranunculus.

Melilotus officinarum Germaniæ B. pin. Germanica Obſ. & Icon. Lob. ♀
Melilotus procera ſiliquis longioribus. An meliloti tertium genus Dod. deſcr. ♀
Melilotus major candida Tragi B. pin. ♀
Melilotus Italica folliculis rotundis B. pin. Italica Cam. ♀

Meliſſa Dod. hortenſis B. pin. ☉
* Meliſſa Romana hirſutior. ♃ ☉
Meliſſa peregrina folio oblongo B. pin. Moldavica flore cœruleo, albo hort. Eyſt. ♀
Meliſſa Moluccana odorata B. pin. Conſtantinopolitana Matth. ed. B. ♀
* Meliſſa Moluccana ſpinoſior Lob. Ad. ♀
Meliſſa Bætica. ♃ ♀

Melo hiſt. Lugd. vulgaris B. pin. ♀
Melo rotundus parvus B. pin. ſaccharatus Icon. Lob. ♀

Melopepo clypeiformis B. pin. Cucurbita laciniata hiſt. Lugd. ♀
Melopepo compreſſus B. pin. compreſſus alter Tab. ♀

Mentha

Mentha hortenſis corymbifera B. pin. Coſtus hortenſis hiſt. Lugd. ⊙

Mentha hortenſis verticillata Ocimi odore B. pin. Romana anguſtifolia, ſ. Mentha cardiaca Ad. & Icon. Lob. ⊙

Mentha criſpa verticillata B. pin. ſativa Dodonæi prima hiſt. Lugd. ⊙

Mentha rotundifolia criſpa ſpicata B. pin. altera Dod. ⊙

Mentha criſpa Danica, aut Germanica ſpecioſa Park. ⊙

Mentha rotundifolia ſpicata altera B. pin. Siſymbrium hortenſe Matth. ed. B. ⊙

Mentha ſpicata folio variegato B. pin. Menthaſtrum niveum Anglicum Penæ & Lobelii hiſt. Lugd. ⊙

Mentha ſylveſtris longiore folio B. pin. Menthraſtum Dod. ⊙

Mentha montana verticillata B. pin. Menthaſtrum ſecundum montanum Cluſ. ⊙

Mentha tuberoſâ radice B. pin. Menthaſtrum tertium tuberoſâ radice Cluſ. ⊙

* Menthaſtrum tuberoſâ radice flore violaceo majus. ♃ ⊙

* Menthaſtrum tuberoſâ radice flore violaceo minus. ♃ ⊙

Menthaſtrum glabrum floribus violaceis. ♃ ☉

Mentha rotundifolia paluſtris .ſ. aquatica major B. pin. Siſymbrium Dod. ☉

Mentha cataria vulgaris & major B. pin. Cattaria herba Dod. ☉

Mentha cataria minor B. prodr. ☉

* Mentha cataria folio longiori Dod. ☉

* Mentha cataria media tenuifolia Hiſpanica Cluſ. ☉

Mercurialis mas Dod. teſticulata .ſ. mas Dioſcoridis & Plinii B. pin. ♀

Mercurialis fœmina Dod. ſpicata .ſ. fœmina Dioſcoridis & Plinii B. pin. ♀

Mercurialis montana teſticulata B. pin. Cynocrambe, Mercurialis ſylveſtris Dod. ☉

Mercurialis montana ſpicata B. pin. Cynocrambe fœmina Matth. ed. B. ☉

Meſpilus Dod. Germanica folio laurino non ſerrato B. pin.

Meſpilus aronia Dod. Apii folio laciniato B. pin.

Meſpilus Apii folio ſylveſtris ſpinoſa .ſ. Oxyacantha B. pin. Spina acuta Dod.

Meum

Meum Dod. foliis Anethi B. pin. ☉
Meum adulterinum Matth. ed. B. latifolium adulterinum B. pin. ☉

Milium Dod. ſemine albo B. pin. ♀
Milium ſemine luteo B. pin. ♀
Milium ſemine nigro B. pin. nigrum Dod. ♀
Milium arundinaceum ſubrotundo ſemine Sorgo nominatum B. pin. Melica .ſ. Sorgum Dod. ♀

Millefolium vulgare album B. pin. militaris .ſ. Millefolium terreſtre vulgare Obſ. Lob. ☉
Millefolium vulgare purpureum minus B. pin. terreſtre purpureis floribus Obſ. & Icon. Lob. ☉
Millefolium rubro flore Cluſ. purpureum majus B. pin. ☉
Millefolium odoratum minus Monſpelienſium. ♃ ☉
Millefolium album foliis minus diſſectis. ♃ ☉
Millefolium tomentoſum luteum B. pin. luteum Obſ. Lob. ☉
* Millefolium Taneceti foliis flore purpureo. ♃ ☉

* Milletolium Tanaceti foliis flore albo. ♂ ☉

Mollugo Dod. montana anguſtifolia, vel Gallium album latifolium B. pin. ☉

Moly latifolium lilifforum B. pin. Moly Theophraſti Cluſ. Idem flore albo. ☉
Moly latifolium Indicum B. pin. Indicum Cluſ. ☉
Moly montanum latifolium flavo flore Cluſ. hiſt. app. alt. latifolium luteum odore Allii primum B. pin. ☉
Moly Dioſcoridis Cluſ. anguſtifolium umbellatum B. pin. ☉
Moly moſchatum capillaceo folio B. pin. & prodr. ☉
* Moly Indicum Dod. caucaphon vocatum Lob.

Morus Dod. fructu nigro B. pin.
Morus candida Dod. fructu albo B. pin.

Myagrum ſiliquâ longâ B. pin. alterum Thlaſpi effigie Obſ. Lob. ♃
Myagrum ſativum Matth. ed. B. ejuſdem pin. ♃ * Mya-

* Myagrum ſylveſtre minus B. pin. ♀
Myagrum monoſpermon latifolium B. pin. & prodr. ♀
Myagro ſimilis ſiliquâ rotundâ B. prodr. ♀
Myrrhis Dod. major, vel Cicutaria odorata B. pin. ☉
Myrrhis ſylveſtris ſeminibus lævibus B. pin. Myrrhis Fuchſ. hiſt. Lugd. ☉
* Myrrhis perennis alba fol. hirſutis. ♃ ☉
* Myrrhis perennis alba foliis hirſutioribus. ♃ ☉
* Myrrhis perennis alba foliis hirſutiſſimis. ♃ ☉
* Myrrhis perennis lutea Daucoïdes. ♃ ♀
* Myrrhis annua nodoſa ſemine aſpero. ♃ ♀
Myrrhis ſylveſtris ſeminibus aſperis B. pin. ſylveſtris nova Æquicclorum Col. ♀
Myrtus latifolia Romana B. pin. Romana Matth. ed. B. ☉
Myrtus flore pleno Cornuti. ☉
Myrtus minor vulgaris B. pin. Tarentina hiſt. Lugd. ☉
* Myrtus latifolia maxima Park. Cluſii hiſt. fol. confertim naſcentibus Laurinis.

N.

NAPUS Dod. ſativa B. pin. ♀

Narciſſus mediopurpureus Dod. albus circulo purpureo B. pin. ⊙

Narciſſus Orientalis medius totus albus polyanthos hort. Eyſt. Orientalis medius albus B. pin. ⊙

Narciſſus totus albus major B. pin. Orientalis polyanthos minor totus albus hort. Eyſt. ⊙

Narciſſus medioluteus Dod. pallidus circulo luteo B. pin. ⊙

Narciſſus medioluteus Donas Narbonenſium Icon. Lob. medioluteus copioſo flore odore gravi B. pin. ⊙

Narciſſus de Argiers polyanthos totus luteus hort. Eyſt. luteus polyanthos Luſitanicus B. pin. ⊙

Narciſſus latifolius flore prorſus flavo Cluſ. luteus Conſtantinopolitanus B. pin. ⊙

Narciſſus anguſtifolius flavus magno calice B. pin. juncifolius primus amplo calice Cluſ. ⊙

Nar-

Narciſſus juncifolius major Dod. juncifolius oblongo calice luteus major B. pin. ⊙

Narciſſus juncifolius flore rotundo roſeo B. pin. juncifolius flore rotundo circinato Swert. ⊙

Narciſſus maximus pallidus amplo calice flavo Cluſ. cur. poſt. deſcr. latifolius pallidus calice amplo alter B. pin. ⊙

Narciſſus latifolius ſulphureus, vel albus brevi calice B. pin. ſulphurei coloris brevi calice Swert. ⊙

Narciſſus luteus ſylveſtris Dod. ſylveſtris pallidus calice luteo B. pin. ⊙

Narciſſus major totus luteus calice prælongo B. pin. Pſeudonarciſſus major Hiſpanicus Cluſ. ⊙

Narciſſus totus luteus florum foliis reflexis B. pin. totus luteus oblongo calice, & reflexis foliis hort. Eyſt. ⊙

Narciſſus oblongâ tubâ totus albus Swert. albus oblongo calice B. pin. ⊙

Narciſſus montanus alter flore fimbriato B. pin. montanus juncifolius minimus alter flore luteo fimbriato Mutoni Icon. Lob. ⊙

Narciſſus nobilis Orientalis polyanthos mixtus hort. Eyſt. Orientalis flore multiplici, & pleno albo luteo B. pin. ⊙

Narciſſus incomparabilis flore pleno partim flavo, partim croceo. ⊙

Narciſſus exoticus .ſ. Conſtantinopolitanus Cam. major flore multiplici hort. Eyſt. latifolius albus medioluteus B. pin. ⊙

Narciſſus totus niveus multiplex B. pin. albus pleno flore Swert. ⊙

Narciſſus ſylveſtris ſtellatus hort. Eyſt. ſylveſtris multiplex calice carens B. pin. ⊙

Narciſſus Septentrionalis flore pleno luteo hort. Eyſt. flore pleno variegato B. pin. ⊙

Narciſſus juncifolius luteus flore pleno hort. Eyſt. juncifolius aureus multiplex Anemones formâ B. pin. ⊙

Narciſſus maritimus B. pin. Pancratium hiſt. Lugd. ⊙

Narciſſus Conſtantinopolitanus tertius Mat. ed. B. Illyricus Liliaceus B. pin. ⊙

* Narciſſus Indicus puniceus gemino latiore folio Bap. Ferr. ⊙

* Narciſſus Indicus ſphæricus flore Liliaceo Bap. Ferr.

Nardus

Nardus montana radice olivari B.pin. montana Matth. ed. B. ☉
Nardus montana radice oblongâ B. pin. montana altera Matth. ed. B. ☉

Naſturtium hortenſe vulgatum B. pin. hortenſe Dod. ☿
Naſturtium hortenſe criſpum B. prodr. ☿
Naſturtium pratenſe magno flore B. pin. Cardamine altera ſimplici flore Cluſ. ☉
Naſturtium pratenſe flore pleno B.pin. Cardamine altera flore pleno Cluſ. ☉
Naſturtium pratenſe floſculis minimis. ♃ ☿
Naſturtium aquaticum ſupinum B. pin. aquaticum Dod. ☉
Naſturtium aquaticum minus B. pin. Cardamine quarta Dal. hiſt. Lugd. ☉
Naſturtium Alpinum trifolium B. pin. Cardamine Alpina trifolia Cluſ. ☉
Naſturtium Alpinum Bellidis folio minus B. pin. Plantula Cardamines alterius æmula Cluſ. ☉
Naſturtium ſylveſtre tenuiſſimè diviſum B. pin. Sophia Chirurgorum Obſ. & Icon. Lob. ☿
Naſturtium Indicum Dod. Indicum majus B. pin. ☿

* Naſturtium hortenſe latifolium B. prodr. ♀
* Naſturtium Alpinum inſipidum Ger. Tab. Cardamine Alpina inſipida Col. ♀
* Naſturtium ſylveſtre minus B. prodr. ♀

Nerion floribus rubeſcentibus B. pin. flore rubro hort. Eyſt.
Nerion floribus albis B. pin. Nerion .ſ. Rhododendron flore albo hort. Eyſt.

Nicotiana major latifolia B. pin. Nicotiana .ſ. Tabacum hiſt. Lugd. ♀
Nicotiana major anguſtifolia hort. Eyſt. B. pin. ♀
Nicotiana minor B. pin. minor .ſ. Hyoſcyamus luteus hort. Eyſt. ♀

Nigella arvenſis cornuta B. pin. Melanthium ſylveſtre Dod. ♀
Nigella latifolia flore majore ſimplici cœruleo B. pin. Melanthium Hiſpanicum majus hort. Eyſt. ♀
Nigella flore majore pleno cœruleo B. pin. Melanthium Damaſcenum flore pleno hort. Eyſt. ♀

Nigella

Nigella flore minore pleno & albo B. pin. citrina Ad. Lob. deſcr. Eadem flore ſimplici ſemine nigro. ♀

* Nigella ſylveſtris flore luteo pallido. ♃ ♀

Nummularia Dod. major lutea B. pin. ☉

Nummularia flore purpuraſcente B. prodr. ☉

Nux juglans Dod. Nux juglans .ſ. Regia vulgaris B. pin.

* Nux juglans putamine fragili & fructu tenero C. B. P.

* Nymphæa maj. alba C. B. Pin. alba Matth. Dod. candida Fuchſ.

* Nymphæa maj. lutea C.B.P. lutea Dod. Fuchſ.

* Nymphæa minima alba C. B. P. morſus Ranæ Dod. Lob. Icon.

O.

OChrus folio integro capreolos emittente B. pin. Ervilia ſylveſtris Dod. ♆

Ocimum caryophyllatum maximum B. pin. latifolium magnum hort. Eyſt. ♆
Ocimum Citri odore B. pin. Citratum .ſ. Citri odore hort. Eyſt. ♆
Ocymum vulgatius B. pin. Baſilicum .ſ. Ocimum medium vulgatius hort. Eyſt. ♆
Ocimum medium foliis ex nigro vireſcentibus flore albo & violaceo. ♆
Ocimum Aniſi odore B. pin. hort. Eyſt. ♆
Ocimum minimum B. pin. minimum caryophyllatum hort. Eyſt. ♆
Ocimum minimum foliis rubeſcentibus. ♋ ♆
* Ocimum rubrum pleno flore. ♋ ♆

Oenanthe Apii folio B. pin. ſecunda Matth. ed. B. hiſt. Lugd. ☉
Oenanthe anguſtifolia Obſ. & Icon. Lob. ☉

Oenanthe

Oenanthe aquatica B. pin. Juncus odoratus aquatilis Dod. ☉

* Oenanthe Chærophylli foliis C.B.P. Cicutæ facie Ad. Lob. Succo viroſo cicutæ facie Lobelio J. B. ☉

Olea Dod. ſativa B. pin.

Olea ſylveſtris folio duro ſubtus incano B. pin. ſylveſtris Cluſ.

Olea Bohemica .ſ. Elæagnos Matth. ed. B. ſylveſtris folio molli incano B. pin.

Onobrychis Dod. foliis Viciæ, fructu echinato major B. pin. Caput gallinaceum Belgarum Obſ. & Icon. Lob. ☉

Onobrychis fructu echinato minor B. pin. Caput gallinaceum minus ejuſdem prod. deſcr. ☉

* Onobrychis flore ſaturate rubro ☉

Onobrychis ſemine clypeato aſpero major B. pin. Hedyſarum clypeatum flore ſuaviter rubente hort. Eyſt. ♀

Onobrychis arvenſis, vel Campanula arvenſis erecta B. pin. Viola pentagonia, Pentagonium Tab. ♀

* Onobrychis viola pentagonia perfoliata. ♃ ♀

Digitized by Google

Onobrychis viola pentagonia latiori folio oblongo. ♃ ♀

Ophioglossum Dod. vulgatum B. pin. ☉

Ophrys Plin. Ophris Fuchs. bifolia B. pin. ☉

Orchis morio mas foliis maculatis B. pin. Testiculus morionis mas Dod. ☉

Orchis morio fœmina B. pin. Testiculus morionis fœmina Dod. Eadem flore roseo & albo. ☉

Orchis flore simiam referens B. pin. altera oreades cercopithecophora Col. ☉

Orchis odore hirci longiore, angustioréque folio B. pin. saurodes, vel scincophora lacertarum æmulatione Gemmæ Obs. & Icon. Lob. ☉

Orchis hermaphroditica secunda Gemmæ s. psychodes diphylla Obs. & Icon. Lob. bifolia altera B. pin. ☉

Orchis hermaphroditica flore flavescente. Forte, Serapias batrachites vel myodes Icon. Lob. ☉

Orchis s. Testiculus minor violaceis floribus Cord. fucum referens major foliolis superioribus

perioribus candidis & purpuraſcentibus B. pin. ⊙

Orchis melitias, .ſ. Apis cadaverulum exprimens Obſ. & Icon. Lob. fucum referens flore ſubvirente B. pin. ⊙

Orchis muſcæ corpus referens minor, vel galeâ & alis herbidis B. pin. Orchis Serapias tertius Dod. ⊙

Orchis ornithopora, vel ornithes folio lævi Obſ. & Icon. Lob. papilionem expanſum herbacei coloris referens B. pin. ⊙

Orchis ſpiralis major. Tetrorchis, vel Triorchis alba Gemmæ hiſt. Lugd. Triorchis, vel Tetrorchis alba odorata major B. pin. ⊙

Orchis ſpiralis minor .ſ. Teſticulus odoratus Obſ. & Icon. Lob. Triorchis alba odorata minor B. pin. ⊙

Orchis palmata pratenſis latifolia longis calcaribus B. pin. Satyrium baſilicum mas Dod. ⊙

Orchis palmata pratenſis maculata B. pin. Satyrium baſilicum fœmina Dod. Eadem flore candido Dod. deſcr. ⊙

Orchis palmata minor calcaribus oblongis B. pin. Palma Chriſti minor mas Tab. Eadem flore albo. ⊙

Orchis palmata paluſtris latifolia B. pin. Satyrium baſilicum folioſum Dod. ⊙

* Orchis hermaphroditica flore luteo pallido. ⊙

* Orchis palmata flore Fuſco. ♃ ⊙

Origanum ſylveſtre .ſ. vulgare Dod. ſylveſtre, Cunila bubula Plinii B. pin. Idem flore albo, quod

Origanum album Tab. ſylveſtre album B. pin. ⊙

Origanum heracleoticum Matth. ed. B. hiſt. Lugd. heracleoticum, Cunila gallinacea Plinii B. pin. Idem flore albo. ⊙

Origanum fiſtuloſum Canadenſe Cornuti. ⊙

Origano cognata Zatarendi B. pin. Zatar hendi. Origanum Indicum Veſl. Obſ. in Proſp. Alp. ⊙

* Origanum ſylveſtre humile B. prodr. ⊙

Ornithogalum umbellatum medium anguſtifolium B. pin. Bulbus leucanthemus minor .ſ. Ornithogalum Dod. ⊙

Ornithogalum anguſtifolium majus floribus ex albo vireſcentibus B. pin. Aſphodelus bulboſus Galeni Dod. ⊙

Ornithogalum

Ornithogalum ſpicatum flore viridi lacteſcente B. pin. pannonicum albo flore Cluſ. ☉

Ornithogalum Neapolitanum Cluſ.app.alt. exoticum magno flore, minore innato B. pin. ☉

* Ornithogalum Arrabicum Cluſ. Eyſt. ☉

Ornithogalum anguſtifolium bulbiferum B. pin. anguſtifolium campeſtre luteum minimum bulbiferum Col. ☉

Ornithopodium Dod. majus B. pin. ♀

* Ornithopodium πολυκέφαλον nobis, ſcorpioïdes leguminoſa Lob. ♀

* Ornithopodium διφάλον minus nobis, ſcorpioïdes altera leguminoſa J. B. ♀

* Orobanche planta clandeſtina Lugd. app. ☉

* Orobanche radice dentata maj C. B. P. Dentaria ἄφυλλος 3 Cluſ. ☉

* Orobanche radice dentata minor C.B.P. Dentaria ἄφυλλος 3 Cluſ. ☉

Orobus ſiliquis articulatis ſemine majore B. pin. receptus herbariorum Obſ. & Icon. Lob. ♀

Orobus pannonicus primus Cluſ. Sylvaticus purpureus vernus B. pin. ⊙

Orobus ſylvaticus foliis Viciæ B. pin. Aſtragaloïdes altera herbariorum Icon. Lob. ⊙

Oxyacantha Dioſcoridis .ſ. Spina acuta pyri folio B. pin. Pyracantha Ad. Lob.

P.

PÆONIA mas Dod. folio nucis Geſn. hort. folio nigricante ſplendido quæ mas B. pin. ⊙

Pæonia communis vel fœmina B. pin. rubra flore ſimplici hort. Eyſt. ⊙

Pæonia promiſcua .ſ. neutra Obſ. Lob. fœmina altera B. pin. ⊙

Pæonia folio ſubtus incano flore albo, vel pallido B. pin. mas flore albo hort. Eyſt. ⊙

Pæonia polyanthos flore rubro hort. Eyſt. fœmina flore pleno rubro majore B. pin. ⊙

Pæonia

Digitized by Google

Pæonia pleno flore albeſcente, minor Cluſ. flore exalbido pleno, major & minor B. pin. ☉

Palma minor B. pin. Camæriphes Dod. ☉

Panaces carpimon .ſ. racemoſa Canadenſis Cornuti. ☉

Panax Paſtinacæ folio, An Syriacum Theophraſti B. pin. Panaces peregrinum Dod. ☉

Panicum Dod. Germanicum, .ſ. paniculâ minore B. pin. ♀

Panicum Italicum, .ſ. paniculâ majore B. pin. ſativum hiſt. Lugd. ♀

Papaver hortenſe ſemine albo B. pin. album ſativum Icon. Lob. ♀

Papaver hortenſe nigro ſemine B. pin. nigrum ſativum Dod. ♀

Papaver criſtatum floribus & ſemine album B. pin. fimbriatum, aut criſtatum album Icon. Lob. ♀

Papaver criſtatum floribus rubris ſemine nigro B. pin. ſativum rubrum fimbriatum Dal. hiſt. Lugd. ♀

Papaver laciniatum rubrum unguibus purpureis hort. Eyſt.

Papaver hortenſe flore ſimplici dilutiſſimè violaceo. ♀

Papaver hortenſe flore ſimplici placidè rubro unguibus plumbeis. ♀

Papaver flore multiplici purpuraſcente hort. Eyſt. ♀

Papaver flore miniato pleno hort. Eyſt. ♀

Papaver flore pleno violaceo. ♀

Papaver flore pleno, argentei coloris hort. Eyſt. ♀

Papaver flore pleno album B. pin. polyanthos Tab. ♀

Papaver flore pleno incarnato hort. Eyſt ♀

Papaver flore pleno rubro hort. Eyſt. ♀

Papaver flore pleno dilutè violaceo. ♀

Papaver erraticum majus B. pin. erraticum Dod. ♀

Papaver erraticum flore pleno B. pin. Idem flore pleno igneo, flore pleno rubro, & criſpo. ♀

* Papaver erraticum Pyrennium flore flavo C.B.P. Argemone Cambrobritannica Park. ⊙

Papaver corniculatum luteum B. pin. corniculatum majus Dod. ⊙ * Pa-

*Papaver corniculatum phœniceum hirſutum C. B. flore phœniceo Cluſ. ♀
*Papaver corniculatum violaceum C. B. Lob. ♀
Papaver ſpinoſum B. prodr. ♀

Parietaria Dod. officinarum, & Dioſcoridis B. pin. ☉

Paſtinaca tenuifolia ſylveſtris Dioſcoridis, vel Daucus officinarum B. pin. Daucus vulgaris Cluſ. ♀
Paſtinaca ſylveſtris hirſuta echinato fructu B. pin. Daucus λαπόκαρυλος Thal. ♀
*Paſtinaca tenuifolia lucida. ♂ ♀
Paſtinaca tenuifolia ſativa, radice lutea, vel alba B. pin. tenuifolia ſativa Dod. ♀
Paſtinaca tenuifolia ſativa, radice atrorubente B. pin. ſativa rubens Dod. ♀
Paſtinaca tenuifolia radice & umbellâ luteâ. ♂ ♀
Paſtinaca ſativa latifolia Dod. B. pin. ♀
Paſtinaca ſylveſtris latifolia Dod. Elaphoboſcum Matth. ed. B. ♀

Pedicularis Alpina Filicis folio minor B. pin.

pin. Alectorolophos ſecunda Cluſ. deſcr. ☉

Pedicularis pratenſis purpurea B. pin. Fiſtularia Dod. ♀

Pedicularis pratenſis lutea, vel criſta Galli B. pin. Criſta Galli Dod. ♀

*Pentaphylloïdes fragariæ folio. ♃ ☉

Peplus Dod. Peplus .ſ. Eſula rotunda B. pin. ♀

* Peplis Cluſ. hiſt. maritima folio obtuſo C. B. P. ♀

Pepo major oblongus Dod. oblongus B. pin. ♀

Pepo rotundus major Dod. compreſſus major B. pin. ♀

Pepo rotundus minor Dod. compreſſus minor B. pin. ♀

Pepo rotundus Melonis effigie B. pin. Indicus minor rotundus Tab. ♀

Pepo rotundus Aurantii formâ B. pin. Cucurbita Aurantii formâ hort. Eyſt. ♀

Pepo minor clypeatus B. pin. Indicus minor clypeatus Tab. ♀

Pepo minor anguloſus B. pin. Indicus minor anguloſus Tab. ♀

Perfoliata Dod. vulgatiſſima .ſ. arvenſis B. pin. ☉

Perfoliata montana Cam. montana latifolia B. pin. ☉

Perfoliata minor anguſtifolia Bupleuri folio B. pin. minima Bupleuri folio Col. deſcr. ♀

Periclymenum non perfoliatum Septentrionalium Obſ. & Icon. Lob. non perfoliatum Germanicum B. pin. ☉

Periclymenum perfoliatum ſerotinum flore rubro. ☉

Periclymenum perfoliatum B. pin. perfoliatum calidarum regionum Obſ. & Icon. Lob. ☉

Perſicaria Dod. mitis maculoſa, & non maculoſa B. pin. Eadem flore albo. ♀

Perſicaria urens .ſ. hydropiper B. pin. Hydropiperi Dod. ♀

Perſicaria puſilla repens Obſ. & Icon. Lob. minor B. pin. ♀

Peta-

Petaſites Dod. major & vulgaris B. pin. ☉
Petaſites flore albo hort. Eyſt. minor B. pin. ☉
Petaſites minor folio Tuſſilaginis. ♃ ☉

Peucedanum majus Italicum Lobelii hiſt. Lugd. B. pin. ☉
Peucedanum Dod. Germanicum B. pin. ☉
* Peucedanum minus fol. Lobatis anguſtis. ♃ ☉
* Peucedanum minus C. B. Pin. ſelinum montanum pumilum Cluſ.

Phalangium parvo flore non ramoſum B. pin. non ramoſum Dod. ☉
Phalangium parvo flore ramoſum B. pin. ramoſum Dod. ☉
* Phalangium Allobrogicum. ☉
Phalangium Virginianum Tradeſcanti Ger. Idem flore albo. ☉
* Phalangium Virginianum amplo flore rubro. ☉
* Phalangium Virginianum amplo flore cœruleo ſ. Azureo. ☉

Phalaris Dod. major ſemine albo B. pin. ♀
* Phalaris

* Phalaris minore ſemine nigro. ♃ ☿

Phaſeolus vulgaris Obſ. & Icon. Lob. Dolichus ſ. Phaſeolus Dod. fructu albo, cœruleo, nigro, flavo, obſoleto, amethyſtino, cineraceo, albo nigris lituris diſtincto, purpureo nigris lituris variegato. ☿

Phaſeolus flore puniceo Cornuti. Idem fructu albo, & variegato. ☿

Phaſeolus minimus fructu viridi ovato. ♃ ☿

* Phaſeolus minor Coloſſinus ſiliquis longiſſimis. ♃ ☿

* Phaſeolus minor albus, macula nigra umbilico notato J. B. ☿

Phillyrea Cluſ. latifolia ſpinoſa B. pin.

Phillyrea folio leviter ſerrato B. pin. ſecunda Cluſ.

Phillyrea folio Liguſtri B. pin. tertia Cluſ.

Phillyrea anguſtifolia prima B. pin. quarta Cluſ.

Phyllum marificum Cluſ. teſticulatum B. pin. ☉

Phyllum

Phyllum fœminificum Cluſ. ſpicatum B. pin. ⊙

Picea hiſt. Lugd. major prima, ſ. Abies rubra B. pin.

Piloſella major Dod. major repens hirſuta B. pin. ⊙
Eadem folio variegato. ⊙
Piloſella major repens minus hirſuta B. pin.
Lactucella ſylveſtris repens Cam. deſcr. ⊙
Piloſella Indica Cornuti deſcr. Hieracium peregrinum hort. Eyſt. ⊙
* Piloſella umbelifera Macrocaulos Col. ⊙
* Piloſella major glabra. ♃ ⊙

Pimpinella, Saxifraga major umbellâ candidâ B. pin. major Fuchſ. ⊙
Pimpinella, Saxifraga major altera B. pin. minor Fuchſ. ⊙
Pimpinella, Saxifraga minor B. pin. criſpa ſ. petræa Tab. ⊙
Pimpinella ſylveſtris, Sanguiſorba major Dod. Pimpinella, Sanguiſorba major B. pin. ⊙

Pimpinella maxima Canadenſis Cornuti. ☉
Pimpinella maxima Canadenſis ſpicâ longâ rubrâ. ♃ ☉
Pimpinella maxima Canadenſis ſpicâ rubrâ glomeratâ. ♃ ☉
Pimpinella, Sanguiſorba minor hirſuta B. pin. Pimpinella, Sanguiſorba Dod. ☉
Pimpinella, Sanguiſorba minor lævis B. pin. ☉
Pimpinella agrimonoïdes. ♃ ☉

Pinus domeſtica Matth. ed. B. ſativa B. pin.
Pinus ſylveſtris montana Matth. ed. B. ſylveſtris B. pin.

Piper Indicum vulgatiſſimum B. pin. Indicum medium hort. Eyſt. Capſicum, Piper Indicum Obſ Lob. ☿
Piper Indicum propendentibus ſiliquis oblongis recurvis B. pin. Capſicum recurvis ſiliquis Dod. ☿
Piper Indicum ſiliquis flavis hort. Eyſt. ☿
Piper Indicum propendentibus ſiliquis rotundis B. pin. Indicum cordatum hort. Eyſt. ☿
Piper Indicum ſiliquis ſurrectis, & oblongis

B. pin. Indicum medium longum erectum hort. Eyst. Idem fructu flavo. ♀

Piper Indicum minimum erectum hort. Eyst. ♀

Piper Indicum siliquis surrectis Cerasi formâ B. pin. Capsicum minimis siliquis Dod. ♀

Pistacia Matth. ed. B. peregrina fructu racemoso .s. Terebinthus Indica Theophrasti B. pin.

Pistacia sylvestris B. pin. Staphylodendron Matth. ed. B. ☉

Pistacia Virginea sylvestris trifolia. ♃ ☉

Pisum majus quadratum B. pin. majus Dod. ♀

Pisum umbellatum B. pin. proliferum Tab. ♀

Pisum hortense majus B. pin. ramulare hist. Lugd. Idem fructu rubro variegato B. pin. descr. ♀

Pisum arvense B. pin. minus Dod. ♀

Pisa sine cortice duriore B. pin. leptoloba Cam. ♀

Pisa nana rotunda. ♃ ♀

* Piſum ſylveſtre Viciæ fructu. ♃ ♀
Piſum repens perenne, aliud maritimum Britannicum Park. ☉
Piſum veſicarium fructu nigro alba macula notato B. pin. cordatum Ad. & Icon. Lob. ♀

Plantago maxima tota glabra B. prodr. maxima Hiſpanica Cornuti. ☉
Plantago latifolia ſinuata B. pin. major Dod. ☉
Plantago latifolia incana B. pin. media Dod. ☉
Plantago latifolia roſea flore expanſo B. pin. roſea hort. Eyſt. ☉
Plantago latifolia roſea floribus quaſi in ſpica diſpoſitis B. pin. roſea ſpicata hort. Eyſt. ☉
Plantago latifolia incana ſpicis variis B. pin: peregrina latifolia Gareti Cluſ. ☉
Plantago quinquinervia Obſ. & Icon. Lob. anguſtifolia major B. pin. ☉
* Plantago marina, quinquenervia incana ♃ ☉
Plantago aquatica latifolia B. pin. aquatica Dod. ☉

Plantago aquatica anguſtifolia B. pin. aquatica humilis anguſtifolia,& longifolia Icon. Lob. ☉

Plantago aquatica ſtellata B. pin. Damaſonium ſtellatum hiſt. Lugd. ♀

Platanus Dod. Cluſ.

* Platanus Virginiana Park. ☉

Polium Lavendulæ folio B. pin. recentiorum fœmina Lavendulæ folio Obſ. & Icon. Lob. ☉

Polium montanum luteum B. pin. luteum Dal. hiſt. Lugd. ☉

Polium montanum album B. pin. montanum primum Cluſ. ☉

Polium montanum ſecundum flore albo Cluſ. deſcr. ☉

* Polium rubrum majus flore dilutiori. ♃ ☉

* Polium rubrum minus, montanum 5 purpureum Cluſ. hiſt. ☉

Polygala vulgaris B. pin. Flos ambarvalis Dod. Eadem quæ

Polygala cœrulea Tab.

Polygala alba Tab. ♀

Polygala repens nivea B.pin. repens nuperorum Obſ. & Icon. Lob. ♀
Polygala Valentina Cluſ. altera B. pin. ⊙

Polygonatum Dod. latifolium vulgare B. pin ⊙
Polygonatum latifolium flore majore odoro B. pin. latifolium Fuchſ. Dod. ⊙
Polygonatum latifolium hellebori albi foliis B. pin. tertium latiore folio Cluſ. deſcr. ⊙
Polygonatum latifolium ramoſum B. pin. ramoſum Dod. ⊙
Polygonatum anguſtifolium non ramoſum B. pin. anguſtifolium Fuchſ. hiſt. Lugd. ⊙
Polygonatum ſpicatum ſterile Cornuti. ⊙
Polygonatum ramoſum flore luteo majus Cornuti. ⊙

Polygonum latifolium B. pin. Polygonum mas Dod. ♀
Polygonum maritimum latifolium B. pin. marinum prius Dal. hiſt. Lugd. ♀
Polygonum floſculis herbaceis ex alis proeuntibus. Polycarpum hiſt. Lugd.. ♀

Polygonum floſculis albis in ſummitate. Saxifraga Anglica Ad. Lob. ♀

Polygonum Caryophylli holoſtii flore. ♂ ♀

Polygonum paluſtre alſinefolium. Saxifraga paluſtris alſinefolia Ger. ♀

Polygonum montanum minimum niveum Obſ. Lob. minus candicans B. pin. ♀

Polygonum littoreum minus floſculis ſpadiceo-albicantibus B. prodr. deſcr. ♀

Polygonum minus .ſ. Millegrana major B. pin. Herniaria Dod. ♀

Polygonum minus .ſ. Millegrana major foliis glabris B. pin. deſcr. ♀

Polygonum minus maritimum incanum. ♂ ♀

Polygonum minimum .ſ. Millegrana minima B. pin. Millegrana minima Icon. Lob. ♀

* Polygonum marinum minus Serpilli fol. C. B. P. ♀

Polypodium majus Dod. vulgare B. pin. ☉

Populus alba Dod. B. pin.

Populus nigra Dod. B pin.

Populus tremula B. pin. Lybica Dod.

Forrum

Porrum commune capitatum B. pin. capitatum Fuchſ. ♀

Porrum folio latiſſimo B. pin. Syriacum Tab. ♀

* Porrum ſylveſtre Vinearum C.B.P. Ampelopraſum Lugd. ♀

Porrum ſectivum juncifolium B. pin. Sehœnopraſum Dod. ☉

Portulaca hortenſis Fuchſ. latifolia .ſ. ſativa B. pin. ♀

Portulaca hortenſis folio flavo. ♃ ♀

Portulaca ſylveſtris Dod. anguſtifolia .ſ. ſylveſtris B. pin. ♀

* Portulaca minima alba. ♃ ♀

* Potamogeton ſalicis fol. C. B. Pin. potamogeïton Dod. ☉

Potentilla Fuchſ. Argentina .ſ. Potentilla Dod. ☉

Poterion Obſ. & Icon. Lob. Poterio affinis, foliis Pimpinellæ, ſpinoſa B. pin. ☉

Prunus Dod.

Prunus fructu rotundo nigropurpureo dulci B. pin. Prunus myrobalanus Cluſ.
Prunus ſylveſtris Obſ. & Icon. Lob. B. pin.

Pſeudodictamnus verticillatus inodorus B. pin. Pſeudodictamnum Dod. ☉

Pſyllium Plinianum forte, radice perenni ſupinum Obſ & Icon. Lob. majus ſupinum B. pin. ☉
Pſyllium Dod. majus erectum B. pin. ♀

* Ptarmica Virginiana folio Helenii. ♃ ☉
* Ptarmica foliis ſcabroſæ B. prodr. ☉
Vide dracunculus.

Pulegium latifolium B. pin. regium hiſt. Lugd. Idem flore albo. ☉
Pulegium anguſtifolium B. pin. cervinum hiſt. Lugd. Idem flore albo. ☉
* Pulegium denſi ſaraclis Park. ☉

Pulſatilla Dod. folio craſſiore, & majore flore B. pin. ☉
Pulſatilla flore albo B pin, albo Lobelii hiſt. Lugd. ☉

Py-

Pyrola Dod. rotundifolia major B. pin. ⊙

Pyrus Dod. ſativa B. pin.
Pyrus ſylveſtris B. pin. Pyra ſylveſtria Tab.

Q.

* QUADRIFOLIUM albo flore phæum Fuchſ. Lugd. ⊙ A luxuriante natura à Trifolio in quadrifolium maculatum, degenerans.

Quercus cum longo pediculo B. pin. Hemeris etymodrys hiſt. Lugd.
* Quercus vulgaris brevi pediculo.
* Quercus, incanis ſubtûs foliis, pediculo brevi.
* Quercus incanis ſubtûs foliis, longo pediculo.
* Quercus calyce echinato, glande majore.

Quinquefolium majus repens B. pin. majus luteum Fuchſ. ⊙

Quinquefolium minus repens lanuginoſum luteum B. pin. Pentaphyllum .ſ. Quinquefolium minus Tab. ⊙

Quinquefolium folio argenteo B. pin. erectum incanum Matth. ed. B. ⊙

Quinquefolium rectum luteum B. pin. Pentaphyllum majus Obſ. Lob. ⊙

Quinquefolium montanum erectum hirſutum luteum B. prodr. ⊙

* Quinquefolium ſiliquoſum Proſp. Alp ♀

* Quinquefolium Alpinum repens ſerotinum. ♃ ⊙

* Quinquefolium Canadenſe erectum. ♃ ⊙

Quinquefolium fragiferum B. pin. fragiferum quintum Cluſ. ⊙

Quinquefolium fragiferum repens Tab. Pentaphyllum ſupinum Potentillæ facie Cluſ. Quinquefolio fragifero affinis B. pin. ⊙

Quinquefolium paluſtre rubrum B. pin. rubrum Dodonæi hiſt. Lugd. ⊙

R.

RANUNCULUS ſylvarum Cluſ. nemoroſus Obſ. Lob. Anemone nemoroſa flore majore B. pin. Idem flore roſeo. ⊙

* Ranunculus nemoroſus flore pleno albo. ⊙

Ranunculus nemoroſus luteus B. pin. Icon. Lob. ⊙

Ranunculus auricomus dulcis Tragi. Icon. Lob. nemoroſus vel ſylvaticus folio rotundo B. pin. ⊙

Ranunculus nemoroſus Thalictri folio B. pin. præcox ſecundus Thalictri folio Cluſ. ⊙

Ranunculus Thalictri folio minor Aſphodeli radicibus Park. Oenanthe Myconi hiſt. Lugd. malè à B. pin. Oenanthe hederæ foliis. ⊙

Ranunculus nemoroſus Muſcatellina dictus B. pin. Moſchatella Cord. hiſt. ⊙

Ranunculus pratenſis ſurrectis cauliculis Obſ. & Icon. Lob. pratenſis erectus acris B. pin. ⊙

Ranunculus pratenſis erectus dulcis B. pin. hortenſis erectus flore ſimplici luteo hort. Eyſt. ☉

Ranunculus pratenſis reptante cauliculo Obſ. & Icon. Lob. pratenſis repens hirſutus B. pin. ☉

Ranunculus pratenſis repens hirſutus maculatus. ♃ ☉

Ranunculus bulboſus Obſ. & Icon. Lob. pratenſis radice verticilli modo rotundâ B. pin. ☉

Ranunculus hortenſis erectus flore pleno B. pin. erectus hortenſis flore pleno luteo hort. Eyſt. ☉

Ranunculus pratenſis reptante cauliculo flore pleno. An Ranunculus hortenſis inclinans B. pin. ☉

Ranunculus dulcis flore pleno B. pin. dulcis multiflorus Tab. ☉

Ranunculus tuberoſus flore multiplici Dod. radice tuberosâ flore pleno, & prolifero B. pin. ☉

Ranunculus folio Aconiti flore albo multiplici B. pin. niveus polyanthes Lobelii hiſt. Lugd. ☉

Ranunculus arvenſis echinatus B. pin. arvorum Icon. Lob. ☿

Ranunculus Creticus echinatus latifolius Prosp. Alp. de exot. Stellatus echinatus Creticus B. pin. ♀

Ranunculus paluſtris Apii folio lævis B. pin. paluſtris rotundiori folio Obſ. & Icon. Lob. ♀

Ranunculus longifolius paluſtris major B. pin. lanceatus major Tab. ⊙

Ranunculus longifolius paluſtris minor B. pin. flammeus aquaticus anguſtifolius Obſ. & Icon. Lob. ⊙

Ranunculus paluſtris ſerratus B. pin. aquatilis anguſtifolius ſerratus Obſ. & Icon. Lob. ⊙

Ranunculus montanus folio gramineo B. pin. pumilus gramineis foliis Icon. Lob. ⊙

Ranunculus Illyricus major Cluſ. deſcr. lanuginoſus anguſtifolius grumoſa radice major B. pin. ⊙

Ranunculus Illyricus Dod. lanuginoſus anguſtifolius grumoſa radice minor B. pin. ⊙

Ranunculus lanuginoſus Apii folio, Aſphodeli radice B. pin. alter ſaxatilis Aſphodeli radice Col. Idem flore pleno. ⊙

Ranunculus Aſiaticus grumoſa radice flore phœniceo ruberrimo hort. Eyſt. Idem flore flavo, ſubalbido. ⊙

Ranunculus Creticus grumoſa radice flore niveo hort. Eyſt. ⊙

Ranunculus creticus latifolius Cluſ. Aſphodeli radice Creticus B. pin. ⊙

Ranunculus Aſiaticus grumoſa radice flore flavo rubris venis diſtincto Cluſ. cur. poſt. ⊙

Ranunculus Aſiaticus flore albo colore ſuaviter rubente per oras mixto B. pin. deſcr. ⊙

Ranunculus Aſiaticus grumoſa radice, flore pleno Cluſ. Aſphodeli radice flore pleno ſanguineo B. pin. ⊙

Ranunculus flore pleno ruberrimo hort. Eyſt. Aſphodeli radice flore ſubphœniceo rubente B. pin. ⊙

Ranunculus Aſiaticus grumoſa radice, flore pleno ſanguineo prolifero hort Eyſt. Aſphodeli radice prolifer miniatus B. pin. ⊙

Ranunculus Aſiaticus grumoſa radice flore ex flavo, & rubro mixto pleno. ⊙

Ranunculus Conſtantinopolitanus gigas flore coccineo. ♃ ⊙

Ranunculus Conſtantinopolitanus gigas flore byſſino. ♃ ☉

Ranunculus montanus Aconiti folio albus flore majore B. pin. albus flore ſimplici Icon. Lob. ☉

Ranunculus montanus Aconiti folio albus flore minore B. pin. Aconitum batrachioïdes Ad. & Icon. Lob. ☉

Ranunculus montanus Aconiti folio flore globoſo B. pin. Ranunculus flore globoſo Dod. ☉

* Ranunculus annuus arvenſis flore minimo luteo. ♃ ☿

* Ranunculus grumoſa radice .ſ. pthora valdenſium Cluſ. ☉

* Ranunculus præcox rutæ folio 1ª Cluſ. ☿

*Ranunculus ferulaceus ſpica falcata nobis perperâm, Melampyrum perpuſillum Lob. ☿

* Raphanus ſylveſtris major flore luteo. ♃ ☉

* Raphanus aquaticus minimus luteus. ♃ ☉

Raphanus niger B. pin. Icon. Lob. ☿

Raphanus minor oblongus B. pin. Radicula ſativa minor Dod. ☿

Raphanus ruſticanus .ſ. vulgaris Matth. ed. B. ruſticanus B. pin. ☉
Raphanus aquaticus foliis in profundas lacinias diviſis B. pin. & prodr. ☉
Raphanus aquaticus alter B. pin. & prodr. Rapiſtrum aquaticum Tab. ☉

* Rapiſtrum monoſpermum B. prodr. ♀
Rapiſtrum arvorum Obſ. & Icon. Lob. flore luteo B. pin. ♀
Rapiſtrum flore albo, ſiliquâ articulatâ B. pin. flore albo Erucæ foliis Icon. Lob. ♀
Rapiſtrum flore albo, ſiliquis in rotundum articulatis, alterum arvorum Park. ♀
Rapiſtrum maximum monoſpermon Cornuti. ♀

* Rapum vulgare album rotundum. ♀
* Rapum vulgare album oblongum. ♀
* Rapum luteum. ♃ ♀

Rapunculus eſculentus B. pin. Rapunculum Dod. ♀
Rapunculus minor foliis inciſis B. pin. Erini .ſ. Rapunculi minimum genus Col. phytob. ♀

Rapunculus

Rapunculus ſpicatus B. pin. alopecuroïdes longiori ſpicâ Cluſ. Idem flore albo. ☉

Rapunculus folio oblongo ſpicâ orbiculari B. pin. Rapuntium corniculatum montanum Col. ☉

Rapunculus ſcabioſæ capitulo cœruleo B. pin. Rapuntium montanum capitatum Col. ♆

Rapunculus perſicifolius magno flore B. pin. Campanula perſicifolia noſtras Obſ. & Icon. Lob. Idem flore albo. ☉

Rapunculus nemoroſus anguſtifolius magno flore, major B. pin. ſylveſtris major hort. Eyſt. ♆

Rapunculus hortenſis latiore folio, ſ. Pyramidalis B. pin. lacteſcens campanula major Lobelii hiſt. Lugd. ☉

* Rapunculus Creticus campanula Pyramidalis altera C. B. P. Petromarula Cretica J. B. ☉

Rapuntium maximum coccineo ſpicato flore Col. in hiſt. Mex. ſ. Flos Cardinalis Barberini. Trachelium Americanum flore rubro Bapt. Ferr. ☉

Rapuntium urens Solonienſe violaceo flore. ♃ ☉

Reſeda

Reſeda vulgaris B. pin. Plinii neotericorum Belgis, Eruca peregrina vel Cantabrica Icon. Lob. ♀

Reſeda latifolia flore flavo. ♃ ♀

Reſeda foliis Calcitrapæ flore albo. ♃ ♀

Reſedæ affinis, Phyteuma B. pin. Phyteuma Monſpelienſium Icon. Lob. ♀

Reſeda alba minor B. pin. Seſamoïdes parvum Salmanticum Cluſ. ☉

Rhamnus ſpinis oblongis flore candicante B. pin. primus Cluſ.

Rhamnus Salicis folio anguſto fructu flaveſcente B. pin. Germanicus Matth. ed. B.

Rhamnus folio ſubrotundo fructu compreſſo B. pin. Paliurus Dod.

Rhamnus catharticus B. pin. hiſt. Lugd. ſolutivus Dod.

Rhamnus catharticus minor B. pin. Spina infectoria pumila prima Cluſ.

Rhaponticum Proſp. Alp. de exot. folio lapathi majoris glabro B. pin. ☉

Rhodia radix Dod. Cluſ. ☉

Rhus folio ulmi B. pin. Rhus Obſoniorum & Coriariorum Cluſ. ☉
Rhus Virginianum B. pin. app. ☉
Rhus myrtifolia Monſpeliaca B pin. myrtifolia Monſpelienſium Ad. & Icon. Lob. ☉
Rhus myrtifolia Belgica B. pin. Chamælæagnus Dod. ☉

Ricinus Dod. vulgaris B. pin. ♆
Ricinus Americanus major ſemine nigro B. pin. Syriacus Cam. deſcr. ♆
* Ricinus minimus Americanus minor C. B. P. ♆

Roſa rubra B. pin. provincialis Dod. deſcr. ☉
Roſa rubra pallidior B. pin. Damaſcena rubra hiſt Lugd deſcr. ☉
Roſa purpurea B. pin. pallida Ad. Lob. ☉
Roſa verſicolor B. pin. prænestina verſicolor Cluſ. deſcr. ☉
Roſa maxima multiplex B. pin. centifolia Batavica Cluſ. ☉
Roſa alba vulgaris major B. pin. alba flore ſimplici hort. Eyſt. ☉

Roſa flore pleno albo hort. Eyſt. ⊙
Roſa moſchata ſimplici flore B. pin. muſcata alba Tab. ⊙
Roſa moſcata flore pleno B. pin. muſcata alba multiplex Tab. ⊙
Roſa ſpinis carens flore majore B. pin. ſine ſpinis Cluſ. ⊙
Roſa ſylveſtris vulgaris flore odorato incarnato B. pin. ſylveſtris Dod. ⊙
Roſa ſylveſtris flore pleno B. pin. ſylveſtris odorata flore multiplici Icon. Lob. ⊙
Roſa ſylveſtris foliis odoratis B. pin. Eglenteria hort. Eyſt. ⊙
Roſa odore Cinamomi flore pleno B. pin. Cinamomea pleno flore Cluſ. ⊙
Roſa lutea ſimplex B. pin. lutea hiſt. Lugd. ⊙
Roſa lutea multiplex B. pin. lutea maxima flore pleno hort. Eyſt. ⊙
Roſa punicea Cornuti deſcr. ⊙
Roſa ſylveſtris pomifera minor B. pin. pinpinellæ folio Ger. ⊙
* Roſa verſicolor .ſ. Striata omnium minima. ♃ ⊙

Roſmarinus hortenſis anguſtiore folio B. pin.

pin Roſmarinum coronarium Dod. ☉
Roſmarinus hortenſis folio aureo. ♃ ☉
* Roſmarinum latifolium Park. ☉

Rubia tinctorum ſativa B. pin. Rubia major Cluſ. ☉
Rubia ſylveſtris hiſt. Lugd. ſylveſtris aſpera, quæ ſylveſtris Dioſcoridis B. pin. ☉
Rubia cynanchica B. pin. Gallium tetraphyllon montanum cruciatum Col. ☉
Rubia anguſtifolia ſpicata B. prodr. ♀
* Rubia ſpicata Cretica Cluſ. ♀

Rubeola arvenſis cœrulea repens B. prod. ♀
Rubeola repens lutea foliis ſpinoſis B. pin. Rubia minima Obſ. & Icon. Lob. ♀
* Rubeola maritima lutea. ♃ ☉
* Rubeola minima alba. ♃ ♀

Rubus Dod. vulgaris .ſ. Rubus fructu nigro B pin. ☉
Rubus vulgaris ſpinis carens. ♃ ☉
Rubus repens fructu cæſio B. pin. reptans Ad. Lob. ☉
Rubus idæus ſpinoſus B. pin. Rubus idæus Dod. Idem fructu albo Cluſ. deſcr. ☉

Rubus idæus lævis B. pin. Rubus idæus Placæ Cluſ. ☉
Rubus odoratus Cornuti. ☉

Ruſcus Matth. ed. B. Ruſcum Dod. ☉

Ruta hortenſis latiſolia B. pin. graveolens hortenſis Dod. ☉
Ruta ſylveſtris major B. pin. ſylveſtris graveolens Dod. ☉
Ruta ſylveſtris minor B. pin. montana legitima Cluſ. app. alt. Peganion Narbonenſium .ſ. Rutula Ad. Lob. ☉
* Ruta Syriaca Adv. Lob. Harmala Dod. ♀

Ruta muraria Dod. B. pin. ☉

S.

SABINA Dod. folio Tamariſci Dioſcoridis B. pin. ⊙

Salix vulgaris alba arboreſcens B. pin. alba perticalis vulgaris hiſt. Lugd.
* Salix latifolia inferne incana J. B.
* Salix latifolia pumila folio rotundo J. B.
* Salix anguſtifolia hirſuta alba J. B.
* Salix anguſtifolia fol. criſpis ſubtûs albicantibus J. B.
* Salix anguſtifolia Amygdalinis fol. J. B.
* Salix anguſtifolia lutea rubrave J. B.
* Salix anguſtifolia humilis repens.

Salvia major Matth ed. B. Dod. major, An ſphacelus Theophraſti B. pin. Eadem foliis ex luteo variegatis, ex purpureo & albo variegatis. ⊙
Salvia minor aurita, & non aurita B. pin. minor Dod. Eadem foliis ex albo vireſcentibus. ⊙
Salvia Cretica non pomifera Cluſ. ⊙

* Salvia ſerrata criſpa J.B. latiſolia ſerrata B. prodr. ⊙

Sambucus Fuchſ. fructu in umbella nigro B. pin. ⊙
Sambucus fructu in umbella viridi B. pin. Sambuci altera ſpecies Dod. deſcr. ⊙
Sambucus racemoſa rubra B. pin. montana racemoſa Ad. & Icon. Lob. ⊙
Sambucus folio laciniato B. pin. lacinioſo folio Dod. ⊙
Sambucus aquatica flore ſimplici B. pin. aquatica hiſt. Lugd. ⊙
Sambucus aquatica flore globoſo pleno B. pin. roſea hort Eyſt. ⊙
Sambucus humilis. ſ. Ebulus B pin. Ebulus Dod. ⊙

Sanicula Dod. officinarum B. pin. ⊙
Sanicula Alpina lutea B. pin. Auricula urſi Dod. Eadem flore pallido & aureo. ⊙
Sanicula Alpina purpurea B. pin. Auricula urſi flore purpureo Icon. Lob. Eadem flore atropurpureo, albo. ⊙
Sanicula Alpina flore variegato B pin. Auricula urſi tertia Cluſ. deſcr. ⊙

* Sani.

* Sanicula montana flore calcari donato C. B. P. pinguicula Cluſ. hiſt.

Sanicula Alpina foliis Borraginis villoſa B. pin. Auricula urſi Myconi hiſt. Lugd. ☉

Sanicula montana latifolia laciniata B. pin. Cortuſa Matthioli hiſt. Lugd. ☉

Sanicula montana Americana .ſ. Cortuſa Americana ſtapelii in Theoph. ♃ ☉

Sanicula montana rotundifolia major B. pin. montana ſecunda Cluſ. guttata Cam. ☉

Sanicula montana crenata umbilico pallido. ♃ ☉

Sanicula montana crenata umbilico rubro. Sedum montanum ſerratum guttato flore Park. ☉

Saponaria Dod. major lævis B. pin. ☉

Saponaria flore pleno Cornuti deſcr. ☉

Saponaria concava Anglica B. prodr. ☉

Satureia Dod. domeſtica hort. Eyſt. hortenſis .ſ. Cunila ſativa Plinii B. pin. ☿

Satureia montana B. pin. Thymbra Dod. perenni radice. ☉

Saxifraga rotundifolia alba B. pin. alba Chelidonīdes Obſ. & Icon. Lob. ☉
Saxifraga aurea Dod. rotundifolia aurea B. pin. ☉
* Saxifraga aurea longifolia humilior. ♃ ☉

Scabioſa vulgaris major Dod. pratenſis hirſuta quæ officinarum B. pin. Eadem flore albo. ☉
Scabioſa montana calidarum regionum Obſ. & Icon. Lob. fruticans anguſtifolia alba B. pin. ☉
Scabioſa prolifero flore Cluſ. prolifera B. prodr prolifera folio latiore ejuſdem pin. ♀
Scabioſa Alpina centauroïdes hort Eyſt. Alpina foliis Centaurii majoris B. pin. ☉
Scabioſa latifolia rubra non laciniata ſecunda B. pin. rubra Auſtriaca Icon Lob. ☉
Scabioſa argentea anguſtifolia B. prodr. ☉
Scabioſa peregrina rubra capitulo oblongo B. pin. Indica flore ſaturatè rubro hort. Eyſt. Eadem quæ
Scabioſa Indica flore dilutè rubro hort Eyſt. ♀

Sca-

Scabioſa capitulo globoſo major B. pin. minor .ſ. Columbaria Obſ. & Icon. Lob. ☉

Scabioſa capitulo globoſo minor B. pin. anguſtifolia flore ſubcœruleo hort. Eyſt. Eadem flore albo, & roſeo. ☉

Scabioſa ſtellata folio laciniato major B. pin. major Hiſpanica, .ſ. prima Cluſ. ♀

Scabioſa ſtellata folio non diſſecto B. pin. arborea Proſp. Alp. de exot. ☉

* Scabioſa gramineis foliis glabris. ♃ ☉

* Scabioſa diſſectis fol. flore amplo ſemine lanuginoſo. ♃ ☉

* Scabioſa 2 Hiſpanica Cluſ. ♀

* Scabioſa minima annua ſemine anguloſo. ♃ ♀

Scammonia Syriaca B. pin. Scammonium Dod. ☉

Scammonia Monſpeliaca foliis rotundioribus B. pin. Scammonium Monſpelienſe Ad. & Icon. Lob. ☉

Scandix ſemine roſtrato vulgaris B. pin. Scandix, Pecten Veneris Dod. ♀

Scilla Dod. vulgaris radice rubra B. pin. ☉

Scolymus Theophraſti, Eryngium luteum Monſpelienſium Obſ. & Icon. Lob. Scolymus chryſanthemos B. pin. ☉
Scolymus chryſanthemos annuus. ♃ ♀

Scordium Dod. B. pin. ☉
Scordium alterum .ſ. ſalvia agreſtis B. pin. Salvia ſylveſtris Bock. ☉
Scordium ſemine talari, flore cœruleo. Idem flore roſeo. ♃ ☉

Scorpioïdes Bupleuri folio B. pin. repens Bupleuri folio Obſ. & Icon. Lob. ♀
Scorpioïdes ſiliquâ craſsâ Boëlii Ger. app. ♀
Scorpioïdes ſiliquâ ſtriatâ minus hirſutâ. ♃ ♀
Scorpioïdes Matthioli Dod. portulacæ folio. B. pin. ♀

Scorzonera latifolia ſinuata B. pin. major Hiſpanica prima Cluſ. ☉
Scorzonera foliis nervoſis B. pin. Germanica anguſtifolia Tab. ☉

Scor-

Scorzonera pulverifera. ♃ ☉

Scorzonera latifolia humilis nervosa B. pin. humilis latifolia pannonica secunda Clus. ☉

Scorzonera angustifolia prima B. pin. humilis angustifolia pannonica tertia Clus. ☉

* Scorzonera angustifolia subcœrulea C. B.P angustifolia elatior pannonica quarta Clus. ☉

* Scorzonera sem. lanuginoso nobis, Tragopogon hirsutum Apulum humile Col. ☉

Scrophularia Dod. nodosa fœtida B.pin. ☉

Scrophularia aquatica major B. pin. Betonica aquatilis Dod. ☉

Scrophularia folio Urticæ B.pin. peregrina Cam. ♀

Scrophularia flore luteo B. pin. Lamium pannonicum secundum exoticum Clus. ♀

Scrophularia foliis & caule hirsutis. ♃ ☉

Scrophularia ruta canina dicta vulgaris B. pin. Ruta canina Clus. ☉

* Scrophularia Sambuci fol. Park. ☉

* Scutellaria Americana Malvæ folio. ♃ ☉

Secale Fuchſ. hybernum vel majus B. pin.

Securidaca lutea major B. pin. vera Cluſ. ♀
Securidaca dumetorum major flore vario ſiliquis articulatis B. pin. Hedyſarum purpureum Tab. ☉
Securidaca minor lutea ſiliquis articulatis. ♃ ♀

Sedum majus arboreſcens floſculis candidis B. pin. Sempervivum .ſ. Sedum arboreſcens majus Dod. ☉
Sedum majus Fuchſ. majus vulgare B. pin. ☉
Sedum minus teretifolium album B. pin. Sempervivum minus alterum Dod. ☉
* Sedum longifolium Citrino flore. ♃ ☉
* Sedum arvenſe flore rubente B. prodr. ♀
* Sedum teretifolium majus album. ♃ ☉
Sedum minus teretifolium album alterum Sempervivum minus æſtivum Obſ. & Icon. Lob. ♀

Sedum

Sedum minus folio circinato B. pin. Aizoon daſyphyllum Dal. hiſt. Lugd. ☉

Sedum minus luteum folio acuto B. pin. Aizoon hæmatoïdes Icon. Lob. Idem flore albo. ☉

Sedum minus luteum ramulis reflexis B. pin. Aizoon ſcorpioïdes Icon. Lob. ☉

Sedum minus cauſticum hort Eyſt. Sempervivum minus vermiculatum acre B. pin. ☉

Sedum minus .ſ. Vermicularis inſipida hort. Eyſt. Sempervivum minus vermiculatum inſipidum B. pin. ☉

Sedum minus fruticoſum B. pin. minimum arboreſcens vermiculatum Obſ. & Icon. Lob. ☉

Sedum Alpinum hirſutum luteum B. pin. Phyllum thelygonon hiſt. Lugd. ☉

Sedum Alpinum foliolis crenatis aſperis B. prodr. ☉

Sedum Pyrenæum ſerratum longifolium. ♃ ☉

Sedum Pyrenæum ſerratum flore albo punctis purpureis conſperſo. ♃ ☉

Sedum montanum tomentoſum B. pin. Sempervivum rubrum montanum γναφαλῶδες Col. ☉

Sedum

Sedum tridactylites Alpinum majus album B. pin. Saxifraga alba petræa Pon. descr. mont. Bald. ☉

Sedum tridactylites tectorum B. pin. Paronychia altera rutaceo folio Obs. & Icon. Lob. ♀

Selinum Sii foliis Ger. Park. ♀

Senecio minor vulgaris B. pin. Erigerum minus Dod. ♀

Senecio incanus pinguis B. pin. Erigerum majus Dod. ♀

* Senecio Jacobææ fol. nobis, fortê folio non laciniato J. B. ♀

* Senecio Canadensis acris alba. ♃ ♀

Serpyllum vulgare majus B. pin. album primum vel secundum Tab. ☉

Serpyllum foliis citri odore B. pin. citratum hort. Eyst. ☉

Serpyllum vulgare minus B. pin. vulgare Dod. Idem flore albo, quod

Serpillum minus flore albo Tab. ☉

Serpyllum latifolium hirsutum B. prodr. ☉

Serpyllum angustifolium hirsutum B. pin. pannonicum tertium Clus. ☉ Ser-

Serratula Matth. ed. B. Dod. ☉
Serratula foliis non inciſis. Tinctoria prima Tab. ☉

Seſeli montanum Cicutæ folio glabrum B. pin. primum montanum pannonicum Cluſ. ☉
Seſeli Creticum Dod. Creticum minus B. pin. Vide tordylium. ♀
Seſeli Æthiopicum frutex Dod. Æthiopicum Salicis folio B. pin. ☉
Seſeli Maſſilienſe fœniculi folio B pin. Maſſilienſe fœniculi folio craſſiore Icon. Lob. ☉
Seſeli foliis tenuioribus. Fœniculus tortuoſus Dal. hiſt. Lugd. ☉
Seſeli paluſtre lacteſcens B. prodr. & pin. ☉
Seſeli pratenſe Lobelii hiſt Lugd. pratenſe, Silaus forte Plinii B. pin. ☉

Sideritis foliis hirſutis profundè crenatis B. pin. Monſpeliaca ſcordioïdes floribus luteis Ad. & Icon Lob. ☉
Sideritis vulgaris prima Cluſ. ☉
Sideritis Alpina hyſſopifolia B. pin. montana Icon. Lob. ☉

Sideritis

Sideritis Alſines trixaginis foilo B. pin. & prodr. ♀
Sideritis arvenſis latifolia glabra B. pin. Alyſſum majus Tab. ♀
Sideritis arvenſis anguſtifolia rubra B. pin. Ladanum ſegetum hiſt. Lugd. Eadem flore albo. ♀
Sideritis glutinoſa odorata. ♃ ☉
* Sideritis ſalviæ folio noſtras donato Park. ☉
* Sideritis verticillis ſpinoſis J. B. ♀

Siliqua edulis B. pin. Ceratonia Dod. Carobe vulgò.
Siliqua ſylveſtris rotundifolia B. pin. Arbor Judæ Dod.

Sinapi Rapi folio B. pin. ſativum Erucæ aut Rapi folio Obſ. & Icon. Lob. ♀
Sinapi Erucæ folio B. pin. ſylveſtre Dod. ♀
Sinapi ſiliquâ hirſutâ B. pin. ſativum alterum Icon. Lob. app. ♀
* Sinapi ſiliquâ hirſutâ, ſemine rubro. ♃ ♀

Siſarum Dod. Germanorum B. pin. ☉

Siſon

Siſon .ſ. Petroſelinum Macedonicum Dodonæi hiſt. Lugd. Siſon quod Amomum officinis noſtris B. pin. ♀

Sium Dod. Sion .ſ. Apium paluſtre foliis oblongis B. pin. ⊙
Sium alterum Oluſatri facie Obſ. & Icon. Lob. Sium erucæ folio B. pin. ⊙
Sium repens umbellis ad geniculos diſpoſitis. ♃ ⊙
* Sium minimum ferulaceis foliis. ♃ ⊙

Smilax aſpera rutilo fructu Cluſ. aſpera fructu rubente B. pin. ⊙
Smilax aſpera nigro fructu Cluſ. aſpera minûs ſpinoſa fructu nigro B. pin. ⊙

Smyrnium Fuchſ. Hippoſelinum Theophraſti, vel Smyrnium Dioſcoridis B. pin. ⊙
Smyrnium Creticum Matth. ed. B. Amani montis Dod. peregrinum rotundo folio B. pin. ♀

Solanum officinarum B. pin. hortenſe Fuchſ. Dod. Idem baecis nigris. ♀
Solanum veſicarium Dod. B. pin. Idem foliis variegatis. ⊙

Solanum veſicarium Indicum B. pin. halicacabum, ſive Solanum Indicum Cam. hort. Eyſt. ♀

Solanum Virginianum veſicarium fructu luteo. Virginianum .ſ. halicacabum, .ſ. Alkekengi Virginenſe Park. ♀

Solanum lethale, Belladonna Cluſ. Solanum μανικὸν B. pin. ☉

Solanum ſomniferum Cluſ. ſomniferum verticillatum B. pin. ☉

Solanum fruticoſum bacciferum B. pin. Amomum Plinii Obſ. Lob. ☉

Solanum quadrifolium bacciferum B. pin. Herba Paris Dod. ☉

Solanum ſcandens .ſ. Dulcamara B. pin. Dulcamara Dod. Idem flore albo quod Dulcis amara flore albo hort. Eyſt. ☉

Solanum ſcandens foliis ex albo variegatis. ♃ ☉

* Solanum Virginianum flore cœruleo. ♃ ☉

* Solanum Virginianum flore albo. ♃ ☉

* Solanum marinum humile flore cœruleo. ♃ ☉

* Solanum marinum humile flore albo. ♃ ☉

Solanum

Solanum tuberoſum eſculentum Matth ed. B. Arachidna Theophraſti forte, Pappas Perüanorum Cluſ. ☉

Solanum racemoſum Indicum. Magnum Virginianum rubrum Park. ☉

Solanum foliis & caule ſpinoſis. ♃ ☉

Solanum pomiferum fructu rotundo ſtriato molli B. pin. Pomum amoris hiſt. Lugd. ♀

*Solanum Pomum amoris fructu ſtriato molli aureo. ♃ ♀

Solanum pomiferum fructu rotundo parvo molli. ♃ ♀

Solanum pomiferum fructu rotundo ſtriato duro B. pin. Mala Æthiopica Dod. ♀

Solanum pomiferum fructu oblongo B. pin. Mala inſana Dod. ♀

Solanum pomo ſpinoſo rotundo longo flore B. pin. Strammonia Fuchſ. Dod. ♀

Solanum fœtidum pomo ſpinoſo oblongo B. pin. Datura Turcarum hort. Eyſt. Idem flore violaceo. ♀

Solanum fœtidum pomo grandibus aculeis donato. ♃ ♀

Solanum fœtidum pomo ſpinoſo rotundo ſemine pallido B. pin. Datura Ægyptia

Veſl. Obſ. in proſp. Alp. Idem triplici calice. ♀

Solaniſolia Circæi dicta major B. pin. Circæa lutetiana Obſ. & Icon. Lob. ☉

Solanum Mexiocanum flore magno B. pin. Admirabilis Perüana rubro flore Cluſ. Idem flore luteo, luteo ex rubro mixto, albo rubris lineis diſtincto. ♀

Soldanella Dod. maritima minor B. pin. ☉

Soldanella Alpina minor Cluſ. Alpina rotundifolia B. pin. ☉

Sonchus aſper non laciniatus B. pin. aſperior Lobelii hiſt. Lugd. ♀

Sonchus aſper laciniatus & non laciniatus B. pin. aſper Dod. ♀

Sonchus lævis laciniatus latiſolius B pin. lævis Dod. ♀

Sonchus lævis laciniatus muralis parvis floribus B. pin. alter folio, profundis laciniis ſinuato hederaceo Obſ. & Icon. Lob. ☉

Sonchus anguſtiſolius maritimus B. prodr. ♀

Sonchus lanatus Dal. hiſt. Lugd. villoſus luteus major B. pin. ♀

* Sonchus Auſtriacus lævis tertius cœruleo flore Cluſ. hiſt. ☉

Soncho affinis, Lampſana domeſtica B. pin. Lampſana Dod. ☿

Sorbus Dod. ſativa B pin.

Sorbus ſylveſtris foliis domeſticæ ſimilis B. pin. Ornus Dod.

* Sparganium ramoſum, & non ramoſum C. B. P. platanaria Dod. ☉

Spartium arboreſcens ſeminibus lenti ſimilibus B. pin. Geniſta Italica Dod. ☉

* Spartum 2 Cluſ. hiſt. Spartum Monoſpermon Hiſpanicum flore luteo Park. ☉

Sphondilium Dod. vulgare hirſutum B. pin. ☿

* Sphondylium hirſutum fol. anguſtioribus B. pin. ☿

* Spiræa Theophr. fortè Cluſ. frutex ſpicatus foliis Salignis ſerratis C. B. Pin.

Stachys Fuchſ. Dod. major Germanica B. pin. ☉

Stachys Cretica B. pin. Pſeudoſtachys Cretica ejuſdem prodr. ⊙

* Stachys Cretica anguſtifolia nobis, Pſeudoſtachys B. prodr.

* Stachys fol. obſcure virenti flore ferrugineo. ♃ ⊙

Stachys paluſtris fœtida B. pin. Clymenum minus Dal. hiſt. Lugd. ⊙

Staphiſagria Fuchſ. Dod. ☿

Stœbe major foliis cichoraceis mollibus lanuginoſis B. pin. Salmantica prima Cluſ. Eadem flore albo. ⊙

Stœbe caliculis argenteis B. pin. Salmantica tertia Cluſ. argentea Salamantica minor Obſ. & Icon Lob. Eadem flore albo. ⊙

Stœbe incana Cyano ſimilis tenuifolia B. pin. Auſtriaca humilis Cluſ. deſcr. ⊙

* Stœbe humilis Cluſ. incana tenuifolia cyano ſimilis C. B. P. ⊙

Stœchas Arabica Cord. hiſt. brevioribus ligulis Cluſ. purpurea B. pin. ⊙

* Stœchas folio ſerrato Ger. Dod. folio criſpo Cluſ. hiſt. ⊙

Styrax

Styrax hiſt. Lugd. folio mali cotoneæ B. pin.

Suber latiſolium perpetuo virens B. pin. Phellos .ſ. ſuber Dod.

Succiſa Fuchſ. glabra B. pin. Eadem flore albo, roſeo. ⊙

Symphytum, Conſolida major B. pin. majus .ſ. Conſolida major Matth. ed. B. Idem flore purpureo, pallido. ⊙
Symphytum tuberoſum majus Cluſ. majus tuberoſa radice B. pin. ⊙
Symphytum maculoſum .ſ. Pulmonaria latifolia B. pin. Pulmonaria vulgaris maculoſo folio Cluſ. Idem flore albo. ⊙
Symphytum .ſ. Pulmonaria anguſtifolia rubente cœruleo flore B. pin. Pulmonaria foliis Echii Icon. Lob. ⊙

Syringa cœrulea B. pin. flore cœruleo hort. Eyſt. Lillach Dod. Eadem flore albo quæ Syringa flore lacteo hort. Eyſt.
Syringa Italica flore ſimplici albo hort: Eyſt. alba .ſ. Philadelphus Athenæi B. pin.

T.

TAMARIX Narbonenſis hiſt. Lugd. altera folio tenuiore .ſ. Gallica B. pin.

Tanacetum Dod. vulgare luteum B. pin. ☉
Tanacetum criſpum Dod. foliis criſpis B. pin. ☉
Tanacetum minus album odore Camphoræ B. pin. minus Dod. Achillea .ſ. millefolium nobile Tab. ☉
Tanacetum montanum purpureum. Vide Millefolium. ☉
Tanacetum montanum inodorum minore flore B. pin. inodorum Dod. ☉
Tanacetum inodorum flore majore B. pin. inodorum primum Cluſ. Vide bellis. ☉
Tanacetum .ſ. flos Africanus major flore pleno B. pin. Othonna major polyanthos Obſ. & Icon. Lob. ♀
Tanacetum Africanum majus ſimplici flore B. pin. Flos Africanus major ſimplici flore luteo Icon. Lob. ♀
Tanacetum Africanum .ſ. flos Africanus minor

nor B. pin. Caryophyllus Indicus flore luteo rubeſcente ſimplici minor hort. Eyſt. Idem flore pleno qui Caryophyllus Indicus plenus flore luteo rubeſcente minor hort. Eyſt. ♀

Tanacetum .ſ. flos Mexicanus flore fiſtuloſo ſimplici B. pin. Caryophyllus Mexicanus fiſtuloſo flore, ſimplex Col. ♀

Tanacetum .ſ. flos Mexicanus flore fiſtuloſo pleno B. pin. Caryophyllus Mexicanus alter polyanthos fiſtuloſo flore Col. ♀

Taxus Dod.

Telephium vulgare B. pin. album Fuchſ. ☉

Telephium purpureum majus B. pin. purpuraſcens Fuchſ. ☉

Telephium purpureum minus B. prodr. ☉

Telephium latifolium peregrinum B. pin. primum Hiſpanicum Cluſ. ☉

Telephium repens folio deciduo B. pin. Sextum, Cepæa Pancii Cluſ. ☉

* Telephium Portulacæ folio. ♃ ☉

* Telephium peregrinum rubens. ♃ ☉

Telephium repens folio non deciduo B. pin. legitimum Imperati ſeptimum Cluſ. ☉

Tere-

Terebinthus Dod. vulgaris B. pin.

Teucrium B. pin. vulgare fruticans .ſ. primum Cluſ. Chamædrys aſſurgens Dod. ⊙

* Teucrium fruticans flore purpureo. ♃ ⊙

* Teucrium Bæticum fruticans Cluſ. hiſt. Bæticum Ger. Tab. ⊙

Thalictrum majus ſiliquâ angulosâ, aut ſtriatâ B. pin. Ruta pratenſis herbariorum Obſ. & Icon. Lob. ⊙

Thalictrum pratenſe maximum Monſpelienſium aſperioribus foliis. ♃ ⊙

Thalictrum majus florum ſtaminibus purpuraſcentibus B. pin. quartum Dod. deſcr. ⊙

Thalictrum Canadenſe Cornuti. ⊙

Thalictrum minus Dod. B. pin. Hypecoum hiſt. Lugd. ⊙

Thalictrum minimum fœtidiſſimum B. prodr. ⊙

Thapſia latifolia villoſa B. pin. prima Cluſ. ⊙

Thapſia Carotæ folio B. pin. vulgaris Carotæ effigie Obſ. & Icon. Lob. ☉

Thlaſpi arvenſe ſiliquis latis B. pin. latius Dod. ♀

Thlaſpi vulgatum ſerrato folio Matth. ed. B. ♀

Thlaſpi arvenſe perfoliatum majus B. pin. Oleraceum Tab. ♀

Thlaſpi umbellatum arvenſe Iberidis folio B. pin. amarum Tab. ♀

* Thlaſpi Iberidis folio tenuiſſimê diviſo. ♃ ♀

Thlaſpi umbellatum Creticum Iberidis folio B. pin. Candiæ umbellatum Iberidis folio Obſ. & Icon. Lob. ♀

Idem flore albo, quod Cappadocicum flore albo hort. Eyſt. ♀

Thlaſpi Alpinum Bellidis cœruleæ folio B. pin. montanum Burſæ paſtoris fructu Col. ☉

Thlaſpi montanum ſempervirens B. pin. montanum candidum hiſt. Lugd. ☉

Thlaſpi parvum ſaxatile flore rubente B. pin. Lithothlaſpi quartum carnoſo rotundo folio Col. ☉

Thlaſpi montanum incanum luteum B. prodr. Supinum luteum Icon. Lob. ☉
Thlaſpi, Alyſſum dictum campeſtre minus B. pin. Alyſſum minus Dal. hiſt. Lugd. ♀
* Thlaſpi Alyſſum minimum. ♃ ♀
Thlaſpi maritimum Dal. hiſt. Lugd. Thlaſpi Alyſſon dictum maritimum B. pin. ♀
Thlaſpi clypeatum Cluſ. biſcutatum aſperum hieracifolium & majus B. pin. ☉
Thlaſpi biſcutatum annuum. Leucoïum Alyſſoïdes umbellatum montanum Col. ♀
Thlaſpi clypeatum Serpylli folio B. pin. Ionthlaſpi minimum ſpicatum lunatum Col. ♀
Thlaſpi fruticoſum ſpinoſum B. pin. fruticoſum ſpinoſum Narbonenſe Obſ. & Icon. Lob. ☉
Thlaſpi incanum Machlinienſe Cluſ. fruticoſum incanum B. pin. ☉

Thuya Theophraſti B. pin. Arbor vitæ Dod.

Thymelæa Dod. foliis lini B. pin. ☉
Thymelæa foliis candicantibus ſerici inſtar mollibus B. pin. Tartonraire Galloprovinciæ

vinciæ Maſſilienſium Ad. & Icon. Lob. ☉

Thymus Fuchſ. vulgaris folio tenuiore B. pin. ☉
Thymus vulgaris latiore folio B. pin. Thymum durius Dod. ☉
Thymum cephaloton Dod. Thymus capitatus qui Dioſcoridis B. pin. ☉
Thymum Creticum fruteſcens. Marum forte Matth. ed. B. hiſt. Lugd. ☉

Tilia Dod. fœmina folio majore B. pin.

Tithymalus characias amygdaloïdes Obſ. & Icon. Lob. B. pin. ☉
Tithymalus characias rubens peregrinus B. pin. characias Monſpelienſis Cluſii & Matthioli Icon. Lob. ☉
Tithymalus characias folio lanuginoſo B. pin. characias ſecundus Cluſ. deſcr. ☉
Tithymalus characias anguſtifolius B. pin. myrſinites tertius anguſtifolius Tab. ☉
Tithymalus characias radice repente. ♃ ☉
Tithymalus myrtifolius arboreus B. pin. arboreus Proſp. Alp. de exot. dendroïdes Matth. ed. B. ☉

Ti-

Tithymalus myrſinites Dod. myrſinites latifolius B. pin. ☉

Tithymalus myrſinites fructu verrucæ ſimili B. pin. verrucoſus Dalechampii hiſt. Lugd. ☉

Tithymalus maritimus B. pin. paralius ex locis maritimis Dod. ☉

Tithymalus maritimus .ſ. Eſula rara Venetorum flore albo. ☉

Tithymalus latifolius Hiſpanicus B. pin. platyphyllos Cluſ. ☉

Tithymalus helioſcopius Dod. B. pin. ☿

Tithymalus cypariſſias Fuchſ. Dod. B. pin. Idem capitulo rubente. ☉

Tithymalus leptophyllos Matth. ed. B. hiſt. Lugd. Tithymalus .ſ. Eſula exigua B. pin. ♆

Tithymalus foliis Pini forte Dioſcoridis Pityuſa B. pin. Eſula minor Dod. ☉

Tithymalus paluſtris fruticoſus B. pin. Eſula major Dod. ☉

Tithymalus montanus non acris B. pin. Eſula dulcis Tragi hiſt. Lugd. ☉

Tithymalus foliis brevibus aculeatis B. pin. Pityuſa Matth. ed. B. ♆

Tithymalus Epithymi fructu Col. Peplios altera ſpecies B. pin. ☉

* Tithymalus linariæ fol. flore lunato Col. ☉
* Tithymalus ſalicis fol. ♃ ☉
* Tithymalus caracias pyriformi radice. ♃ ☉
* Tithymalus Eſula exigua fol. obtuſis B. in prodr. ☿

* Tordylion majus vide Caucalis major Cluſ. ☿
* Tordylion minimum Apulum Col. vide Seſeli Creticum minus. ☿
* Tordylion Syriacum ſem. Granulato nobis, Gyngidium 1. Dod. Gyngidium latifolium Syriacum Park. Caucalis Syriaca cum maximo ſem. J. B. ☿

Tormentilla Dod. ſylveſtris B. pin. ☉
Tormentilla Alpina folio ſericeo B. pin. candida Dal. hiſt. Lugd. Heptaphyllon Cluſ. ☉

Trachelium umbelliferum cœruleum Pon. Cervicaria Valerianoïdes cœrulea B. pin. ☉

Tragacantha

Tragacantha Matth. ed. B. hiſt. Lugd. B. pin. ☉

Tragopogon pratenſe luteum majus B. pin. luteum Obſ. & Icon. Lob. ♀
* Tragopogon pratenſe minus Park. deſcr. ♀
* Tragopogon laciniatum luteum C. B. P. ♀
* Tragopogon Coronopi folio minus. B. pin. ♀
Tragopogon Coronopi folio B. pin. ♀
Tragopogon gramineo folio radice villoſa B. pin. Barbula hirci minor Tab. ♀
Tragopogon purpureocœruleum Porri folio B. pin. purpureum hort. Eyſt. Idem flore dilutiore. ♀
Tragopogon gramineis foliis hirſutis B. pin. alter gramineo folio ſuaverubente flore Col. ♀

Tragoriganum Proſp. Alp. exot. ☉

Tribulus terreſtris Dod. terreſtris Ciceris folio fructu aculeato B. pin. ♀

Trichomanes Fuchſ. Dod. Trichomanes ſ. polytrichum officinarum B. pin. ☉
* Trichomanes ramoſum C. B. P. ☉
Trichomanes minus & tenerius B. pin. minor mas Tab. candidum Thal. ☉

Trifolium bituminoſum Dod. bitumen redolens B. pin. ☉
Trifolium pratenſe purpureum B. pin. pratenſe Dod. ☉
Trifolium pratenſe album Fuchſ. B. pin. ☉
Trifolium pratenſe luteum capitulo breviore B. pin. pratenſe luteum Fuchſ. ☉
Trifolium pratenſe luteum capitulo Lupuli vel agrarium B. pin. luteum minimum Penæ & Lobelii hiſt. Lugd. ♀
Trifolium pratenſe majus caule fiſtuloſo foliis ſubtùs punctatim nigris. ♃ ♀
Trifolium montanum purpureum majus B. pin. majus ſecundum Cluſ. deſcr. ☉
Trifolium montanum purpureum folio acuto crenato B. prodr. ♀
Trifolium montanum album B. pin. majus primum Cluſ. ☉
Trifolium Alpinum flore magno radice ſubamarâ. Anguſtifolium Alpinum Pon.

deſcr. mont. Bald. ☉

Trifolium ſpicâ ſubrotundâ rubrâ B. pin. Lagopus major folio Trifolii Dod. ♀

Trifolium ſpicâ oblongâ rubrâ B. pin. majus tertium Cluſ. ☉

Trifolium montanum ſpicâ longiſſimâ rubente B. pin. Lagopus altera folio pinnato Obſ. & Icon. Lob. ☉

Trifolium lagopoïdes flore pallidè luteo. ♃ ☉

Trifolium montanum anguſtiſſimum ſpicatum B. pin. Lagopus altera anguſtifolia Ad. & Icon. Lob. ♀

Trifolium arvenſe humile ſpicatum, .ſ. Lagopus B. pin. Lagopus .ſ. Pes leporis Matth. ed. B. Obſ. & Icon Lob. ♀

Trifolium ſtellatum B. phytop. ♀

Trifolium Salmanticenſi affine. ♃ ♀

Trifolium fragiferum Cluſ. cur. poſt. Idem flore albo. ☉

Trifolium album anguſtifolium floribus velut in capitulum congeſtis B. pin. Dorycnium Monſpelienſium Icon. Lob. ☉

Trifolium nodiflorum. ♃ ♀

Trifolium cochleatum fructu latiore B. pin. cochleatum primum Dod. ♀

Trifolium

Trifolium cochleatum turbinatum .ſ. fructu compreſſo oblongo B. pin. Medicæ ſpecies ſecunda Ad. Lob. ♀

Trifolium cochleatum fructu globoſo plano. ♃ ♀

* Trifolium cochleatum ſpinoſum majus ſpinis ſurſum, & deorſum tendentibus. ♃ ♀

* Trifolium cochleatum ſpinoſum minus nigrum ciliare. ♃ ☉

* Trifolium cochleatum ſpinoſum minus, polycarpon minus hiſpidum. ♃ ♀

* Trifolium cochleatum ſpinoſum minus coronatum Cherleri J. B. ♀

Trifolium cochleatum maritimum tomentoſum B. pin. Medica marina Cluſ. ☉

Trifolium cochleatum fructu nigro hiſpido B. pin. cochleatum alterum Dod. ♀

Trifolium cochleatum folio cordato maculato B. pin. Medica Arabica Cam. ♀

Trifolium cochleatum echinatum minus B. prodr. ♀

Trifolium cochleatum glomerulis Perſonatæ Cherleri J. B. ♀

Trifolium echinatum arvenſe B. pin. Tribulus terreſtris minor repens hiſt. Lugd. ♀

Trifolium ſubterraneum Tricoccon. ♃ ♀
Trifolium ſiliquâ cornutâ .ſ. Medica B. pin. Medica Dod. ☉
Trifolium ſylveſtre luteum ſiliquâ cornutâ, vel Medica fruteſcens B. pin. Medica flavo flore Cluſ. flore albo, cinereo, violaceo nigricante, herbaceo, purpureo. ☉
* Trifolium veſicarium rubello flore. ♃ ♀
* Trifolium capitulo glomerato rigido J. B. ♀
* Trifolium corniculatum luſitanicum flore rubro Park. in app. ad Theatr. ♀
Trifolium acetoſum vulgare B. pin. acetoſum Dod. Idem flore purpuraſcente. ☉
Trifolium acetoſum corniculatum B. pin. Oxys flavo flore Cluſ. ♀
* Trifolium acetoſum corniculatum Indicum erectum. ♃ ♀
Trifolium hepaticum flore ſimplici B. pin. Hepatica trifolia cœruleo flore Cluſ. rubro flore ejuſdem flore albo hort. Eyſt. ☉
Tritolium hepaticum flore pleno B. pin. Hepatica trifolia cœruleo pleno flore Cluſ. ☉

Tripolium Dod. majus cœruleum B. pin. ☉

Triticum Hyberum ariſtis carens B. pin. Triticum Dod. ☿
Triticum ſpicâ multiplici B. pin. Icon. Lob. ☿
Triticum ariſtis longioribus ſpicâ albâ B. pin. Typhinum quartum Tab. ☿

Tulipa Dod. Tulipa Turcarum Geſn. app. ad Cord. hiſt. ☉
Tulipa ſerotina πολυκλαδὴς major Cluſ. ſerotina ramoſa major B. pin. ☉
Tulipa ſerotina flore pleno B. pin. ☉
Tulipa dubia pumilio Cluſ. pumilio latiſolia rubra B. pin. ☉
Tulipa Apenninea Cluſ. minor lutea Italica B. pin, vulgò Bononienſis. ☉
Tulipa Perſica hort. Eyſt. variegata Perſica B. pin. ☉

Tuſſilago Fuchſ. vulgaris B. pin. ☉

V.

VALERIANA hortenſis Dod. hortenſis, Phu folio Oluſatri Dioſcoridis B. pin. ☉

Valeriana foliis calcitrapæ B. pin. annua .ſ. æſtiva Cluſ. ♀

Valeriana ſylveſtris major B. pin. ſylveſtris Dod. ☉

Valeriana Græca Dod, cœrulea B. pin. Eadem flore albo quæ

Valeriana Græca flore albo hort Eyſt. ☉

Valeriana paluſtris minor B. pin. Phu minimum Matth. ed. B. ☉

* Valeriana Urticæ folia flore albo Corn.

* Valeriana Alpina ſylveſtris prima latifolia Cluſ. ☉

Valeriana rubra Dod. B. pin. Eadem flore albo. ☉

Valeriana rubra anguſtifolia B. prodr. ☉

Valeriana campeſtris inodora major B. pin. album olus Dod. ♀

Valerianella altera nudo umbilicato,& brevi ſemine Col. ♀

Valerianella cornucopioïdes echinata Col. echinata B. pin. ♀

Valerianella altera tenuifolia ſemine ſcabioſæ ſtellato Col. ſemine ſtellato B. pin. ♀

* Valerianella noſtras major ſcabioſæ ſemine. ♂ ♀

* Valerianella ſemine hirſuto minore rotundo. ♂ ♀

* Valerianella Indica Cluſ ♀

* Valerianella vulgaris major ſerotina. ♂ ♀

* Valerianella umbilicato nudo ſemine oblongo. ♂ ♀

Valerianella ſemine, acinis uvarum ſimili .ſ. hirſuto, extremo umbilicato. ♂ ♀

Verbaſcum mas latifolium luteum B. pin. mas & candela regia Obſ. Lob. ♀

Verbaſcum fœmina flore luteo magno B. pin. album fœmina luteo flore Dod. Gall. ed. ♀

Verbaſcum ramoſum floribus luteis parvis. Candidum fœmina Fuchſ. ♀

Verbaſcum ramoſum floribus parvis albis. ♂ ♀

Verbaſcum nigrum Dod. nigrum flore ex luteo purpuraſcente B. pin. ♀

Verbaſcum perenne nigrum floribus albis. ♃ ⊙

*Verbaſcum nigrum Papaveris corniculati folio C. B. P. laciniatum Lugd. ♀

Verbaſcum ſalviſolium fruticoſum luteo flore Ad. & Obſ. Lob. latis Salviæ foliis B. pin. ⊙

Verbaſcum ſalviſolium fruticoſum flore purpureo. ♃ ⊙

Verbaſcum anguſtis Salviæ foliis B. pin. Phlomis Lychnitis Cluſ. ⊙

Verbaſculum pratenſe odoratum B. pin. Primula veris flavo flore elatior Cluſ. ⊙

Verbaſculum pratenſe, vel ſylvaticum inodorum B. pin. Primula veris pratenſis inodora luteo-pallida Obſ. & Icon. Lob. ⊙

Verbaſculum ſylvarum majus ſingulari flore B. pin. Primula veris pallido flore humilis Cluſ. ⊙

Verbaſculum Conſtantinopolitanum. Carchichæc Turcarum Cornuti. Idem flore albo, ſulphureo, dilutè violaceo ⊙

Verbaſculum ſylveſtre magno, plenoque flore

flore B. pin. Primula veris flore pleno hort. Eyſt. ⊙

Verbaſculum proliferum B. pin. Primula veris gemino flore hort. Eyſt. ⊙

* Verbaſculum majus umbellatum purpureum. ⊙

* Verbaſculum majus umbellatum album. ⊙

* Verbaſculum proliferum C. P. B. flore alteri, altero innato purpureo. ⊙

* Verbaſculum proliferum C. P. B. flore alteri, altero innato luteo. ⊙

* Verbaſculum minus Alpinum umbellatum purpureum. ⊙

* Verbaſculum minus Alpinum umbellatum album. ⊙

Verbena communis flore cœruleo B. pin. Verbenaca recta Dod. ⊙

Verbena Urticæ folio ♃ ⊙

* Verbena nodiflora Matth. ed. B. repens nodiflora Park. ⊙

Veronica mas ſupina & vulgatiſſima B. pin. mas ſerpens Dod. ⊙

* Veronica ſpicata anguſtifolia flore carneo. ♃ ⊙

Vero-

Veronica ſpicata minor B. pin. recta minima Cluſ. Eadem ſpicâ roſeâ. ☉
Veronica pratenſis ſerpyllifolia B. pin. minor ſerpyllifolia Obſ. & Icon. Lob. ☉

Viburnum Matth. ed. B. Lantana Dod. ☉

Vicia Dod. ſativa vulgaris ſemine nigra B. pin. ♀
Vicia maxima dumetorum B. pin. Cracca major Tab. Eadem flore albo. ☉
Vicia Sepium folio rotundiore acuto B. pin. Aphace Fuchſ. ♀
Vicia multiflora B. pin. Aracus Tab. ☉
Vicia ſylveſtris lutea ſiliquâ hirſutâ B. pin. ♀
Vicia lutea foliis Convolvuli minoris B. pin. Orobanche legumen hiſt. Lugd. ♀
Vicia ſemine rotundo nigro B. pin. Craccæ primum genus Dod. ♀
Vicia ſegetum cum ſiliquis plurimis hirſutis B. pin. Aracus .ſ. Cracca minima Obſ. & Icon. Lob. ♀
Vicia ſiliquis glabris flore violaceo. Minima cum ſiliquis Glab. J. B. ♀
Vicia minima præcox Solonienſis. ♋ ♀

Vicia

Vicia Seſamacea Apula Col. ♀
* Vicia maxima annua foliis ſerratis perperàm authoribus, faba Græcorum. ♀

Viola martia purpurea flore ſimplici odoro B. pin. nigra .ſ. purpurea Dod. ☉
Viola martia alba B. pin. candida Dod. deſcr. martia flore albo hort. Eyſt. ☉
Viola martia inodora ſylveſtris B. pin. ſylveſtris inodora Dod. ☉
Viola martia multiplici flore B. pin. Viola flore multiplici Dod. Eadem flore pleno cinereo. ☉
Viola Alpina rotundifolia lutea B. pin. montana prima Cluſ. ☉
* Viola Granadillæ folio. ♃ ☉
* Viola longifolia montana lutea grandiflora Cam. C. P. B. Violæ montanæ 3. alterum Genus Cluſ. hiſt. ☉
Viola tricolor erecta B. pin. aſſurgens tricolor Dod. ☉
Viola tricolor Dod. tricolor hortenſis repens B. pin. ♀
Viola bicolor arvenſis B. pin. Sylveſtris Icon Lob. ♀

Viola lunaria major ſiliquâ rotundâ B. pin. Lunaria Græca annua hort. Eyſt. ☿

Viola lunaria major ſiliquâ oblongâ B. pin. Lunaria Græca perennis hort. Eyſt. ☉

Virga aurea major vel Doria B. pin. Herba Doria Dod. ☉

Virga aurea latifolia ſerrata B. pin. aurea margine crenato Dod. ☉

Virga aurea foliis lævibus non ſerratis. ♃ ☉

Virga aurea anguſtifolia ſerrata B. pin. Solidago Sarracenica Dod. ☉

Virga aurea Americana foliis ſerratis ſubtus nervoſis. ♃ ☉

Virga aurea Americana hirſuta. ♃ ☉

Virga aurea Limonii folio paniculâ uno verſu diſpoſitâ. Mexicana B. pin. app. ☉

Virgo aurea humilis. ♃ ☉

* Virga aurea longis glabris foliis nervoſis. ♃ ☉

* Virga aurea ſerratis foliis hirſutis. ♃ ☉

* Virga aurea Solidago aurea foliis minùs ſpiſſis. ♃ ☉

Vitex

Vitex Dod. foliis anguſtioribus Cannabis modo diſpoſitis B. pin. Eadem flore albo. ☉

Vitis vinifera Fuchſ.
Vitis Corinthiaca.
Vitis laciniatis foliis Cornuti.
Vitis ſylveſtris, Labruſca B. pin. Labruſca Tragi hiſt. Lugd.

Vitis idæa foliis oblongis crenatis fructu nigricante B. pin. Vaccinia nigra Dod. ☉
Vitis idæa foliis ſubrotundis non crenatis baccis rubris B. pin. Vaccinia rubra Dod. ☉

Ulmus Dod. campeſtris & Theophraſti B. pin.
Ulmus montana B. pin. ſecunda Dod. deſcr. latifolia.

Urtica urens maxima B. pin. ſylveſtris aſperior Obſ. Lob. ☉
Urtica urens minor B. pin. minor acrior Obſ. & Icon. Lob. ♀
Urtica urens pillulas ferens B. piu. Romana Fuchſ. ♀ Ur-

Urtica racemoſa Canadenſis. ♃ ☉

Urtica aculeata foliis ſerratis B. pin. Cannabis ſylveſtris hiſt. Lugd. ♀
Urtica aculeata foliis ſerratis altera B. pin. ſylveſtris ſpuria tertia Icon. Lob. ♀

X.

XYRIS Matth. ed. B. Spatula fœtida Fuchſ. ☉

Y.

YUCA foliis Aloës B. pin. Yuca Ad. Lob. part. alt. ☉

Z.

Z.

ZEA, Briſa dicta vel monococcos Germanica B. pin. Monococcon Dod. ♀

*Zea Amylea, vel Zeopyrum Amyleum C. B. P. Triticum trimeſtre Cordi. ♀

Zeopyron Dod. Zeopyrum ſ. Triticoſpeltum B. pin. ♀

PLAN-

PLANTARUM
IN
HORTO REGIO
BLESENSI

Auctore, Contentarum:

Nemini hucusque Scriptarum:

Brevis & succincta Delineatio.

A.

ACANTHIUM lusitanicum: Vulgari Acanthio simile, nisi quod folia habeat magis incana, albidiora, densioreque lanugine obdu-

cta; caules emittit altiores, aliquandò quinos, ſenoſve pedes altos. Capita habet minora, cœterâ vulgari ſimile.

♃ Acanthium Illyrico accedens capitibus minoribus, flexuoſis: Illyrico adeo accedit, ut niſi propiùs intuenti, non animadvertitur differentia, caules emittit, non, tam ſpinis armatos: Capita habet minora, quaſique, cernua & flexuoſa.

♃ Aconitum, ſpica florum Pyramidali: Quoad folia & radices; aliorum Aconitorum more, provenit. Solummodò ſpicam fert, florum cœruleorum cucullatorum, ad baſin latiorem; quæ tandem, in Mucronem deſinit Pyramidalem; atque hac notâ quam facillime à cœteris ſuæ ſortis diſtinguitur.

♃ Ageratum purpureum villoſum: Ab Agerato purpureo Dalechampii hiſt. Lugd. neutiquam diſcrepat, niſi quod folia & caules ejus, villis obducuntur; vulgaris autem purpurei Dalechampii, folia ſunt viridia glabra.

♃ Alaternus, foliis ex albo variegatis: Folia habet partìm alba partìm viridia.

♃

♃ Alaternus, foliis ex luteo variegatis: Hujus folia, partìm ſunt lutea, partìm viridia. Utriuſque folia ſic ſunt variegata, ex calce radici ſuppoſita, aut pari mixtura calcis cum terra ruderoſa, ita plurimæ aliæ plantæ, ex hac, aut ſimili mixtura, variegantur in foliis; ut ſolanum lignoſum ſeu dulcamara, clematis daphnoïdes, Ilex baccifera, liguſtrum, Hyſſopus, Roſmarinum, piloſella & cœteræ ludente ſic natura: inſuper nota ex ſeminibus prædictarum, ſatis; provenire plantas, quæ priſtinum recuperant ſtatum, viridémque induunt colorem; éſtque quaſi morbus ſuperveniens foliis, à ſucco, radicibus attracto; qui pangendo, in perennibus durat; ſatione euaneſcit.

♃ Alcea veſicaria capitis bonæ ſpei: A vulgari Alcea venetorum, in paucis differt, niſi quod caulem producat rectum, firmum, pedalem aut ſeſquipedalem; folia habet firmiora rotundiora & Glabriora florémque majorem, elegantioriſque coloris, cœterâ ei par.

☞ Alchimilla minor: Vulgari Cluſ. in omnibus accedit, niſi partium parvitate. Eſt enim triplo vulgari, in omnibus minor, ejuſque folia dilutius virent.

☞ Alſine inaperto flore .ſ. unicaulis noctiflora: Caules producit triunciales, quorum nodis apponuntur bina foliola, è regione diſpoſita, mucronata, linguæ paſſerinæ ſimilia, in caulium ſummis naſcuntur flores, ſinguli, quinis petalis albis, ad medium bifidis (aliarum Lychnidum & alſinarum more) conſtantes. Qui vix interdiu, lucente ſole conſpiciuntur: propter tenuitatem partium: hoc valde familiare eſt, pluribus Lychnidum. Creſcit in arvis, ſylvoſis humidis & udis.

☞ Alſine ſpergula, ſemine foliaceo nigro; circulo membranaceo albo, cincto: Vulgari ſpergulæ, quoad reliquas omnes partes, adeò eſt ſimilis, ut paſſim in agris arenoſis conſpicere licet, nec diſtingui à vulgari poterit niſi ſemine, quod eſt compreſſum foliaceum ſeu membranaceum, album exteriùs, circularitèr cingens in medio, ſemen nigrum.

nigrum. Reperitur & alia, prope rupellam: cujus ſemen in medio eſt compreſſum fuſcum, circulo paritèr foliaceo albo cinctum; vulgaris autem ſpergulæ ſemen eſt rotundum, Alſines vulgaris modo.

♃ Althæa peregrina, ſemine roſtrato, flore luteo minimo: Hujus folia ſunt quaſi carpini foliorum æmula, bina è regione, caulem ambientia, rotundum, ſeſquipedalem, in cujus ſummis faſtigiis, innaſcuntur floſculi lutei, minores, quibus deciduis ſuccedunt, arctè in circulum diſpoſita, ſemina, aculeis pungentia; Unde roſtrati ei nomen impoſuimus.

♃ Amaranthus ſpica erecta: Eſt flore obſoletè rubro, viridique, cujus utriuſque ſemen eſt nigrum; vel ſpica eſt erecta, cujus ſemen eſt album.

♃ Amaranthus auriculatus: Species eſt Amaranthi; quinquageſimo ſeptimo anno, ſupra milleſimum, ſexcenteſimum, detecta; & ex Americæ inſulis adjacentibus, in Angliam delata. Cujus ſingula folia, ad appoſitionem, bina habent foliola Lobata, ſ. auriculata, qua nota, ab

omnibus aliis (multos ante annos) detectis, differt éſtque à flore .ſ.ſpica florum, aut luteus, aut ſtramineus, ruber aut phœniceus, ſpicatus vel criſtatus.

℥ Amygdalus pumila: Bipedalem habet altitudinem, vix pollicarem craſſitiem, obtinent ejus caudices, in quorum ſummis alis, ſub exitum hyemis proveniunt floſculi Coloſſini coloris, minores; Amygdali majoris more; quibus ſuccedunt paritèr fructus parvi triplo minores, Amygdali vulgaris fructu.

℥ Anagallis omnium minima: Planta eſt in humidis proveniens, vix biuncialis, floſculos gerit cœruleos, vulgaris Anagallidis triplò minores; capſulas ſeminales paritèr minores.

℥ Anonis purpurea non ſpinoſa: A vulgari nulla nota differt niſi quod ſpinis Careat.

℥ Anonis montana præcox, purpurea fruteſcens: Suffrutex hic, eſt Bipedalis, Trifolii ſpecies, perpetua fronde virens, flores fert pendentes leguminoſos .ſ. Papileaceos, elegantèr purpureos: exeunte hyeme floret; quorum ſingulis ſuccedunt

cedunt ſingulæ ſiliquæ, hirſutæ biuncia-les, natricis Plinii ſiliquis, multò craſſi-ores, ſemina in ſe continentes majora; deliquia (cœterarum Anonidum, ſemi-num more) patientia, qua nota ſpecifica, indigitantur Anonides, à cœteris, trifo-liorum claſſibus. Perperàm ergò Pari-ſienſibus Botannicis, cicer perenne (me diſcipulo ſub Veſpaſiano Robino) di-cebatur.

♃ Anonis purpurea, erecta annua mitis: Hujus caules ſunt Bipedales nonnun-quam Tripedales erecti, in quorum ſum-mis faſtigiis ſpicatìm, proveniunt floſ-culi. Dilutè purpurei, quibus fatiſcen-tibus ſuccedunt, Siliculæ lentiformes, erectæ, vulgaris Anonidis ritu, ſemina in ſe continentes, deliquium patientia, quæ eſt nota eſſentialis, & Generica, A-nonidum omnium. Et præcipuè, hac nota ab omnibus aliis trifoliis diſtingui-tur: ut clarius docebimus, in univerſali noſtra Hiſtoria Plantarum, per Tabulas cognationis diſpoſitarum.

♃ Apocynum Camadenſe foliis Andro-ſemi: Planta eſt radice longè latéque

per terram ſubtûs repente, prædita; caules ſuggerente Bipedales, quibus binatìm, adhærent folia Androſæmi foliorum adinſtar: in caulium ſummis enaſcuntur flores obſoletè rubri. Quoad formam, aliorum Apocynorum floribus accedentes. Quibus euanidis ſuccedunt, ſiliquæ oblongæ, in medio craſſæ, membranaceæ, Aſclepiadis aut Apocyni vulgaris, ritu, ſemina continentes compreſſa, ſubrufa, intermixta, materiæ cuidam pappoſæ ſuccum lacteum Apocynorum more, quælibet pars rupta fundit.

♃ Apocynum Canadenſe, anguſtifolium flore aurantio: Hæc Apocyni ſpecies vix Bipedalem, excedit altitudinem, folia gerit, anguſta longiora multò Apocyno ſurrecto umbellifero Cornuti; in caulium ſummis faſtigiis, flores aureos aut aurantii coloris gerit. Quibus ſingulatìm, ſuccedunt ſiliquæ membranaceæ, pappoſæ materiæ, intermixta ſemina compreſſa rufa (cœterorum ritu) gerentes.

♃ Apocynum Virginianum, flore herbaceo ſiliqua longiſſima: Hujus folia

ſunt

ſunt lata glabra Bedelzar Arabum foliis, magnitudine accedentia. Caules habet ternos quaternos,pedes altos:ſubrubros, in ſummis alis caulium proveniunt floſculi herbacei, ejuſdem cum cœteris ſuæ ſortis,formæ ; ſed multò minores,quibus flaccidis ſuccedunt ſiliquæ, Dodrantales tenuiſſimæ, pappoſam materiam, & ſemina compreſſa, aliorum more continentes. Lacteſcit etiam aliorum Apocynorum & Aſclepiadum modo.

♃ Argemone capitulo longiori,glabro : Folia habet magis diſſecta , quam ſunt papaveris Rhæadis folia, minùs autem, quam ſunt Argemones;capitulo hirſuto. Florem fert Phœniceum, cui ſuccedit capitulum biunciale, glabrum, per loculamenta triangularia , continens minuta ſemina, aſpera, aliorum papaverum tum Argemonarum modo.

♃ Artemiſia foliis variegatis: Sola variegatione foliorum,à vulgari Artemiſia differt.

♃ Arum Indicum bicubitale, amplo flore:Folia habet ampla,cannacori æmula,è quorum gremio,ſurgit caulis bicubitalis, in

in ſummo gerens florem amplum, album Lilii candidi flore multò majorem, uniſolium cui deciduo, ſuccedunt, racematim, in ſpica, grana plurima rubra, ſpicatim ſeinvicem contingentia, vulgaris (ut paucis dicam) ritu, diſpoſita.

♃ Aſter ſerotinus, anguſtifolius, Canadenſis flore cœruleo: Planta eſt longè latéque ſerpens, caules emittit tripedales, foliis anguſtis glabris, ſaturè virentibus, è regione diſpoſitis onuſtos. E ſummis caulium exeunt floſculi, cœrulei; ſub exitum æſtatis, aut ineunte hyeme, aliquando tardiús: cœterorum more, in pappum abeuntes.

♃ Aſter præcox, flore cœruleo majore: Hæc, Canadenſis ſeu exotici aſteris, eſt ſpecies, caulis humilior flos major (ſi comparetur cum ſequenti congenere) media æſtate floret cœrulei coloris eſt.

♃ Aſter elatior flore cœruleo minore, Bellidis minoris: Hic longe ſub terram hac, illac, repit ſerpitve; caules emittit tripedales, foliis anguſtis glabris, ſaturè virentibus, è regione diſpoſitis, onuſtos. Flores ſuſtinentes, (quoad magnitudinem)

nem) Bellidis minoris, ſed cœrulei coloris, verſus hyemis initium (cœterorum more) in pappos abeunt.

☞ Aſter Tripolii flore latifolius: Hic omninò accedit minori, Narbonenſium. Si non quod folia producat glabriora & latiora, floreſque paulò majores, deniq; tota Planta eſt major, quoad omnes ipſius partes.

☞ Aſter Conyzoïdes Monſpelienſium: Eſt planta Bipedalis, cujus folia ſunt villoſa .ſ. lanuginoſa, Conyzæ in modum, quin & contrectata ejus folia, Conyzæ odorem ſpirant. In ſummis, caulium faſtigiis, flores fert luteos, multos, imo ad alas foliorum, quibus deciduis, ſuccedunt ſemina ſolida Anguloſa, qua nota debet jure merito appellari, Chryſanthemum Conyzoïdes Monſpelienſium. Conyzoïdes apte dicitur, quia Conyzæ habet odorem. Monſpelienſis dicitur quia frequentèr circa oppidum illud reperitur. Omnis aſteris flos, abit in pappos, qua ſola nota eſſentiali & generica, differt aſter, à Corymboſis ſeu Corymbiferis, quippe producentibus, flores

flores etiam ſtellatos, quemadmodum aſter, ſed ſemina ſolida, ergò potiore jure Chryſanthemum Conyzoïdes Monſpelienſium denominabitur, ut clariùs demonſtrabimus in tabulis noſtris cognationis. Conyzæ fundit odorem, ſed nec odor nec ſapor, nec color indigitant plantas, per ſe, ſed per accidens tantúm: ad deſignandas ſpecies, ſubalternas inſerviunt. Idem dicendum de aſtere luteo, foliolis ad florem rigidis, itèm de aſtere cernuo col. ſeu foliis circa florem mollibus; paritèr de aſtere Attico ſupino Dod. & multis aliis hujus Farinæ, ab Authoribus, aſterum claſſi aſſignatis, ut prolixius in Tabulis affinitatis ſeu cognationis, clare demonſtrabimus.

♃ Aſter ramoſus annuus Canadenſis: Hæc planta folia producit oblonga, in margine ſinuata, ſupernè hirta, caules edit Bipedales ramoſos, flores albos, Bellidis inſtar. Qui abeunt in pappos: ergò aptiùs retulimus, ad ſuæ claſſis conſortes. Nempe pappeſcentes, vide Bellis ramoſa umbelliſera Cornuti in horto Regio Bleſenſi.

Atriplex

♃ Atriplex folio haſtato ſeu Deltoïde: Hæc atriplicis ſpecies paſsìm in foſſis. Suburbanis provenit, folio eſt Triangulari acuto, mucronato, haſtæ cuſpidi ſimili: caulibus procumbentibus, pedalibus & ſeſquipedalibus; pediculis longis innixo. In quorum ſummis alis, ſemina multa proveniunt: ſingula binis foliolis (ut in Atriplice ſativo, Polygani folio, marino, aliiſque) incluſa.

♃ Atriplex procumbens folio ſinuato craſſo: Planta pedalis, procumbens, folia habet ſinuata in margine, ut plurimùm & caules & folia ſunt rubra; ſemine onerantur ſummi caules, racemoſo, cœterarum ſylveſtrium more.

B.

ꝺ BELLIS pratenſis duplici petalorum ſerie: Non aliâ ratione differt à Bellide pratenſi vulgari, niſi quod natura luxuriante, duplicem petalorum latorum, umbonem luteum cingentium, ſeriem profert.

ꝺ Bellis hortenſis flore Herbaceo: Hortenſi in omnibus convenit, niſi quod florum petala, in folia herbacea degenerent.

ꝺ Bellis hortenſis flore prolifero: Hæc notu facilima, ex primo intuitu, quia plures flores exeuntes, ex uno, (Calendulæ proliferæ more) producit.

ꝺ Bellis ramoſa umbellifera Cornuti: Cum ejus flos abit in pappos, ad Aſteris claſſem retulimus vide ſuprâ, Aſter ramoſus annuus Canadenſis.

ꝺ Bellis Virginiana denſiori folio: Folia habet latiora, ſerrata, denſiora, quam ulla, Bellidum ſylveſtrium majorum: majoréſque fert flores. Forte Bellis alpina Caſ. Banh. in prodr. ꝺ

℥ Bellis radice repente, foliis latioribus ſerratis: Hæc omnium Bellidum majorum, folia habet latiora magiſque ſerrata, flores itidem maximos, radices longe latéque ſubter terram, repentes.

℥ Bellis montana minor & major Tanaceti folio: Vide Tanacetum montanum inodorum flore majore & minore in horto Regio Blæſenſi aucto. An non potiore jure Bellis Tanaceti folio, deſignatur. Examinent quicunque voluerint.

℥ Beta radice vulgaris, coſta ampla aurea: Hujus folia ſunt Platyneura, aliquando aurea, aliquandò rubra, nonnunquam alba.

℥ Blattaria folio verbaſci, flore luteo amplo: Hæc folia habet verbaſci, Incana mollia, mucronata, è quorum gremio exit caulis, Bipedalis aut Tripedalis, in faſtigio, gerens flores luteos, amplos verbaſci inſtar. Quibus tamen deciduis ſuccedunt, capſulæ bivalves, ſeu binis loculamentis, diſtinctæ. Non in Pyramidem verbaſci capſularum more, ſed ſunt craſſæ, rotundæ; qua nota ſola à verbaſco diſtinguitur Blattaria.

℥

♃ Blattaria purpurea radice perenni: Folia edit incanè viridia, ſinuata, floſculos cœruleos, quatuor petalis conflatos, quibus deciduis ſuccedunt capſulæ, prioris capſularum æmulæ, ſimiliaque in ſe, continentes ſemina. Datur & hujus ſpecies foliis majoribus, Symphyti foliis, quam proxime accedens, quæ flores edit elegantèr carneos vel gilvos. Capſulaſque edit minores; natura eſt adeò intenta productioni, hujus: longè latéque rependo ut vix in Gallia, multò minùs in Anglia, ſemen ſuum perficeat.

♃ Blitum Polyſpermon, folio elatines ſubrotundo: Hæc Bliti ſpecies folia habet, hirta, ſeu villoſa, ſubrotunda, quemadmodùm elatine folio ſubrotundo; binatìm è regione, cingunt, caules pedales & ſeſquipedales; ad alas, enaſcuntur racematìm ſemina multa ſubnigra aut fuſca, lata tumidiuſcula: viridibus capſulis minimis (Atriplicis aut Bliti modo) contenta.

♃ Brunella Alpina, glabro folio, anguſto, integro: Folia edit mucronata, glabra, anguſta rigida, caulibus ſemi-

pedalibus

pedalibus è regione adnaſcentia: quorum ſummis apicibus verticillatìm, densè diſpoſiti, innaſcuntur flores, galeati cœrulei, nonnunquam purpurei, qui decidui, capſulas verticillatas, in ſpicam degenerentes, relinquunt: aliarum prunellarum more.

♃ Brunella Bætica annua foliis laciniatis: Folia inferiora profert ſinuata, per terram ſtrata, caulescente planta: folia edit laciniata, è regione diſpoſita. Summis caulibus innaſcuntur, verticillatìm ampli flores cœrulei, qui exſiccati, relinquunt ſpicatìm, ſeu denſiſſimè diſpoſitas capſulas verticillatas quinquefariam in margine partitas; vide Meliſſa Bætica in horto Regio Bleſenſi aucto, cum qua nullam habet affinitatem ſed omninò Brunellæ accedit. Ut clariùs poſtea docebitur.

♃ Bugloſſum luſitanicum bullatis foliis: Hæc Bugloſſi ſpecies, folia (ſi ſeratur autumno) exeunte hyeme profert viridia, bullis albis elegantèr conſperſa: è quorum medio, exit caulis, dodrantalis, aliquando pedalis, per terram ſtra-

tus, in ſummitate, ſuſtinens flores elegantèr cœruleos, parvos quadripartitos, quibus deciduis, ſuccedunt ſemina, viperini capitis æmula, aliorum Bugloſſorum more. Hujus eſt, & alia varietas, priori in omnibus ſimilis, niſi, quod flores edit rubros ſtriis albis, elegantèr pictos.

♃ Bugloſſum procumbens annuum, pullo minimo flore: Hujus cauliculi Bipedales per terram ſtrati ſunt, folia gerunt villoſa, floreſque minimos; pulli ſeu obſcurè ferruginei coloris; Bugloſſi in modum, undè aptiùs Bugloſſum minimum procumbens, pullo flore, dicitur. Quam Echium; quia Echium, ſeu Lycopſis, flore donatur corniculato integro: qua nota, ab invicèm diſtinguentur, Bugloſſum: Echium ſeu Lycopſis, ut in Tabulis noſtris cognationis, clarius patebit. Vide Echium procumbens, floſculis atrorubentibus; in horto Regio Blæſenſi aucto.

♃ Buxus foliis aureis: Sola variegatione foliorum, à vulgari differt: quod in multis aliis arboribus, fruticibus, & ſubfruticibus, plantiſque, accidit. C.

C.

☞ CAchrys peucedani foliis, ſemine fungoſo ſulcato: Hujus folia ſunt diſſecta, peucedani inſtar, caules Bipedales, umbellas ſuſtinent luteas, quibus fatiſcentibus, floribus ſingulis, ſuccedunt bina ſemina, ſtriata, fungoſa ſubſtantia incluſa, oblonga, nucis Myriſticæ, ferè magnitudine, Melonum inſtar ſulcata, à lævitate una, ſemine ſulcato lævi, altera ſemine ſulcato aſpero dicitur. Hinc, duæ ſpecies, ſola ſeminis fungoſa ſubſtantia, lævi & aſpera ſulcata, diſtinctæ, utraque ſpecies, folia gerit peucedani. Vide Libanotis Cachriſera, ſeu Cachrys verior, in horto Regio Bleſenſi aucto; cujus folia ſunt ferulacea, aut fœniculacea ſemina itidèm ſtriata gerit, ſubſtantia fungoſa, plana, incluſa, nucis Myriſticæ paritèr ad formam, & quaſi magnitudinem.

☞ * Calceolus mariæ luteus: Folia vulgaris edit, flores ejuſdem formæ niſi

quod luteos gerit, éſtque Helleborines ſpecies, à forma floris, ſic denominatur Calceolus mariæ.

♃ Campanula .ſ. viola pentagonia perfoliata : Hujus folia ſunt ſerrata, ſubrotunda, binatim ex adverſo, caulem ambiunt Dodrantalem: ita,ut ab ipſo caule perforari, videantur. In ſummis, caulium, naſcuntur flores, Campanulati, violæ Pentagoniæ vulgaris æmuli, quibus ſuccedunt Siliculæ pares, Triplici loculamento diſtinctæ, ſemina Pellucida, ſubrufa, paria,cum vulgaris, Campanulæ ſeminibus : continentes.

♃ Campanula .ſ.viola Pentagonia,folio oblongo latiori : Hæc vulgari perquam ſimilis, niſi quod folia edat longiora, latioraque : totaque planta eſt major vulgari. Reperimus & vulgarem, intra ſata, Regiæ Chambort, flore cœruleo pleno, utraque hæc ſpecies lacteſcit aliarum Campanulatarum more. Vide Onobrychis arvenſis in horto Regio Bleſenſi aucto.

♃ Carduus Leucographis .ſ. mariæ hirſutus,capitulo minore: Hic Carduus folia

edit

edit longiora, & anguſtiora, obſcurè maculata, ſecundo anno caulem emittit Bipedalem, aut Tripedalem, in cujus cimis naſcuntur capita, minora, purpureos ferentia, tubulos, ſeu, flores; quibus fatiſcentibus, ſuccedunt, ſemina rotunda, Craſſa, & ſolida: annua eſt planta.

℔ Carduus mariæ non maculatus: Carduo vulgari mariæ, in omnibus accedit: niſi quod folia edit maculis .ſ. lineis albis, deſtituta.

℔ Carduus Leucoii foliis: Planta eſt pedalis aut ſeſquipedalis, cujus caulem ambiunt ex adverſo diſpoſita, bina folia Leucoii murorum ſeu Keiri inſtar: ſupernè viridia, ſubtûs ſeu infernè incana, ad quorum appoſitionem ſunt binæ ſpinæ. Floſculos gerit minimos, purpureos, qui in pappos reſoluti, relinquunt ſemina Craſſa, magna ſatis, pro plantæ tenuitate.

℔ Carduus exoticus minor, albis maculis notatus: Hic autumno ſeſe ſerit. Atque per hyemem folia per terram ſtrata gerit; laciniata, elegantèr lineis albis notata. Caules pedales & ſeſquipedales,

caputque parvum rubrum fert, cui pappeſcenti, ſuccedunt, ſemina glutinoſa, pappo, incluſa. An Carduus Galactites Joh. Bauh. paſsim ſecus vias, prope Monſpeſſulum reperitur.

♃ Carduus Polyacanthæ æmulus, ſeu alatus; Polyacanthæ, capitulis æmulus: Hic Carduus ad Tripedalem, nonnunquàm ad quadripedalem aſſurgit altitudinem. Caulem habet, quatuor alis undulatis cinctum, folia obſcuriùs virent, in margine criſpa, rarioribus & mitioribus ſpinis, quam Polyacantha, armatur: Hic Carduus paſsim, ſecùs viarum margines, circa Londinum, Blæſas, pariſios, & cœtera (id genus) loca, provenit copioſe. Floſculos capitatos, in caulium faſtigiis gerit, rubicundiores pauloque majores, Polyacanthæ Theophr. ſaturatius rubeſcunt flores: nonnunquam variat flos, colore albo, cœterorum carduorum rubro colore, præditorum; more.

♃ Carduus Sphærocephalus annuus Bæticus minor: Hic cœteris Carduis Sphærocephalis, minor eſt. Vix Bipedalem excedit

excedit altitudinem: flores cœruleos minores, globulatìm fert; capita etiam minora, ſed ſemina multò, cœteris ſuæ familias ſeminibus majora fundit.

ℭ Caryophyllata montana minor, flore nutante purpureo: Hæc à Caryophyllata majori vulgari; flore purpureo nutante deſcripta, differt, in hoc, quod folia edat glabra, minutiúſque diſſecta, totaque planta minor eſt.

ℭ Caryophyllus omnium humillimus & minimus: Hic in montibus Pyrenæis, & alpibus provenit, pediculos profert unciales & biunciales, ſuſtinentes floſculos minutes, rubros, laciniatos, folia gerit tenuia, muſcoſa, undè muſcoſus appellari, meretur.

ℭ Caryophyllus marinus minimus flore globoſo: Hic, folia tenuia, ternas quaternas uncias, longa, flores globoſos, omnium hujus generis, minimos fert, eorum qui in maritimis naſcuntur. Arearum aut Ambulacrorum margines, hoc conſeri (cum propter perennitatem foliorum, tum amœnitatem florum per æſtatem durantium) ſolent.

℥ Centaurium purpureum minimum: Caules biunciales vel triunciales ramatìm fert, flores purpureos, ſemina & capſulas ſeminales, centaurii vulgaris minoris modo, fert. In humidis (circa ſtagna) locis, provenit.

℥ Centaurium luteum minimum: Hæc plantula etiam in humidis vegetat: vix Palmaris eſt; floſculos gerit luteos: capſulas, plurima minuta ſemina (vulgaris more) in ſe continentes, habet: Forte luteum novum colum.

℥ Chamæciſtus halimi folio oblongo: Hic ſubfrutex ad pedalem utplurimùm aliquando Bipedalem aſſurgit altitudinem, folia profert oblonga, mucronata, incana, halimi inſtar. In Glareoſis & Arenoſis, circa caput Cenomanorum, copiosè conſpicitur flores fert luteos, quibus labentibus, ſuccedunt capſulæ, triangulares; minuta multa ſubrufa ſemina in ſe continentes cœterorum Chamæciſtorum ad modum. Vide Petrum Belonium Cenomanum.

℥ Chamæciſtus annuus erectus: Hic caules rectos palmares & ſeſquipalmares,

res, producit; floſculos minimos luteos, quinis petalis, luteis, donatos profert. Quibus decidentibus, ſuccedunt (aliorum more) capſulæ triquetræ. Ad cellam Eremitarum, Milliari, Bleſis diſſitam; prope pagum dictum la Chauſſé & circa pictavium copiosè reperitur; eſt minor ſpecies Ciſti, folio Salicis Cluſ. ad pedalem altitudinem rectè aſcendentem, circa pictavium reperimus.

♃ Chamæmelum inodorum; ſeu cotula inodora ſemine nigro: In ora maritima Armoricæ, ſatis frequens occurrit; hæc planta, cujus folia, ſunt Cotulæ vulgaris fœtidæ foliis, magis diſſecta, Fœniculi, quaſi modo, obſcuriùs virent: quin & florum petala, ſunt multò ampliora, longiora; diſcum luteum cingentia. Quibus fatiſcentibus ſuccedunt ſemina nigra, éſtque hujus ſpecies, quæ in hortis colitur, & Cotula inodora flore pleno dicitur, quæ pangendo, propagatur. Semen itidèm habet nigrum, quemadmodùm, obſervavimus & penes nos impreſentiarùm habemus, plurimæ ſunt plantæ præcipuè annuæ, etiamſi flore pleno

pleno donentur, tamen ſemen producunt. Ut clariùs Enucleabimus in Hiſtoria generali, ſeu univerſali, Plantarum: per Tabulas cognationis, ſeu affinitatis, diſponendarum.

℔ Chelidonium majus, foliis tenuiſſimè diſſectis: Sola foliorum inciſione, multò tenuius, diſſectorum à Chelidonio foliis quernis differt: nam florem fert, laciniatum itidèm; quare ex hujus ſemine, creditur eſſe degener planta. Nam ſemen ex hoc collectum, & à me ſatum: Chelidonium foliis quernis laciniato flore, produxit.

℔ Chryſanthemum Hiſpanicum, tenuifolium: Folia habet minùs tenuiter diſſecta, quam Achillea montana, floremque duplò majorem: utrumque & hoc & Achillea montana: Chryſanthemum, ferulaceis foliis nobis dicitur.

℔ Chryſanthemum Canadenſe latifolium altiſſimum: Ad quinos ſenoſve octonos nonnunquam pedes, aſſurgit caulis, craſſus, ſupra pollicarem craſſitudinem quem cingunt alternatim, folia. rugoſa oblonga, lata, plantaginis majoris

(quoad

(quoad formam) majora. In ſummis caulium faſtigiis, fert flores luteos diſcum luteum ambientes: (numiſma, dimidium taleri) vix æquantes; quibus exſiccatis ſuccedunt ſemina anguloſa, quaſi floris ſolis ſemina; ſed minora.

℥ Chryſanthemum Canadenſe latifolium humilius: Hoc vix Tripedalem excedit altitudinem, foliis, priori, quam proximè accedit, flores multò minores; ſemina paritèr minora: ejuſdem tamen figuræ, cum præcedentis ſemine.

℥ Chryſanthemum Americanum laciniato folio majus: Caules producit tricubitales, ad quorum nodos, apponuntur folia laciniata, magna, dilutè viridia: flores fert Corymbiferos luteos, quibus deciduis, ſuccedunt ſemina quadrangula, oblonga; quæ declarant, non eſſe Doronicum. Cum omnis Doronici flos abit in pappos.

℥ Chryſanthemum Americanum laciniato folio, obſcuriùs virenti, minus: Hoc à priori differt quod ſit humilius, folia gerit minora, magis laciniata, & obſcuriùs virentia, floreſque minores, utrumque

utrumque & ſuperius, & hoc ſemen fert ſolidum quadrangulum; neutrum pro Doronico haberi debet. Ut vult Johan. Parkinſon, cum utriuſque ſemen ſit ſolidum, Chryſanthemi more; multò minùs, eſt Aconitum ut vult Jacobus Cornutus; omnis Aconiti flos eſt Cucullatus, producens ſiliquas membranaceas, quaternas quinas: multa minuta ſemina rugoſa, in ſe continentes .ſ. nigra .ſ. rufa, ergò aptius, denominatur Chryſanthemum, & redarguendi ſunt authores, qui non norunt diſtinguere inter aſterem, Doronicum, Conyzam, Helenium & cœteras pappeſcentes, & Corymboſas: ſolida producentes ſemina. Atque hæc dixiſſe impræſentiarùm ſat. eſt.

♃ Chryſanthemum Canadenſe minus, alato caule: Hæc Chryſanthemi ſpecies, folia habet glabra, anguſta, ſecundum longitudinem caulium, producta, adeò ut alatus caulis dici poſſit. Caules habet Bipedales Tripedales, flores ſuſtinent luteos: quibus deciduis ſuccedunt, ſemina ſolida quadrangula, minimè autem in pappos abeunt Aſteris inſtar.: ergò malè

malè à Jacobo Cornuto Aſter alatus luteus dicitur, ut videre eſt in horto Regio Bleſenſi aucto, & in Parkinſono ſub Aſteris nomine, quam bene Judicent lectores.

♃ Chryſanthemum Canadenſe bidens, alato caule: Omninm Chryſanthemorum exoticorum, altiſſimum eſt hoc; caules edit ſenos ſeptenos, nonnunquam duodenos excedentes pedes, quibus, adhærent folia aſpera, oblonga, ad appoſitionem, alas conſtituentia, ſecundum longitudinem; caulis. Undè alato caule dicitur: flores in cimis caulium, Corymbatim fert, luteos; umbonem luteum cingentes, valdè exiguos, pro magnitudine plantæ vix Calendulæ flores æquant: quibus fatiſcentibus ſuccedunt ſemina, ſubrotunda compreſſa, binis aculeis, in ſuperiore parte ſeminis, prædita. Undè bidens (adinſtar Chryſanthemi Cannabini) appellandum, judicavimus. Malè ſub Helenio Canadenſi in horto Regio Bleſenſi. Quædam horum nominantur ut hoc Helenium Canadenſe prælatum, alato caule Aſter virgineus

gineus luteus, membranaceo caule Park. ex ſemine. Chryſanthemum Canadenſe bidens, melius nominatur; legentium & perpendentium, eſto judicium.

♃ Chryſanthemum Cannabinum Americanum: A noſtrate, cognitu facile, quippe caules edit quadripedales, ſubrubentes firmiores & rectiores, floſculos, ex multis ſtaminulis luteis, conflatos habet: & ſemina bidentia, multò, vulgaris Chryſanthemi Cannabini, ſeminibus majora; veſtibus paritèr adhærentia, pari modo ut noſtras fert: malè in horto Regio Bleſenſi pro Eupatorio, cum authoribus aliis ſcribitur. Ut poſtea ad Oculum demonſtrabitur.

♃ Cicer arietinum, fructu nigro: Solo fructus colore, à Cicere arietino fructu rubro, & fructus ipſius parvitate, & figura, utriuſque enim flores ſunt rubri.

♃ Clematis peregrina flore parvo: Hæc Clematidi peregrinæ in omnibus accedit: niſi quod flores profert duplò minores, aut rubros aut albos: ſemina paritèr minora, ejuſdem cum priorum ſeminibus, formæ.

♃

♃ Clematis Tetraphyllos Americana: Ex inſpectione quaternorum foliorum cruciatim diſpoſitorum, citò dignoſcitur; nec ſemen nec florem nobis conſpicere licuit: ſcandere (adhibito baculo) obſervavimus.

♃ Clematis Hederæ folio: Planta eſt ſcandens & circa perticam ſeſe involvens ad altitudinem quinorum pedum, plus minuſve aſcendit; cujus folia ſunt Hederæ, æmula, latiora; flores fert minores, herbaceos; fructum nunquam vidi. Malè citatur in horto Regio Bleſenſi. Convolvulus Hederaceus perennis, cum Convolvulo propriè dicto, lacteſcente, florem unifolium gerente, nullam habet affinitatem, niſi quod ſcandit.

♃ Cochlearia, armorica flore parvo, ſubpurpureo: Hæc minor eſt ſpecies, Batavicæ; folia ei ſimilia edit, ſed minora, anguloſa, flores etiam minores, ſubpurpureos, ſeu dilutè purpureos: quibus ſuccedunt, Siliculæ bivalves, minores aliarum Cochleariarum more, ſemina in ſe minora paritèr continentes.

♃

℥ Conſolida regalis hortenſis, flore minore verſicolore: Seu Delphinium minus verſicolor: folia habet, hortenſis Delphinii vulgaris foliis tenuius diſſecta, flores minores rubros, Striis nigris maculatos; & eo ipſo nomine, ab aliis quam facillimè diſtinguitur.

℥ Convolvulus minor argenteus ſ. incanus radice repente: Radices ut cæteri, ejus claſſis, ſub terram, repentes, habet: folia producit ſubrotunda, incana: flores, Coloſſinos, ſeu purpureos & albos, quorum ſingulis ſuccedunt ſemina, nunc plura, nunc pauciora, anguloſa: paſsim, ad oram maritimam, circa Rupellam, ſpontè provenit.

℥ Conyza acris, annua, alba: Planta eſt Bipedalis, cujus folia ſunt oblonga, ſerrata in margine; villoſa: flores gerit ramatim, in ſummis caulibus, albos: erigeri, ſeu Conyzæ acris cœruleæ, more, quin Acria Guſtui, ejus folia maſticata; ſtatim apparent. Flores decidui in pappos abeunt, & vento Aſportata ſemina, copioſè ſeſe ſerunt, per agros, tam in ſylvis quam vinetis Galliæ, Angliæ paſsim.

Digitized by Google

sìm. Aſter Canadenſis annuus, malè in horto Regio Bleſenſi aucto nuncupatur, quin & pariſienſibus, Aſter Canadenſis (me diſcipulo) dicebatur, ſed cum non ſtellatum gerit florem, ſed Erigeri aut potius Conyzæ cœruleæ, cui ſapor acris accedit, aptius (mea opinione) Conyza acris alba; quam appoſitè Canadenſis dicitur (cum paſsìm reperire eſt) Judicent lectores.

ℭ Cotyledon, radice tuberoſa, longa, repente: Hæc Cotyledonis ſpecies priori, ab Authoribus diſcriptæ, quam proximè accedit, niſi quod folia gerit majora, ſpiſſiora, nec circularitèr, ſcutellam .ſ. Cotyledonem effingunt: Sed juxta, pediculi, ſuperiorem partem aperta; nec circularitèr clauſa, prioris & majoris modò, quin & radicem, non promit rotundam & tuberoſam, rapi inſtar, ſed longitudinalitèr porrectam, & ſub terram repentem, à qua exeunt fibræ multæ exiles, quibus nutritur radix. Flores & ſemina vulgaris modo, habet.

Cynogloſſum folio molli, incano, flore cœruleo ſtriis rubris variegato: Hæc Cynogloſſi ſpecies, folia vulgaris foliis, multò producit minora, mollia, incana. Caules profert Bipedales, quorum ſummis innaſcuntur: flores dilutè cœrulei, ſtriis rubris variè ducti: quorum ſingulis deciduis, ſuccedunt quaterna ſemina (vulgaris & aliorum more) aſpera veſtibus adhærentia, multò minora, rotunda: paſsim ſecus vias; circa Blæſas provenit.

D.

DENTARIA Monophyllos, Anapodaphyllon pariſienſibus: Nominabatur; planta eſt vernalis, caulem è terra Palmarem erigens, cujus apici innitur folium ſub exortum complicatum proceſſu temporis extenſum, orbiculariter, rotundum, ſinuatum, in margine, ad caulis prædicti exortum, è terra prorumpit pediculus, triuncialis plus minuſve;

Digitized by Google

nuſve; ſuſtinens. Florem, quinis petalis albis, donatum; cui deciduo ſuccedit capſula, nucis Avellanæ magnitudine, multa minuta in ſe continens ſemina.

♃ Dentariæ affinis, Echii flore, capſula rotunda: Hæc planta ineunte vere, folia promit, ſub exortum, Aſtranſiæ aut Diapenſiæ ſimilia, tandem Dentariæ referunt folia, inter quæ exit caulis Palmaris, flores ſuſtinens quaternos, quinos, albos. Echii floribus ſimiles; quibus evaneſcentibus ſuccedunt capſulæ rotundæ Anagallidis capſularum inſtar: ſemina ſolida in ſe continentes, aſperulæ odoratæ ſeminibus quaſi paria.

♃ Digitalis Virginiana Lyſimachiæ facie, foliis ſubrubentibus, flore violaceo minore: Hæc Digitalis ſpecies, caules promit Bipedales ſubrubentes, quos ambiunt per intervalla bina folia mucronata, Limonii majoris (primo ortu) foliis haud abſimilia, inter quæ ſeſe erigit caulis cubitalis, lignoſus, rotundus, purpuraſcens: cui ad ſingula genicula, ex adverſo bina adnaſcuntur folia, Lyſimachiæ quodammodò, æmula, arctè adeo

Digitized by

caulem cingunt, ut perfoliata planta, dici meritò poſſit: ex alis ad genicula ſuperiora, enaſcuntur pediculi, alios procreantes, quibus innaſcuntur flores, calice & forma Dactylothecæ ſeu digitalis ferrugineæ minoris, æmuli, dilutè violacei, ore non alitèr, quinque inciſuris, tribus propendentibus, duabus erectis, diviſo; per margines hirſuto: in fundo autem tubi ſeu floris, quatuor ſtaminula arcuata latent, horum ſingulorum extremo, aliarum digitalium more, duo apponuntur teſticuli rubelli, lentiformes, in ſuperiore floris parte. Eſt autem ſtilus (propendens verſus inferiorem partem) luteus; lanuginoſus. Succedunt, poſt evanidum florem Thecæ bicapſulares in mucronem acutum bifurcatum deſinentes: continentes in ſe ſemina infinita minuta, digitalium omnium more.

♃ Digitalis foliis maximê ſerratis: Vide, galeata & verticillata, Perſicæ foliis, digitalis æmula: poſt Horminum & Hyſſopum infrâ.

♃ Digitalis latifolia flore ferrugineo: Hæc Digitalis, folia profert oblonga, ſulcata,

Digitized by Google

ſulcata, nervoſa, multò digitalis ferrugineæ anguſtifoliæ foliis latiora, è quorum gremio ſeſe erigit caulis Bipedalis, ſuſtinens faſtigiatim complures flores, (Dactylothecæ vulgaris, inſtàr, elegantis ferruginei coloris) majores; in quorum ſingulorum locum ſuccedunt Thecæ, bicapſulares mucronatæ, & acutè bifidæ, craſſiores; digitalis anguſtifoliæ, flore ferrugineo, capſulis. Semina itidèm paria, hinc inde in duplici loculamento, rufa, ſed majora, continentes.

♃ Doronicum foliis ſubtûs lanuginoſis: Hujus caules ſunt pedales, folia ſuprà obſcurius virentia, ſubtûs incana, oblonga: flores fert luteos ſtellatos (aliorum more) ſeminaque abeuntia in pappos.

E.

ECHIUM procumbens, floſculis atro-rubentibus, procumbens annuum: Vide Bugloſſum pullo minimo flore; in horto Regio Bleſenſi aucto.

Eruca tanaceti foliis: Hæc folia profert incana viridia; tanaceti foliis (quoad diſpoſitionem) haud abſimilia, ex quorum gremio exit caulis, ſeſquipedalis, in cujus ſummis, naſcuntur floſculi lutei, quatuor petalis conflati, quibus pereuntibus ſingulis, ſingulæ, ſuperveniunt ſiliquæ biunciales, bivalves, ſepto membranaceo medio, utramque ſeriem ſeminum, hinc inde terminante. Tota planta Guſtui calida & acris: quemadmodùm reliquæ erucarum ſpecies. Antequàm hæc planta cauleſcat, cauleſque & ſiliquas producat, folia ſunt adeo tanaceti foliis proxima, ut niſi guſtus, in opem vocetur, difficile erit terminatu. Sub quo genere militare debet: ſed flores & ſiliquæ bivalves, ſtatim ad Tetrapetalas,

Digitized by Google

trapetalas, ſiliquoſas, referri debere, indicant.

☞ Eruca vinealis, parvis luteis floribus: Hæc Erucæ ſpecies eſt annua, inter vineta proveniens, folia edit glabra ſinuata, binas vel ternas uncias longa: ex horum ſinu, exeunt, cauliculi per terram ſtrati, in ſummis caulium faſtigiis, gerentes floſculos parvos, luteos, quatuor petalis, conſtantes: quibus ſuccedunt ſiliquæ unciales & biunciales: ſeriem duplicem ſeminum continentes (aliarum ſiliquoſarum more) ſepto medio membranaceo eandem, dirimente. Nec tactum manuum patiuntur maturæ, aliarum Erucarum more. Nec mihi videtur ſylveſtris minor lutea folio Burſæ paſtoris, Banh. in Prodr. cum ejus folia neutiquàm ad Burſæ paſtoris, folia accedunt.

☞ Eruca Barbareæ æmula major: Hujus folia ſunt majora, caules altiores, ſiliquæ craſſiores, ſeminaque grandiora. Etenim planta eſt perennis: quibus notis, quam facillimè, ab Eruca Barbarea, dicta: diſtinguitur.

Digitized by Google

℞ Eryngium ex plani ſemine degener : Hoc Eryngium ex plani Eryngii ſemine, ſato pullulavit, & folia produxit non integra, ſerrata ut planum, ſed multifida, & variè ſecta, florem cœruleum & ſemina paritèr, neutiquam ab Eryngio, plano diſcrepantia, edidit.

℞ Eryngium pannonicum latifolium, album, differt à pannonico Cluſii ſolo, florum colore.

℞ Euphraſia lutea viſcata, ſerrata pratenſis : Planta eſt Dodrantalis in pratis & hùmentibus agris proveniens, intra ſepta Regiæ, Chambort, paſsímque alibi, cujus folia ſunt ſerrata longa, tactui viſcoſa, in ſummis alis caulium floſculos gerit, ſatis amplos (pro tenuitate plantæ) luteos, Galeatos, qui flaccidi relinquunt, capſulas oblongas, continentes infinita ſemina minuta : Euphraſiæ vulgaris modo.

Digitized by Google

F.

℞ FRAGARIA ſterilis incana: Hæc Fragariæ ſpecies ſterilis dicitur; quia fraga non edit, ſed ſemina ſicca, ut reliquæ pentaphylli ſpecies, folia habet tripartita ut Fragaria, ſed incana aut pubeſcentia, nec alia nota, à Fragaria ſterili vulgari differt: copiosè in ſylvis adjacentibus, pictavio, reperitur. An leucas Dioſcoridis Lob. dubitatur.

℞ Fragaria pentaphylli fructu: Hæc Fragariæ ſpecies, ſub exortum, folia omninò tripartita, & Fragariæ vulgaris æmula gerit; ſed caules emittit pedales, & ſeſquipedales, erectos: ut pentaphyllum majus; Fragariæ folio, meritò dici poſſit. Flores luteos, capſulas ſeminales, & ſemina ibidem contenta, pentaphylli ſunt æmula. Vide pentaphylloïdes hort. Regii Bleſ. aucti.

℞ Fraxinus florifera Botryoïdes: Hæc à vulgari Fraxino parùm differt; primo intuitu, quoad foliorum diſpoſitionem,

&

& corticem; ſed florum profert racemos; Dodrentales albos, pendentes: ſiliquas ſeminales nondùm vidimus.

♃ Fritillaria ramoſa, ſeu Lilium Perſicum minus: folia profert omninò, Lilii Perſici florum æmula, ſed anguſtiora, caules edit pedales, in quorum ſummitate proveniunt ramatìm ſeu Corymbatìm, floſculi quini, ſeni pendentes, pullo colore præditi, Lilii Perſici inſtàr.

♃ Frumentum Indicum, ſpica articulata, radice perenni Spicam longam ſemipedalem per articulos, ſemiunciales aut unciales, diſpoſitam gerit, qua ſola nota (juncta duratione) facile cognoſcitur.

♃ Fumaria tenuifolia, ſemine compreſſo, lini inſtár: Hæc Fumariæ ſpecies, folia omnium tenuiſſima gerit, caules promit palmares & ſeſquipalmares, in quorum ſummis proveniunt (aliarum more) floſculi Corniculati, ex albo elegantèr purpurei, quibus decidentibus, ſubveniunt ſemina, compreſſa, tumidiuſcula lini inſtàr, nec capſulas vulgaris Fumariæ ritu, producit. Et ideò indeſcriptam eſſe judicavimus: à ſemine primo intuitu, cuilibet

cuilibet obvia: creſcit in Galloprovincia, & Gallia Narbonenſi.

G.

GALLIUM album tripetalon: Hæc Gallii ſpecies folia gerit, oblonga & anguſta, vulgaris Gallii haud abſimilia, flores fert multò minores. Tribus tantùm petalis donatos. Qua nota, ab aliis ſuæ claſſis (quippe quatuor petalis donantur) facilè diſtinguitur.

Geranium Batrachoïdes, minus, perenne verſicolor: Hoc folia habet minora, minúſque incana, Ranunculi Batrachoïdis inſtàr, maculis nigris ſ. fuſcis notata; flores fert in ſummis ramis, ſtriatos, ex albo & nigro; quemadmodùm Romanum verſicolor Park. floribus cadentibus ſuccedunt ſemina quina minora, ſubrufa, roſtrum brevius aculeatum terminantia.

Geranium Batrachoïdes minus annuum purpurocœruleum: Hujus folia omninò

ninò Ranunculi Batrachoïdis; ſunt æmula, flores fert magnos, purpurocœruleos, quibus deciduis ſuccedunt ſemina paritèr quina, nigra; roſtrum Gruinum biunciale, terminantia, ad baſin. Hoc ſolum Batrachoïdes, annuum eſt, & ideò inter cœtera notu, facilius.

♃ Geranium Malachoïdes, ſeu Malvæ folio rotundo, flore albo: Hoc ſolo florum, colore albo, differt à pede Columbino Dod. ſeu Geranio Malvæ folio rotundo vulgari B. P.

♃ Geranium Robertianum, flore albo: Solo itidèm florum colore, à vulgari rubro differt.

♃ Glaux major, paluſtris, flore herbaceo: Hæc in fluviis & ad margines fluviorum: per prata labentium, provenit; caules habet pedales, aquis innatantes, à quorum geniculis nodiſve, demittuntur fibræ, in aquam, quarum beneficio nutritur tota planta; ad ſingula genicula oriuntur bina folia, pulegii Regalis foliis latiora, glabra, mucronata; in caulium alis ſuperioribus, emergunt flores, herbacei coloris qui relinquunt in capſulis oblongis

oblongis, Caryophylli aromatici magnitudine ; in ſingulis infinita ſemina minuta, ſubviridia : etiam, Euphraſiæ ad modum, continentur.

ℳ Gramen Phalaroïdes perenne, tuberoſa radice: Spicam fert, Phalaridis ſpica, tenuiorem longiorem ; ſeminaque multò ſunt minora. Radices habet tuberoſas, lucentes, perlarum inſtàr, per nodos, Typhini graminis radici ſimiles : perennis eſt planta.

ℳ Groſſularia .ſ. Ribes fructu rubro, Uvæ criſpæ folio : Frutex hic ad tripedalem aſſurgit altitudinem cortice refert Ribem, foliis Uvam criſpam, fructum fert Ribis, rubrum, racematìm, ſeu Botryos in modum diſpoſitum.

Digitized by Google

H.

ꝟ HEDERA trifolia Canadenſis; folio ſinuato: Hæc à Hedera trifolia Canadenſi Cornuti differt in hoc, quod folia, habet minora, tripartita, in margine autem ſinuata, quod non conſpicere licet, in Hedera trifolia Canadenſi Cornuti.

ꝟ Helenium Indicum tuberoſum, radice oblonga: Sola radicum longitudine; (quæ ad ſemipedem aliquandò accedit) ab Helenio (ſeu potius Chryſanthemo) tuberoſo, radice rotunda, differt.

ꝟ Helleborine, herbaceo flore: Ex primo intuitu florum, cognoſcitur. Hæc Helleborines ſpecies; in collibus, ſylvoſis. Ubi ſylvæ ſunt cœduæ, cernere ipſam frequentèr licet.

ꝟ Helleborus niger, flore roſeo, minor Belgicus: Hic, omnibus ſuis partibus, foliis, caulibus, flore, eſt vulgari hortenſi, ſeu legitimo, minor: florem fert ex albo roſeum.

ꝟ

ꝛ Hieracium flore parvo luteo, apuli, flore ſuaverubenti, odore æmulum: Hæc Hieracii ſpecies adeò odore, Hieracio apulo, flore ſuaverubenti Col. accedit. Quippe Amygdalas amaras redolet utrumque: ut idem odoratui appareret, ſed hoc folia habet anguſta, flores minimos luteos, quibus evanidis ſuccedunt, in Calycis, margine exteriori, incurva quaſi roſtra. In viarum marginibus paſsim reperitur circa Bleſas. Hæc duplici nota, odore & roſtris incurvis, in marginibus Calycis, exterioribus: cognitu cuivis obvium.

ꝛ Hieracium, foliis & floribus dentis Leonis bulboſi, ſemine curvo: A ſemine curvo facilè diſtinguitur, hoc Hieracium ab omnibus aliis, ſuæ claſſis.

ꝛ Hieracium anguſtifolium glabrum, ſeu piloſella major glabra: Caules edit Bipedales, ramoſos: folia anguſta glabra. Atque ſola glabritie, quam facillimum erit, à cœteris hujus ordinis, (quippe villis ſcatentibus) ipſum diſtinguere.

ꝛ Hieracium (pulmonaria non maculata, dictum)

dictum) indiſſectis foliis: In ſylvis montoſis, & collibus ſylvoſis: Hæc pulmonariæ ſpecies, provenit. Foliis nullis maculis conſperſis, in margine integris. Hinc à cœteris, notu, non eſt difficile.

♃ Horminum majus, amplo flore violaceo, Lobis florum, punctis aureis notatis: Hoc Horminum folia profert hirſutiora, cœteris ſuæ ſortis; (excepto ſolo Æthiopico) ad baſin anguloſa: inter quæ exeunt, caules Bipedales, aliquandò tripedales, quadrati, quos (ad genicula) amplectuntur, bina folia Dodrantalia (ut ſuperius dictum) auriculata. In ſummis caulium, verticillatim proveniunt flores ampli, violacei, Lobis florum, lateralibus; punctis aureis aut purpureis, notatis; floribus ſingulis evanidis, ſuccedunt in capſulis magnis, ſemina craſſa nigra quaterna utplurimum, majora, cœterorum omnium Horminorum, ſeminibus. Atque punctis aureis aut purpureis & ſemine majore: notu facile, à ſequenti, cum quo majorem habet affinitatem.

♃ Horminum lapathi unctuoſi folio, ſeu, majus

majus haſtato folio: Hoc folia habet hirſuta, anguloſa aut auriculata ad baſin, ut ſuperius. In mucronem deſinunt, haſtæ inſtar cuſpidis, quin & forma lapathi unctuoſi, folia repreſentat. Flores fert magnos, omninò violaceos, ſemina, in omnibus prioris ſeminibus minora, & ſubfuſci coloris.

℥ Horminum napi folio: Ex diviſione foliorum, napi foliis ſimilium & florum parvitate, lavendulæ, inſtar. Satis notu facile.

℥ Horminum Salviæ folio: Hoc Horminum adeò Salviæ majoris foliis habet ſimilia ut principio, à ſatione. Salviæ (primo intuitu) videretur ſpecies: Flores fert purpureos majores. Odore, Horminum ſeſe declarat. Fortè hæc Hormini ſpecies eſt quæ, à proſp. alpino; marum Ægyptiacum denominatur.

℥ Horminum minus, folio ſinuato, flore minore dilutè cœruleo: Caules quadratos, pedales, hirſutos, producit; procumbentes, rariùs ramoſos: foliis nonnihil hirſutis (pollicarem latitudinem vix excedentibus) oneratos; uncias

tres longis, in margine ſinuatis. Cauliculi, floribus rarius diſpoſitis ornantur, dilutè violaceis, Lavendulæ florum æmulis. Quibus decidentibus, ſuccedunt in ſingulis capſulis, quaterna ſemina minora, nigra aut fuſca rotunda. Sylveſtris quarti, quinta ſpecies Cluſ. paſsim hìc in Anglia ſecus fluvium Thamiſin, copiosè provenit, folia habet majora ſinuoſa, ſaturatiùs viridia. Caules fert quadratos (ut verticillatæ Galeatæ fere omnes) pedales & ſeſquipedales erectos. Quorum ſummis ramis, innaſcuntur flores paulò majores magiſque ſaturè violacei, quàm prioris deſcripti, ſunt flores; ſed florum magnitudine, quàm proximè accedunt.

♃ Hyſſopus hirſuta: Ab Hyſſopo latifolia vulgari, ſola hirſutie foliorum: Quemadmodùm, ſtriata ſeu aurea, à variegatione; diſtinguitur à vulgari; hoc accidit in multis aliis tàm arboribus, fruticibus, & plantis, ut ſuperiùs diximus

♃ Galeata & verticillata, Perſicæ foliis, digitalis æmula: Hæc Galeata & verticillata

cillata planta, adeò ſimilia producit folia, foliis, Perſicæ mali, quoad formam & ſerraturam, ut nihil magis, flores autem fert Galeatos, ſubpurpureos, amplos, quorum ſingulis ſuccedunt, quaterna ſemina angulosa, in capſulis quinque partitis in margine, verticillatarum & Galeatarum propriè ſic dictarum more. Nec Thecam bicapſularem, pyramidalemve: Hinc indè, infinita ſemina minuta, ſubruſa, continentem habet. Ergo perperâm Botanicis hucuſque, paſsim pro digitali Perſicæ foliis, habetur. Vide digitalis, foliis maximè ſerratis: in horto Regio Bleſenſi aucto.

I.

JACEA lusitanica sempervirens: Planta est tripedalis, folia habet, cœteris, suæ sortis, latiora, durabiliora, per hyemem totam utplurimùm durant; flores, & semina habent majora. Quibus notis, à reliquis, speciebus facilè distinguitur.

Jacobæa foliis amplioribus incanis: Ad sylvarum & vinetorum margines provenit hæc Jacobæa: foliis Jacobææ vulgaris, foliis multò; sunt latiora, minusque secta, quin & ipsis sunt multò magis incana. At Jacobææ marinæ foliis, multò minus sunt incana: medium inter utramque (quoad canitiem) obtinet locum; flores & semina paria.

Jacobæa Crithmi litorei foliis: Planta est Bipedalis, folia angusta, crassa, oblonga, viridia, Crythmi Chrysanthemi maritimi instar, habet: florem fert luteum Jacobææ, stellatum: cujus petala lata lutea, cingunt discum luteum, ex plurimis staminulis luteis, conflatum: (aliarum

(aliarum Jacobæarum more) flos abit in pappos.

☞ Jacobæa Senecionis folio : Obſerva, Jacobææ notam genericam & eſſentialem (quâ diſtinguitur à Senecione) eſſe, quod flos Jacobææ, conſtet multis petalis latis & planis : cingentibus diſcum, ex plurimis ſtaminulis luteis, conflatum. Senecionis autem flos tantùm componitur, ex ſtaminulis plurimis luteis ; nullis petalis planis, latis, eaſdem cingentibus. qua ratione hæc concluditur eſſe vera & genuina, Jacobææ ſpecies ; cum Jacobææ habeat florem ſtellatum : ut ſupra ſatis dictum. Quoniam folia omninò diſſecta habet, ut Senecio : Jacobæa Senecionis folio, aptè dicitur. Deſumpta denominatione, à nobiliore parte, Flore videlicet.

☞ Iberis Virginiana, latifolia ramoſa ; Folia fert Iberidis vulgaris foliis, latiora. Caulesque edit altiores & fruticoſiores ; annua eſt etiam planta, & hac nota à vulgari facilè diſtinguitur.

Digitized by Google

L.

℥ LAMIUM parietariæ facie : in omnibus Lamio vulgari accedit. Facilè autem dignoſcitur, ex primo intuitu, à foliorum cum parietariæ foliis, ſimilitudine.

℥ Laſerpitium alterum, foliis Lobatis anguſtis, dilutè virentibus : A Laſerpitio Gallico diſtinguitur, quod folia producit dilutius virentia, quam Gallicum, cujus folia, ſunt lucide atrovirentia.

℥ Lathyrus latifolius, luteus, annuus, ſiliqua hirſuta ; Hic Lathyrus inter Sata Bleſis adjacentia, frequentèr ſatis, provenit ; florem fert dilutè luteum (aliorum more) papileaceum cui, ſuccedit ſiliqua longa, hirſuta, qua nota præcipuè diſtinguitur ; à ſuæ claſſis, reliquis.

℥ Lathyrus latifolius, flore obſoleto, ſiliqua Piſi minoris : In arvis ſatis, circa Rupellam, frequens nobis obvius fuit, hic Lathyrus : florem ſubrubrum vel obſoleti coloris gerit ; cui ſuccedit ſiliqua

qua ſeſquiuncialis, duplici ala, cavâ parte (qua ſemina quaterna quina junguntur) prædita.

Lathyrus anguſtifolius λεπτολόϲος, ſemine rotundo, flore rubello: Hæc Lathyri ſpecies, folia ut ſequens, gerit anguſta, Graminea, longa, caulem pedalem & ſeſquipedalem, habet: in cujus cimis naſcuntur flores Rubelli, obſoleti coloris. Quorum ſingulis, ſuccedit ſiliqua rotunda, multò ſequentis, ſiliqua longior: in qua continentur ſemina dena, duodena rotunda.

Lathyrus anguſtifolius λεπτολόϲος, ſemine hexedro, flore cœruleo: Lathyrus hic folia habet anguſtiſſima, Graminea, florem fert cœruleum, papileaceum: (cœterorum more) cui ſuccedit ſiliqua, longa, tenuis: continens minuta ſemina, anguloſa cubum .ſ. hexedram, referentia.

Laurus Alexandrina, fructu longis pediculis, caulibus, alligato: Sola hac nota diſtinguitur; quod fructuum ſeu baccarum pediculi, unciales aut biunciales, à cauliculis pendent, in reliquis

Digitized by Google

autem, folio aut pediculo, baccæ, frutuſque inſident.

ℭ Liliaſphodelus, latifolius flore luteo, ſola magnitudine foliorum & florum ab anguſtifolio differt.

ℭ Lilium .ſ. Martagon Canadenſe: Elegantia florum, elegantèr à natura punctatorum: facilè diſtinguitur; à cœteris ſuæ familias.

ℭ Linaria minima hirſuta, flore obſoleto: Hæc tenuis Linariæ ſpecies, folia habet quodammodò Antirrhini æmula, florem habet ſubnigrum (inſoleti coloris) ſed omninò Linariæ, quippe rictum habet anteriùs, & calcari donatur poſterius capſulamque binis loculamentis, diſtinctam habet. Septo membranaceo utrumque diſterminante: malè ab authoribus pro Antirrhino minimo, appinguitur: in ſatis arvis, paſsim occurrit.

ℭ Linaria minor viſcoſa, flore luteo: Hæc Linariæ ſpecies, folia edit hirſuta, viſcoſa, caules ſemipedales, in quorum cimis, flores fert Linariæ, luteos: quibus evaneſcentibus, ſuccedunt capſulæ, aliarum more. Continentes ſemina ni-

gra

gra compreſſa, ſeu foliacea, non autem ut Linaria vulgaris, rotunda : (hinc nota duo reperiri genera Linariæ, quorum unum fert ſemen compreſſum, nigrum, foliaceumve, alterum rotundum, ut prolixiùs alibi, in Tabulis noſtris, cognationis, deducemus) in Arenoſis maritimis Armoricæ provenit hæc Linaria.

♃ Linaria Armorica, flore ſaturè violaceo : Hæc folia producit glabra, anguſta : caules pedales, quibus innaſcuntur flores, ſaturatè violacei, parvi, parvas relinguentes capſulas, Linariæ in modum.

♃ Linaria anguſtifolia glabra, flore luteo majore : Hujus folia ſunt quaſi Pſyllii foliorum æmula, longiora. Caulem producit, ſeſquipedalem, cujus cimis innaſcuntur, flores lutei multò majores, pro tenuitate plantæ, quâm in ulla alia Linaria. Circa Burdegalam, in ſatis, provenit.

♃ Linum ſylveſtre annuum, luteum, folio latiore & anguſtiore : Spontè in humidis ſylvis, utraque hæc ſpecies creſcit, ſolo hoc diſcrimine quod, unius folia

ſunt

ſunt latiora, alterius anguſtiora. *Vix* Dodrentalem altitudinem excedit, utraque ſpecies.

ꝺ Linum perenne cœruleum, ſativo æmulum: Hoc in omnibus ſativo accedit niſi quod eſt perenne.

ꝺ Lychnis flore muſcoſo capillaceo: Habet caules ſeſquipedales, Lychnidis (ſeſamoïdes Cluſio) dictæ, æmula. Sola hac nota ab omnibus Lychnidum ſpeciebus differt quod non (aliarum more) florem, quinis petalis, ad medium bifidis, gerat. Sed florem edit ex plurimis capillis muſcoſis, herbacei coloris.

ꝺ Lychnis Alpina glabra, Caryophylli holoſtei flore: A foliis glabris Caryophyllorum, foliorum æmulis, biuncialibus aut triuncialibus, notu perquàm facilis, flores fert minores albos, quinis petalis, ad medium bifidis conſtantes; capſulam ſeminalem, & ſemina aliarum more producit.

ꝺ Lychnis minor viſcoſa Saxatilis; Caryophylleis foliis: Hæc inter Saxoſa, provenit, Pyrenæorum montium & in Alpium jugis, folia habet glabra, mucronata,

cronata, Caryophylli inſtar : Caules vix ſemipedales profert : ad ſingula internodia (ut in pluribus aliis Lychnidum ſpeciebus) viſco obductos ; quo muſcæ captantur, unde viſcoſa dicitur.

℔ Lychnis maritima repens, ſtrictiſſimis foliis : Hujus cauliculi per terram ſunt ſtrati, quibus ſtrictiſſima, innaſcuntur folia, flores fert albos minores, quibus pereuntibus, ſuccedunt capſulæ, aliarum Lychnidum more, & ſemina ibidèm contenta. Provenit in ſabuletis maritimis armoricæ, & circa Rupellam. Hæc planta.

℔ Lychnis incana longiori folio, petalis non bifidis : Hæc Lychnidis ſpecies, quod folia edit longiora, incana, & florum petala, habet non bifida, notu facilima, nec novi ullam, hujus generis ſpeciem aliam, quæ petala gerit non biſida : præter auriculam muris, pulchro flore, folio tenuiſſimo. Joh. Bauh. in Ericetis ſabuloſis, ſex milliaria, à Burbonum capite, diſtantibus, reperitur, ſatìs frequentèr.

℔ Lychnis (papaver ſpumeum) latifolia vulgaris,

Digitized by Google

vulgaris, hirſuta: Hæc ſola foliorum hirſutie, à papavere ſpumeo, vulgari glabro, differt; inter ſata, valdè frequentèr occurrit.

♃ Lychnis (papaver ſpumeum) anguſtifolia glabra, flore purpureo, ab anguſtifolia vulgari glabra, floris colore, ſolummodò differt.

♃ Lycopſis Monſpeliaca, floribus dilutè purpureis & albis: Ad tripedalem aſſurgit altitudinem, flores Corniculatos dilutè rubros & albos, aliorum Echiorum aut Lycopſium more habet, quaterna, dicidentes relinquunt ſemina; viperini capitis, figuram referentia. Nota eſſentialis & generica quâ diſtinguitur Echium ſeu Lycopſis. A Bugloſſa ſeu Anchuſa, eſt hæc. Quòd hujus flores quinis petalis conſtent: illius autem flores ſunt Corniculati. Quod cuilibet obſervanti & propiùs intuenti, quam facillimè apparet.

♃ Lyſimacha Corniculata, lutea Canadenſis minor ſeu anguſtifolia: Hæc ſola foliorum anguſtia, aliarum ſuarum partium; florum ſcilicet & capſularum ſeminalium,

minalium, parvitate differt, à Lyſimachia Corniculata lutea majori Cornuti.

℥ Lyſimachia purpurea ſpicata lanuginoſa, ſeu folio ſubrotundo incano: A Lyſimachia ſpicata communi purpurea Cluſ. differt, quod hujus folia ſunt glabra, oblonga, mucronata; illius vero incana ſeu lanuginoſa, ſubrotunda, cœtera ei par.

M.

℥ MALVA Betonicæ folio: Cauliculos profert pedales & ſeſquipedales per terram ſtratos quibus adhærent alternatim, folia oblonga, Betonicæ foliorum inſtár. Flores fert amplos Rubellos; ſemina, nec ad maturitatem pervenêre.

℥ Malva Hiſpanica: Hæc Malvæ ſpecies folia producit mollia, undulata in margine. Superiùs, micis ſulphureis ad ſolem, lucentibus donata ſunt. Caules edit bipedales, aut tripedales, flores & ſemina

ſemina aliarum Malvarum modo. Perennis eſt planta: quin & perpetua fronde viret. Ut Althæa Olbiæ, in meridionalibus ſaltem regionibus.

ℳ Marrubium aquaticum ſeu aquatile foliis tenuiùs diſſectis: Hæc Marrubii ſpecies, longè lateque radices ſuas, hâc, illàc ſerpendo, ſpargit: Caules fert tripedales ut vulgare. Folia autem duplò aut triplò tenuiùs, vulgaris foliis, ſunt diſſecta. In omnibus aliis, vulgari congruit.

ℳ Meliſſa Romana hirſutior: Hæc folia profert multò vulgaris foliis latiora, multóque hirſutiora, quin & tota planta, multò eſt major vulgari, ad ternos, quaternos (aliquandò) aſſurgit pedes. Hujus ſemen, ut & aliarum plantarum (in horto Regio Bleſenſi aucto, contentarum) multa, ad me miſſa fuerunt ſemina. A doctiſſimo & peritiſſimo Botanico, Domino Jacobo Barillier, in ſacra ſancta Theologia Doctore, ordinis prædicatorum ſocio, digniſſimo, ad minervam Romæ, pro eo tempore commorante: cui immortales debemus gratias.

ℳ

Melissa Bætica: Vide Brunella Bætica suprâ: Cum non habet odorem Melissæ, nec flores Galeatos, rariùs per verticillos dispositos ut Melissa. Sed verticillos densissimè dispositos: & quasi (prunellæ in modum) in spicam degenerantes, prout clariùs deducemus in Tabulis nostris, cognationis. Pro prunella habeatur.

Millefolium tanaceti foliis: Hujus folia sub exortum, antequam, caulescat planta, adeò sunt Tanaceti, foliorum æmula, ut non nisi ab oculato Botanico, distingui possint; nisi gustus & olfactus, vocentur in opem. Sed postquàm produxerit caulem & Corymbum, Millefolii in modum, semenque paritèr Millefolii. Sese declarat verum & genuinum Millefolium. Estque flore albo & purpureo. Vide Tanacetum montanum purpureum: in horto Regio Bles. non aucto. Quàm ineptè Botanicis, Tanacetum dicitur, videat lector & examinet mecum.

Myrrhis perennis lutea daucoïdes: Hujus folia quàm proximè dauco, seu pastinacæ

paſtinacæ tenuifoliæ, accedunt, quin & flores habet luteos, ſemina roſtrata ſtriata, Myrrhidis. Montanæ, ſeminibus longiora at minùs tumida habet. Perfecto ſemine, ſeſe renovat ex eadem radice.

♃ Myrrhis nodoſa annua, ſemine aſpero: Hujus caules ad internodia craſceſcunt, nodoſque gerunt pyriformes, flores gerit albos, in ſummis caulium bipedalium. Quibus exeuntibus, ſuccedunt ſemina roſtrata bina, ſtriata, aſpera, in convexa parte; imo multò ſunt, reliquis craſſiora. Unde à nodoſitate caulium flore albo, & ſemine ſtriato aſpero, quàm facillimè, à reliquis Myrrhidum ſpeciebus diſtinguitur. Hoc addito, quod ſit annua planta. Nota quod omnis Myrrhidis, ſemina bina ſint ſtriata; Cereſolii autem lævia nigra, ergò Myrrhis Fuchſ. ſeu ſylveſtris ſeminibus lævibus B. Pin. eſt Chærophyllum ſylveſtre. At Anrhriſcus Plinii hiſtoriæ Lugd. eſt Myrrhis ſylveſtris, non Chærophyllum ſylveſtre. Quia ſemina habet ſtriata, hoc adde quod nodos gerit pyriformes.

♃

♃ Myrrhis perennis montana alba: Hæc Myrrhidis ſpecies aſſurgit, ad binos ternoſve pedes, caules producit ſtriatos, flores (aliarum ſuæ claſſis more) albos: quibus ſuccedunt ſemina roſtrata, ſtriata. Semiuncialia exteriùs lutea: cæva parte qua junguntur, ſubcœrulea; eſt foliis hirſutis & hirſutioribus, cujus folia medio modo, ſecta ſunt, ſatis lata. Eſt & tertia hujus ſpecies, cujus folia ſunt tenuiùs diſſecta.

♃ Myrtus aurantiæ folio: Hujus folia ſunt aurantiæ foliorum, colore & conſiſtentia, adeò æmula, niſi quod ſint multò minora, & anguſtiora. Ut malum aurantiam minorem (antequàm flores & fructum edit) nullus dicere vereretur. An Myrtus latifolia Park. Cluſ. hiſt. dubitatur.

T N.

Digitized by Google

N.

NASTURTIUM pratenſe floſculis minimis: Pedalem & ſeſquipedalem habet altitudinem hæc planta: folia quàm tenuitèr diſſecta, floſculos minimos albidos quatuor petalis (aliarum ſiliquoſarum, tetrapetalarum more) conſtantes producit; quorum ſingulis ſuccedunt ſiliquæ biunciales, bivalves, ſepto medio membranaceo, diſterminante: duplicem ſeriem, ſeminum, cœterorum (ſuæ claſſis) more. Quin & matura ſemina, manuum tactu; per terram ſparguntur. Unde noli me tangere dicitur. Fortè Naſturtium ſylveſt. minus Bauh. in Prodr.

Nigella ſylveſtris, flore luteo pallido; Seu Lychnis (Pſeudomelanthium) dicta ſolo, florum colore, à vulgari Pſeudomelanthio, differt.

O.

Digitized by Google

O.

ꝛ OCYMUM minimum foliis rubeſcentibus: Ex ſatione, tanta eſt varietas horum, ut folia aliquandò rubeſcant, aliquandò ſunt luteopallida, ut plurimùm viridia. Aliquandò integra, nonnunquam ſerrata, ita ut ex ſatione, concludo eſſe luſum naturæ, ut in multis aliis annuis.

ꝛ Onobrychis, flore ſaturatè rubro: Ab Onobrychide vulgari, ſolo florum colore differt.

ꝛ Onobrychis, viola Pentagonia dicta, vide Campanula .ſ. viola Pentagonia.

P.

ꝛ PASTINACA tenuifolia lucida: Vulgaris Paſtinacæ, folia ſunt hirta, hujus folia ſunt glabra & lucida, ſola luciditate & glabritie foliorum, à vulgari facilè diſtinguitur.

Digitized by Google

☞ Paſtinaca tenuifolia, radice & umbella lutea: Facilè ab hortenſi & ſylveſtri, diſtinguitur; quippe harum umbellæ, colore ſunt albo illius umbellæ ſunt colore luteo, præditæ.

☞ Pentaphylloïdes Fragariæ folio: Vide Fragaria Pentaphylli fructu.

☞ Pentaphylloïdes, Argentinæ facie, nobis: Pentaphyllum ſupinum potentillæ facie Cluſ.

☞ Pentaphylloïdes, Ulmariæ facie, nobis: Pentaphyllum fragiferum 5. Cluſ. cum duæ poſteriores plantæ, coſtas ſeu nervos medios, longos habent, quibus quina, interdùm plura foliola, adhærent è contrario, inter ſe foliorum exortu, naſcentia: uno extremum claudente. Potentillæ .ſ. Argentinæ, ritu. Ergo, cum florem & ſemen, aliorum Pentaphyllorum more, perficiunt. Folia autem Pentaphylli non habent, Pentaphylloïdes nuncupari velim.

☞ Petaſites minor, folio tuſſilaginis: Petaſitis folia ſunt ſcabra aſpera. Hujus autem, ſunt prona parte viridia, ſupina, incana, tuſſilaginis inſtàr, & niſi quod florem

florem producit Petaſitis, eſſet vera tuſſilaginis ſpecies.

☞ Peucedanum minus foliis Lobatis anguſtis: Hæc umbella, folia producit quina, ſena, nunc plura, nunc pauciora, è regione Rachi ſeu nervo medio, adnaſcentia, uno extremùm claudente, è quorum medio ſeſe erigit Caulis Bipedalis, ſuſtinens flores luteos, umbellæ in modum; ſemina fert membranacea, rotunda, Peucedani vulgaris inſtár.

☞ Phalangium Virginianum, amplo flore: A Phalangio Virginiano tradeſcanthi, ſola amplitudine, florum & ſeminum differt. Eſtque flore rubro, cœruleo & azureo.

☞ Phalaris, ſemine minore nigro: Hæc ſemen, producit, minus Phalaridis vulgaris, ſemine, quod & nigrum eſt, qua nota, facilè, diſtinguitur à Phalaride Dod.

☞ Phaſeolus minor Colloſſinus, ſiliquis longiſſimis: Hic Phaſeolus ut & ſequens, vix pedalem aut ſeſquipedalem excedunt altitudinem; Hic florem Coloſſinum: fructumque minorem Coloſſinum,

ſiliquas, ſenas, ſeptenas uncias longas fert, ſtrictiſſimas. Non ſcandit nec ſequens.

ꝟ Phaſeolus, ſiliquis rotundis Viciæ, fructu ovali viridi: Hæc Phaſeoli ſpecies, vix pedalem excedit altitudinem, flores fert herbacei coloris; aut ſubvirides, papiliaceos, ſiliquas Viciæ in modum, rotundas, quaternas uncias aliquandò excedentes: ſemina oblonga, viridia, ovalia; in ſe continentes.

ꝟ Piloſella major glabra: Hæc à majori repenti hirſuta Bauh. in Pin. differt ſola glabritie.

ꝟ Pimpinella maxima, ſpica longa rubra: Hæc à ſanguiſorba differt, quod ſpicam gerit, ipſa longiorem, ternis, quaternis unciis conſtantem, hujus eſt & alia ſpecies, ei omninò accedens, niſi quod ſpica ſit brevi, Glomerata, rubra paritèr, utraque, folia gerit longiora, & magis ſerrata, quàm Pimpinella ſanguiſorba.

ꝟ Pimpinella Agrimonoïdes: Hujus folia (prima facie) Agrimoniæ foliis quàm proximè accedunt: aſpera, villoſa ſunt,

Rachi

Rachi ſeu nervo medio (aliarum Pimpinellarum more) adhærent, uno extremum claudente gratiſſimum, contrectata ſpirant odorem, caules pedales habet: in horum cimis proveniunt, ſpicæ Glomeratæ, ſeminibus ſtriatis (gratiſſimi odoris) onuſtæ.

♃ Piſa nana rotunda: Omnium, ſuæ familias ſunt minima, & humillima. Piſa hæc, nec prolixiore egent deſcriptione.

♃ Piſum ſylveſtre Viciæ fructu: Primo intuitu, à folliculo ſeu ſiliqua Viciæ dignoſcitur.

♃ Piſtachia Virginiana, ſylveſtris triſolia: Vel primo intuitu à foliorum (ternatim) diſpoſitione, ſeminumque in veſicis parvitate, notu, à vulgari, perquàm facilis.

♃ Plantago marina quinquenervia incana: Hujus folia Plantaginis quinquenerviæ vulgaris, in omnibus, accedit: niſi quod ſint minora & villis adeò obducuntur, ut incana, aut cinere obducta videantur.

♃ Polium rubrum majus, flore dilutiore: Hujus folia, Polii lutei vulgaris,

foliis, quàm proximè accedunt, & quoad magnitudinem, ſerraturam & incaniti-em. Florem fert majorem, dilutiùs rubrum, quàm quintum Cluſ.

ꝛ Polyganum minus maritimum incanum: Hujus folia ſunt Polygoni marini vulgaris, multò minora, cinerei coloris, aut incana lanugine, obducta videntur.

ꝛ Portulaca minima flore albo: Hæc plantula eſt uncialis, in humidis arvis, ineunte æſtate, per terram ſtrata: folia producit ſemiuncialia anguſta, quodammodò craſſa. Floſculos ad nodos producit albos. Quibus deciduis, ſuccedunt, minutula ſemina, Portulacæ, ſeminibus omnimodè conformia, nigra itidem.

ꝛ Portulaca hortenſis flore flavo: Solo foliorum colore, à Portulaca hortenſi, ſeu culta differt.

ꝛ Potamogeton Virgineanum: Bipedalis aut tripedalis eſt planta: folia gerit latiora, obſcurè virentia; ſpicas triuncíales, albis floribus rarò ibidèm diſpoſitis, onuſtas fert, non autem compactìm ſpicæ in modum. Semina habet triquetra, ut Perſicaria & Polygonum.

ꝛ

℔ Ptarmica Virginiana folio Helenii: Hæc eſt Corymboſa, Bipedalis planta, cujus caulibus, adnaſcuntur, folia lata & oblonga, Helenii foliis haud abſimilia, flores fert (corymbatim) albidos, ſeminaque, Ptarmicæ vulgaris, ſeminibus, omninò ſimilia unde ei nomen impoſuimus.

Q.

℔ QUERCUS vulgaris brevi pediculo: Hæc facilè diſcernitur. In ſylvis annoſis, Galliæ paſsim; à vulgari, longo pediculo, vel ipſo aſpectu. Folia habet ſinuata; prona & ſupina parte, viridia.

℔ Quercus foliis ſubtus ineanis, brevi pediculo. Datur & alia longo pediculo. Utraque folia habet anguſta, ſinuata, ſubtûs incana.

℔ Quercus Calyce Echinato, glande majore: Hæc à glandis magnitudine, & Calyce Echinis donato, inter cœteras ſuæ familias (in ſylvis annoſis, Galliæ) cognitu perfacilis.

℔

♃ Quinquefolium alpinum repens, ſerotinum: Hæc Quinquefolii ſpecies, ſub exitum autumni floret, caules per terram ſtratos gerit, ſuſtinentes flores luteos.

♃ Quinquefolium Canadenſe erectum: Hujus caules ſunt Bipedales, erecti, floſculos gerunt magnos, luteos, capſulas ſeminales, aliorum more.

R.

♃ Ranunculus arvenſis annuus flore minimo luteo: Hic Ranunculus folia gerit hirſuta, tripartita, caules edit per terram ſparſos, quorum ſummis caulibus innaſcuntur, floſculi lutei, minimi, quinis (aliorum more) petalis donati. Quibus deciduis, ſuccedunt racematim, ſemina minora, ſpinoſa aliorum more.

♃ Ranunculus nemoroſus flore albo pleno: Hic ſola plenitudine florum diffèrt, à nemoroſo, flore albo ſimplici.

♃

♃ Ranunculus nemorosus flore pleno violaceo: Hic pariter plenitudine florum, & colore violaceo, differt, à nemoroſo ſimplici. Quoad folia & radices idem.

♃ Ranunculus Ceratophyllos, ſeminibus falcatis, in ſpicam addactis: Foliis eſt ter, tripartitis: Cornu cervi quodammodò referentibus è quorum medio exeunt cauliculi plurimi, trifuriculeles, linguli, ſingulos ſuſtinentes floſculos calyculum Ranunculorum more, quinis lacteis petalis conſtantes; ambientibus rudimentum viride, quod tandem in ſpicam degenerat uncilem; ſeminibus falcatis onuſtam. Annua eſt plantula, præcox. Menſe Martio floret, in ſegetibus, Gallo-Provincia, & Narbonenſi Galliæ, ſatis frequenter: perperam à Lobelio in adverſariis pagina 11. appingitur pro Melampyro perpuſillo; cum Melampyrum (in genere) ſit planta Galeata, toto Cœlo, ab hac Ranunculo, diſcrepans. Cumque omnino propius ad Ranunculorum claſſem accedit. Magis apposite Ranunculus dicitur. Vide Hallucinationes

lucinationes Caſpari Bauh. in Pin. pag. 234. colum. 2. plant. 6.

℥ Raphanus ſylveſtris major flore luteo: Folia ſunt longa, ſerrata: aquis innatant. Flore eſt luteo, ſed dum recedunt aquæ, mutant inciſuram foliorum; florem fert luteum; (ut ante dictum) ſiliquas ſeu potius ſiliculas fert, aliorum (ſuæ claſſis) more.

℥ Raphanus minimus luteus Naſturtii folio: Hæc planta florem fert Tetrapetalum, luteum; Thecamque ſeu ſiliculam, bicapſularem: radicem etiam edit longè latéque repentem, ut Raphanus vulgaris, quin & uda, humidaque amat loca. Quæ omnia Raphani ſeſe declarant ſpeciem: folia autem gerit Naſturtii tenuitèr diviſa, ſeu diſſecta, flores ſatìs amplos luteos, pro exiguitate plantæ. Vix. palmares aut ſeſquipalmares edit caules, quorum ſummis, (poſt deciduos flores) ſuccedunt ſiliculæ bicapſulares.

℥ Rapuntium Solonienſe, violaceo Galeato flore: Hæc Rapunculi ſpecies, radicem habet, aſclepiadis; caules emittit bipedales, quorum ſummis faſtigiis, eorumque

rumque alis, innaſcuntur flores Galeati, cœrulei: Rapuntii Americani, (quod flos Cardinalis Columnæ dicitur) omninò æmuli. Quibus flacceſcentibus, ſuccedunt capſulæ triangulares, (aliorum Rapunculorum more) continentes ſemina pellucida ruſa. Tota planta percalida eſt, & Maſticata linguam, urendo, pungit. Creſcit in humidis ericetis & ſylvis annoſis, paſsim in agro Solonienſi, alibique in Gallia. Malè à Caſparo Bauh. in Pin. pag. 110. collocatur pro draba, flore cœruleo Galeato. Quum nil habet affinitatis, cum ſiliquoſis, cujus draba eſt ſpecies. Rapuntium Galeatum ſeſe declarat, ex ſucco lacteo emanante ex qualibet parte rupta; adde capſulam ſeminalem, & ſemina. Rapunculorum claſſi accedentia.

♃ Reſeda latifolia, flore flavo: Hæc Reſedæ ſpecies, folia gerit lata, obſcurè virentia; caules edit Bipedales, in quorum ſummis naſcuntur flores flavi (aliarum Reſedarum more) ſpicatìm dipoſiti: quibus evanidis ſuccedunt, Thecæ, ſeminales tricapſulares, ut etiam, reliquis, ſuæ familias; contingit. ♃

Digitized by Google

☞ Reſeda foliis Calcitrapæ, flore albo: Hæc Reſeda, folia gerit omnimodè diviſa, quemadmodùm Reſeda vulgaris; caules edit ſeſquipedales, in quorum faſtigiis, naſcuntur, flores albi, ſtaminulis multis purpureis donati. Quibus ſuccedunt Thecæ tricapſulares; ſeu tribus loculamentis diſtinctæ; ſemina multa nigra in ſe continentes.

☞ Ricinus minimus: Omnibus ſuis partibus: cedit, majori & medio, quoad ſemina, folia & caulium altitudinem.

☞ Rubeola maritima lutea: Hæc Rubeolæ ſpecies, in ora maritima, Armoricæ. Radice longè latéque repente, reperire eſt, paſſim. Flore donatur luteo minimo (quatuor petalis conflato) flori unicuique, ſuccedunt bina ſemina teſticulata, Rubeolæ more.

☞ Rubeola minima alba: Plantula eſt in viis aridis, conſpectui, ſatis obvia: flore minimo albo, tetrapetalo aliarum more, foliis ſpinoſis donatur. Florum ſingulis, ſuccedunt bina ſemina teſticulata; cœterarum more.

S.

Digitized by Google

S.

☞ SANICULA montana crenata, umbilico pallido: Species eſt Saniculæ montanæ Guttatæ, cujus folia ſunt oblonga tenuia, ſerrata, è quorum medio; foliis crenatis (Roſæ inſtàr)-ſepto: exit cauliculus Dodrantalis, ruber, ſuſtinens plurimos flores albos, (aliarum more) punctatos, umbilico pallido affectos. Quorum ſingulis cadentibus ſuccedunt ſiliquæ ſingulæ bifurcatæ, aliarum Saniculârum, montanarum, & Saxifragæ Chelidonidis more, nam hæc eſt vera & genuina Saxifraga montana, à flore & capſula ſeminali, ſeminibus inibi contentis, datur plantæ jam delineatæ, ſpecies minor, quoad folia & caules.

☞ Saxifraga aurea Lichenis facie: Hæc Saxifragæ aureæ, ſpecies, ſatìs frequentèr reperitur in rivulis aquæ duleis, à fontibus jugibus manantibus. Folia gerit oblonga, ſerrata hirta per terram ſtrata. Floſculos in alis, luteos; quin & primo intuitu,

Digitized by Google

intuitu, à Saxifraga aurea montana rotundifolia, diſcernitur, à foliorum, denſa diſpoſitione, & forma longa, cum alterius, montanæ, ſint rotunda & rariora.

♃ Scabioſa Gramineis foliis glabris: Hæc adeò eſt ſimilis, Scabioſæ: argenteo, folio, quod hæc, folio glabro ſit. Altera, folio argenteo, ſeu, incano prædita. Utraque eſt Calyce membranaceo, & ſemine lanuginoſo, donata.

♃ Scabioſa foliis diſſectis, flore amplo, Calyce membranaceo, ſemine lanuginoſo: Hæc folia, admodùm tenuitèr diſſecta, Columbariæ adinſtàr habet; ſed flores gerit amplos, dilutè cœruleos, quorum ſingulis ſuccedunt, ſemina lanuginoſa, Calyce amplo membranaceo, donata.

♃ Scabioſa minima, annua, erecta, ſemine anguloſo: Vulgaris inſtàr, hæc Scabioſæ ſpecies eſt. Pedales & ſeſquipedales profert caules, erectos. Folia, vulgaris quodammodò æmula, ſed multò minora. Flores itidèm multò minores, carneos: quin & ſemina habet, vulgaris ritu, anguloſa.

♃

Scorzonera noſtras, pulverifera: Hæc in pratis humidis & udis, Italiæ Galliæ & Germaniæ, inter perfectas noſtrates provenit, in pratis inquam & ſylvis humidis; ſecus fluviorum margines, ſitis; à noſtrate perfectum florem & ſemina gerente, in hoc ſolo, differt, quod in hujus caulium ſummis, naſcitur, globulus, Bellidis majoris more, Calyculo inexpanſo: in cujus ſinu, loco floris continetur pulvis ferrugineus, ad Cynaberis colorem, valde accedens. Subtilitate nulli, peritiſſimi artificis manus induſtria, facto, ſecundus. Imò brevi, partìm pluviis ingruentibus, partìm humidis deciduis (nocte) vaporibus, accedente ſolis diurno calore, in materiam fæci vini ſimilem, exſiccatam, degenerat: Quam Tartarum vini, acetarii, nominant: qua tandem decidente, apparet apex cauliculi, ad modum dentis Leonis (caput Monachi dicti) floribus & pappo ſuo exutus. Scorzoneram ritè dici arguunt partium diſpoſitio & omnimoda cum Germanis ſeu noſtrate conformatio, excepto Apice. Pulverifera dicta,

quod loco floris & conſequentèr ſeminum, pulverem (ferruginei coloris) in Calyculo caulium, Apici innaſcenti, gerat. Viget in omnibus pratis humidis Galliæ indifferentèr cum noſtrate, ſeu Germanica dicta. Apex autem pulverifer, ab Aprili medio, uſque ad idus Maii exactas, perſeverat, nec ultra hoc tempus, ipſum conſpicere licet, cum ſit tantum vitium naturæ, à Cœlo aut ſolo proveniens. iiſdem viribus, & radix & folia ſunt prædita, quibus & noſtras perfecta, ut ex radicis, foliorum colore, ſapore, ſuccoque lacteo, totiuſque plantæ conſiſtentia, colligere licet; idem obſervare licet in Tragopogo & aliis, ut ſæpe nobis fuit obvium, in peregrinationibus noſtris Botanicis, ſereniſſimi, principis Gaſtonis Franciæ, impenſis & liberalitate factis.

℧ Scrophularia foliis & caule hirſutis: Hæc Scrophulariæ ſpecies folia habet hirſuta, lanuginoſa, Urticæ quaſi æmula, quin & caules producit multò humiliores, vulgaris caulibus; paritèr hirſutos, flores & capſulas bivalves Pyramidales, vulgaris more habet. ℧

Digitized by Google

Scutellaria Americana, Malvæ folio: Planta eſt emittens folia, quodammodò Malvæ foliorum æmula (quoad figuram) ſed majora è quibus proſilit caulis, ſeſquipedalis, rectus, plurimos in faſtigio fundens flores verticillatos Galeatos, quibus evanidis ſuccedunt capſulæ, tali inſtar, qui ſupponitur, mulierum Calceis, ſi poſteriùs inſpiciatur capſula, ſi autem ſuperiùs conſideratur capſula ſeminalis, Scutellam omninò refert, cavitate ſua, in ſuperiore parte. Nos hanc exoticam addimus novo generi à nobis deſignato. Scutellariam appellando ex variis nominibus, ab authoribus Botanicis, variis in locis diſpoſitis: ut cuilibet conſideranti, clariſſimè apparebit ex doctrina de Hallucinatione Caſpari Bauh. in Pin. tam in denominandis quàm digerendis plantis. Nos omnes plantas, capſulas ſeminales, Scutellatas, ad eundem modum efficientes Scutellatas appellavimus, & Scutellariæ, novum genus creavimus.

Securidaca peregrina: Fortè Pareſſeux Galicæ id eſt planta ſenſibilis pygra

Legumen eſt, cujus unicuique, foliorum Rachi, ſeu nervo medio; è regione, plurima adnaſcuntur folia, dena duodena, uno extremum claudente. Nec ſiliquam nec ſemen obſervavi. In horto Regio Bleſenſi floruit, ſed per rigorem tempeſtatis, anni periit ſine ſiliqua aut ſemine. Pigrè ſeu lentè ſe movet, tacta.

♃ Securidaca minor lutea, ſiliquis articulatis: Vide Ornithopodium ὀρνιθόποδον minus nobis. Scorpioïdes altera Johan. Bauh. in horto Regio Bleſenſi aucto.

♃ Sedum longifolium Citrino flore: Hæc ſedi ſpecies, à ſedo longo folio vulgari, ſolo, floris Citrini colore; differt. Quippe vulgare teritefolio; aureo ſeu luteo donatur flore, ſemen & capſula ſeminalis, hujus & illius, eadem; frequentèr reperitur in rupibus Saxoſis, ad dextram Viæ communis, eundo Bleſis, in oppidum, pictonum caput; vix milliare, ab oppido diſtantibus.

♃ Sedum teretifolium majus flore albo: A præcedenti ſolo floris colore differt. Circa Rupellam, frequens ſatìs in tectis, & ſummis murorum, reperitur.

♃

℥ Senecio Jacobææ folio: Hujus folia omninò ſunt Jacobææ, in modum ſecta, ipſis in omnibus ſimilia, at dum florem edit, Senecionem ſe prodit, quia Senecio florem fert, multis ſtaminulis luteis conflatum. Sine ſoliis latis, ſtellatìm aut rotatìm, ipſa ſtamina cingentibus, quod videre eſt in hujus flore.

℥ Sideritis glutinoſa, bitumen redolens: Hæc Sideritidis ſpecies, Dodrantalis eſt altitudinis, cujus folia, ex adverſo ſita, ſunt hirta, glutinoſa Caulibus (binatìm) adhærentia, contacta manibus, & naribus admotis: bitumen redolent, flores albos & ſemina reliquarum Sideritidum modo fert.

℥ Sinapi ſiliqua hirſuta, ſemine rubro: Hæc Sinapi ſpecies à vulgari, ſiliqua hirſuta, ſemine albo, ſolo ſeminum colore differt.

℥ Sium minimum ferulaceis foliis: Sium hoc ad margines fluviorum, per ſiccitatem æſtatis, exſiccatorum reperitur. Vix palmarem excedit altitudinem: folia gerit ferulacea ſeu Fœniculacea, craſſa biuncialia, aut triuncialia: pediculos

parùm aut nihil Palmum ſuperantes, edit. In quorum faſtigiis naſcuntur floſculi umbellati albi, quorum ſingulis deciduis ſuccedunt bina ſemina, minuta, ſtriata, aliarum hujus generis umbellarum more.

♃ Solanum foliis & caule ſpinoſis: Hujus folia ſunt lata, anguloſa, ſinuatave, ſpinis armata, quemadmodùm etiam caules, ſpinis armantur, flores fert Solani modo, aut potiùs borraginis, ſemina nunquam obſervare licuit quippe perennis eſt planta, radice per terram longè latéque repentè, dotata. Ideò natura, minùs productioni ſeminum intenta, quod in pluribus repentibus obſervavimus.

♃ Solanum fœtidum pomo grandibus aculeis donato: At nobis, pomo aculeis craſſioribus & longioribus donato: Hæc Solani ſpecies, & folia & flores vulgaris foliis & floribus, multò gerit minora, poma autem aculeis, multò craſſioribus & longioribus prædita, fert.

♃ Solanum pomiferum fructu rotundo, parvo, molli, rubro, plano: Vulgò pomum

mum amoris hoc, dicitur authoribus. Fructu eſt ceraſi inſtar (quoad magnitudinem) nunc, luteo, nunc, rubro eſt colore.

☞ Solanum pomum amoris, fructu, ſtriato molli, aureo: Hujus pomum, ſeu fructus, ſolis ſtriis, à molli aureo plano majore differt.

T.

☞ TELEPHIUM Portulacæ folio: Hæc Telephii ſpecies, ſola craſſitie, ſeu ſpiſſitudine foliorum, Portulacæ æmulorum, à cœteris ſuæ claſſis, facilè diſtinguitur.

☞ Teucrium regium purpureum: Hæc Chamædryos ſpecies, multò altior eſt vulgari, humilior Teucrio fruteſcenti vulgari: floſculos gerit amplos rubros, Galeatos. Quibus pereuntibus ſubveniunt ſemina itidèm magna: folia ſunt Lucida, obſcuréque virentia.

ꝟ Thalictrum pratenſe maximum, Monſpelienſium aſperioribus foliis ſeu : Folio rugoſo ; nobis. Hæc Thalictri ſpecies ad ternos quaternos, pedes aſcendit. Folia, rugis ſcatentia, gerit, in ſuperiori, ſeu prona ſuperficie, dilutiùs virent, quam, pratenſe vulgare. Muſcarium ſeu Muſcoſos flores edit luteos, ſemina rugoſa (aliorum more) & ſtriata, (excepto Thalictro Canadenſi, Cornuti : quod ſemina gerit triangula) omnia Thalictra habent ſemina ſtriata.

ꝟ Thaſpi Iberidis folio, tenuiſſimè diviſo : Hoc, omninò Thlaſpi Iberidis folio accedit ; in omnibus, niſi quod folia habeat, multò tenuius diſſecta. Flores & ſemina, cœteraſque partes minores.

ꝟ Thlaſpi Alyſſum minimum : Hæc Thlaſpios ſpecies, eſt à flore, ſemine omnibuſque ſuis partibus à prioribus, (majore & minore videlicet) notu facilis, propter exiguitatem partium.

ꝟ Tithymalus Caracias lanuginoſus : Hic Tithymalus paſsim in ſylvis Galliæ Solonienſis, provenit, & à foliis lanugi-

ne

ne obductis, à cœteris, ſuæ ſortis facilè cognoſcitur.

ℳ Tithymalus Caracias radice repenté: Folia gerit oblonga, radicem longè latéque ſerpentem producit qua nota diſtinctu eſt facilis.

ℳ Tithymalus Caracias radice pyriformi: Sola radicum tuberoſitate pyriformi, dignoſcitur inter omnes Tithymalos (Caracias) dictos.

ℳ Tithymalus Caracias, Salicis minoris folio: Hic à ſimilitudine, foliorum Salicis longifoliæ vimineæ, notatu facillimus.

ℳ Tithymalus Linariæ folio, lunato flore: A foliis Linariæ ſimilibus & flore, Lunæ creſcentis, referente figuram; abſque longa verborum ſerie, facilè dignoſcitur.

ℳ Thymum Creticum fruteſcens: Planta, eſt ſeu potiùs Suffrutex, ſeſquipedalis, nudos edens cauliculos, foliolis longis, Satureiæ omninò ſimilibus, onuſtos, quin & odorem fragantiſſimum, inter Thymi & Satureiæ, medium, ſuggerit.

ℳ Tragopogon pratenſe luteum minus:

Hæc

Hæc Tragopogi ſpecies, ſola partium ſuarum, præcipuè florum & ſeminum, foliorumque parvitate, à majore differt deſcribitur à Parkinſono, in Theat. Botanico.

♃ Trifolium bitumen redolens majus: Hoc à minore omnibus ſuis partibus magnitudine ſolummodò diſtinguitur, præcipuè florum & ſeminum.

♃ Trifolium pratenſe majus album, caulibus fiſtuloſis, foliis ſubtùs punctatis nigris: Hoc Trifolium, in pratis udis & humidis creſcit. Cauliculos profert procumbentes cavos ſeu fiſtuloſos, pedales & ſeſquipedales, in quorum, ſummis alis exeunt flores ſubalbidi, ſeu Ochroleuco colore præditi. Planta hæc ad extremum pediculorum, producit terna folia ſubtùs maculata, maculis nigris ſeu fuſcis obducta.

♃ Trifolium album, ſubterraneum, Tricoccon, regium: Hoc Trifolium paritèr in udis & humentibus arvis ſeu pratis provenit. Ineunte æſtate flores fert albos, quorum ſingulis deciduis ſuccedunt, involucra, reticuli in modum, fa-

cta.

&ta. Continentia terna ſemina, utplurimùm, ſtatìm evaneſcenti flore, in terram deſcendunt rudimenta ſeminum, ſeſeque ibidem condunt, inibique perficiuntur, ideò ſubterraneum nuncupavimus. A trinis ſeminibus (utplurimùm, niſi Cœlum & ſolum refragentur) Tricoccon diximus; denique regium addidimus, quoniam, ab ipſa celſitudine ejus regia, aurelianenſi duce (ut plurimæ aliæ plantæ hic ſcriptæ) primò detectum fuit.

♃ Trifolium nodoſum capitulis Thymi: Perexigua eſt hæc ſpecies Trifolii ſub exitum hyemis: Palmares & ſeſquipalmares, edens cauliculos per terram ſtratos, ad quorum ſingulos nodos, naſcuntur floſculi minimi, quibus deciduis, ſuccedunt vaſcula plurima, ſimul ad nodos juncta: Thymi capitulis ſimilia.

♃ Trifolium Salmanticenſi affine: Hæc Trifolii ſpecies, Salmanticenſi Cluſ. adeò accedit ut idem videatur, niſi quod, (quoad omnes partes) ſit minus.

♃ Trifolium veſicarium ampullatum, Rubello flore: In humidis Sabuloſis in ora maritima Armoricæ, & prope Rupellam.

Digitized by Google

pellam, copiosè satis pullulat hæc *Trifolii* species, vix Dodrantalem excedit altitudinem, imò frequentèr ad palmarem vix pervenit. Flosculos globatìm gerit elegantes, dilutè rubri coloris. Quibus evanescentibus, succedunt ampullæ, in modum vesicarum membranacearum; continentes, semina subflava, Trifolii fragiferi instár.

♃ Trifolium cochleatum majus, fructu globoso plano: Fructum fert Cochleatum planum, globuli instar: hoc Trifolium. Et eo nomine à cœteris Cochleatis, perquàm facilè notu.

♃ Trifolium Cochleatum, spinosum, majus, spinis sursùm & deorsùm tendentibus: Hujus fructus est Cochleatus, Cœrasi magnitudine, spinis, sursùm & deorsùm tendentibus atque hac nota, facilè cognoscitur; est & hujus alia species fructu Cerasi, itidem magnitudine, spinis rectis cujus folia sunt, cordata maculata. Medicæ Arabicæ, Camerarii instár.

♃ Trifolium Cochleatum, spinosum minus, nigrum Ciliare: Hæc Trifolii spe-

cies,

Digitized by Google

cies, inter Cochleatas, radice eſt perenni, longè latéque repente, medicæ falcatæ vulgaris more; in ſummis caulium Bipedalium, multa vaſcula, ſeu fructus Cochleatos, ſpinoſos congeſtìm fert nigros, Ciliares (guillandino, ſatìs appoſitè, medica Ciliaris dicta) quia ciliorum pilis nigris, perquàm ſunt ſimilia. Unde à radice perenni, repente: Quod ſolummodò, medicæ marinæ incanæ &, Ciliari huic medicæ guillandini, convenit, inter omnes medicas Cochleatas, facilè cognoſcuntur: cœteræ quippe annuæ ſunt.

♃ Trifolium Cochleatum ſpinoſum minus δυσάρεστον, ſeu πολυσάρεστον hæc parva Trifolii ſpecies, mille paſſus à capite Bleſenſium, prope Cellam, Eremitarum, copioſiſſimè provenit. Eſt vix triuncialis, in Aridis illis locis ſeu collibus Saxoſis: fructum fert rotundum, Cochleatum (ad Piſi majoris magnitudinem) ſpinoſum, binatìm è nodis Caulium (per terram ſtratorum) exeuntem.

♃ Trifolium Lagapoïdes, flore pallido luteo: ſeu Ochroleucon: Hoc in ſylvis meridio-

Digitized by Google

meridionalibus, Bleſenſibus, ſatìs copiosè provenit. Folia edit terna, incana lanugine obſita: in faſtigio caulium erectorum, Dodrantalium, proveniunt, ſpicæ oblongæ, Lagopi inſtár: coloris ſubalbidi ſeu potiùs (ut cum Cluſio loquar) ex ὀχρωλεύκῳ compoſitæ. Semina & ſeminales capſulas aliorum, ſuæ familias more, producit.

♃ Trifolium Acetoſum, Corniculatum Indicum, rectum: Hujus caules eriguntur rectâ & ad pedalem, altitudinem, folia, flores & ſiliquas ſeu Cornicula fert, repentis lutei minoris, quoad omnes ſuas partes, multò majora. Nota inſuper hoc exoticum eſſe, cum repens luteum minus paſsìm in Britannica Gallica, ſeu Armorica provenit.

V.

Digitized by Google

V.

ꝟ VALERIANELLA noſtras, major Scabioſæ ſemine: Hæc à Valerianella Scabioſæ ſemine, à doctiſſimo Fabio columna deſcripta & picta, ſola magnitudine ſeminum, præcipuè differt: quod, duplò ſunt majora, ipſius ſeminibus, à colum. deſcripti.

ꝟ Valerianella, ſemine hirſuto, extremo umbilicato: Hæc ſemina gerit (quoad formam ſeu figuram) Uvarum acinis, ſimilia. Sed hirta & ad extremum umbilicata, ſeu magis cava.

ꝟ Valerionella ſemine hirſuto minore rotundo: Hujus ſemen eſt hirſutius, prioris ſeminibus, valdè caucalis minoris ſemini (quoad figuram) accedens. Spontè provenit in picardiæ novalibus.

ꝟ Valerionellæ vulgaris, ſpecies major Serotina, præcox: Vulgari (dicta lactuca agnina Gerardo) præcocior eſt; verna eſt planta, ineunte vere flores fert, ſubcœruleos, ſemina compreſſa, tumidi-

uſcula

uſcula tamen habet, ſtatimque diſparet. At hæc, cum frumento ſub finem æſtatis, matureſcit. Seſquipedalis eſt planta, erecta, inter ſata creſcit, flores fert albos. Semina, prioris vulgaris ſeminibus, duplò majora, & multò craſſiora & rotundiora producit. In ſatis arvis, intra ſepta Regiæ Chambort & alibi circa Bleſas, inter ſata reperitur: ſatis frequentèr.

♃ Verbaſcum ramoſum, floribus albis parvis: Folia gerit quaſi Cinere obducta, cauliculos edit Bipedales, & Tripedales ramoſos: ſuſtinentes plurimos, flores albos, parvos. Quorum, ſingulis deciduis, ſuccedunt: Valvulæ bicapſulares, mucronatæ, duplici loculamento diſtinctæ, minuta ſemina ſubruſa, hinc indè continentes aliorum more.

♃ Verbaſcum perenne nigrum floribus albis: Hæc Verbaſci ſpecies folia gerit ſubviridia, flores gerit, itidem albos, minutos, quibus paritèr, ſuccedunt capſulæ bivalves, minutè, ſemina minuta (ut cœtera ſuæ familias) in ſe continentes perennis eſt planta.

♃

☞ Verbena Urticæ folio: Planta eſt Bipedalis, aliquandò altior, è regione; ad caulium. Nodos, quadratorum, bina naſcuntur folia, mucronata, Urticæ majoris foliis, haud abſimilia; ſed ſaturatiùs virent, quoad flores, & ſemina, Verbenæ vulgari, in omnibus accedit.

☞ Veronica ſpicata, flore purpureo: Hæc à vulgari, flore cœruleo, ſolo florum colore differt.

☞ Vicia maxima annua, foliis ſerratis: Hæc ſolis foliorum inciſuris, ſerræ, ſimilibus, à vulgari Vicia maxima, annua (perperàm, authoribus Faba Græcorum dicta) differt, cum hæc capreolis donetur, Viciam ſeſe declarat, porrò nec ſiliquam, nec fructum, Fabæ, conformes habet. Judicent lectores attentè. Plura addere non expedit impræſentiarúm.

☞ Vicia præcox verna, minima Solonienſis, ſemine Hexedro: Hæc omnium hujus ſeriei, eſt planta minima, ſub exitum hyemis, in pratis, contra Regiam Chambort, floret. Florem fert dilutè rubentem, ſiliquam vix pollicarem edit quin & ſemina anguloſa, minuta, Hexe-

dram referentia, unde ei nomen à nobis inditum, datur & Lathyrus, Gramineis foliis, ejuſdem formæ, quaſi gerens ſemina, Hexedram referentia: de quo ſupra diximus.

♃ Vicia ſiliquis glabris, flore violaceo: Eſt Viciæ .ſ. Craccæ minimæ ſpecies, ſiliquis glabris Joh. Bauh.

♃ Viola Granadillæ folio: Hæc Viola exotica, ſemine & floribus, noſtrati vulgari, Martiæ, omninò conformis eſt ſimilem violaceo colore præditum, fert florem: Sed folia habet lata, ſinuoſa, in plurimos Lobos profundos diviſa Granadillæ ſeu floris paſſionis, Granadillæ majoris modo, unde ei nomen; datum.

♃ Virga aurea foliis lævibus, non ſerratis: Hæc folia gerit læviora longioraque, vulgaris Virgæ aureæ foliis: ſed nullam, in margine patiuntur ſerraturam.

♃ Virga aurea Americana, foliis ſerratis anguſtis, ſubtûs nervoſis: Hujus folia ſunt ſubtûs adeò nervoſa, & eminentèr anguſta, maximéque ſerrata, quin & radicem habet odoratiſſimam; hiſce notis notu facilis. ♃

Digitized by Google

☙ Virga aurea Americana hirſuta, radice odorata, foliis ſerratis: Hujus folia ſuperiùs, quaſi pube obducuntur, & hac nota cognitu inter cœteras, (junge quod ſint multò latiora, multoque profundiùs ſerrata, ſuperioris foliis) facilis eſt.

☙ Virga aurea humilis, Alpina: Hæc in omnibus vulgari accedit; ſola parvitate diſcrepans, eſt enim Virga aurea vulgari triplò minor. Cœtera ei par.

☙ Virga aurea, altera Solidago Saracenica: Foliis minùs ſpiſſis, conſtat, quàm Solidago vulgaris. Ad altitudinem, quatuor plùs minuſve pedum, aſſurgit. Frequentèr nobis fuit obvia, in declivibus, Regionis illius Galliæ (Morvan dictæ) inter Burgundiam & Burboniam ſitæ.

☙ Urtica racemoſa Canadenſis: Hujus folia ſunt multò obſcuriùs virentia, quàm Urticæ noſtratis majoris: inſupèr hujus folia non pungunt, quippe non copioſè pube pungente, armantur: in cœteris, ei, ſemine & floribus par.

Digitized by Google

Digitized by Google

Obſervationes Generaliores
EX
Horto REGIO
BLESENSI
Auctо, Collectæ:
BOTANICÆ
Rei, ſtudioſis, neceſſariæ, & cognitu perutiles.

Arbores, perpetua fronde, virentes.

Pomiferæ.

MALUS Aurantia. Malus Limonia. Malus Citria.

Coniferæ & Reſiniferæ.

Abies. Picea. Pinus. Pinaſter.

Digitized by Google

Cedrus Libani, ſ. Conifera.
Cupreſſus mas.
Cupreſſus fœmina.
Cupreſſus humilis ſ. Cedrus Lycea.

Glandiferæ.

Ilex Glandifera.
Suber.

Bacciferæ.

Laurus vulgaris.
Laurus Indica Baptiſt. Ferrarii.
Laurus tinus e juſque ſpecies.
Agrifolium ſ. Ilex Baccifera.
Juniperus.
Myrtus.
Alaternus.
Phillyrea.
Pyracantha.
Taxus.
Hedera arborea.

Aliæ variæ.

Ceraſus Lauri folio.
Thuya ſ. Arbor vitæ.
Arbutus.
Buxus.
Althæa Syriaca.

Reſiniferæ foliis deciduis.

Larix.
Terebynthus.
Lentiſcus.
Styrax.

Frutices

Digitized by Google

Frutices & ſubfrutices perpetua fronde virentes

BArba Jovis.
Cytiſus Maranthæ.
Geniſta Hiſpanica.
Geniſta vulgaris.
Jaſminum luteum
Roſmarinum.
Roſa ſempervirens.
Sabina.
Rhus Myrtifolia.
Althæa arborea Olbiæ.
Malva Hiſpanica.
Sedum vermicularum
Salvia Verbaſci folio & ejus ſpecies.
Atriplex Halimus, & Portulaca marina.
Chamælea tricoccos.
Laureola.
Erica ejuſque ſpecies.
Aſtragalus Bæticus.
Chamædrys Arboreſcens & vulgaris.
Colutea ejuſque ſpecies.
Ciſtus mas.
Ciſtus fœmina eorumque ſpecies.
Chamæciſtus.
Anonis montana purpurea.
Caryophyllus Creticus.
Cneorum Matth.
Ruſcus.
Hypogloſſum ſ. Laurus Alexandrina ejuſque ſpecies.
Hiſſopus.
Lavendula.
Thymum.
Satureia ſempervirens.
Pimpinella ſpinoſa ſ. Poterion.
Ruta.
Clematis Daphnoïdes major & minor.
Abrotanum.
Abſynthium.
Smilax aſpera.
Stœchas Arabica.
Thymelæa.
Tragacantha.

Herbæ perpetua fronde virentes.

Salvia hortenſis ejuſque ſpecies.
Polium ejuſque ſpecies.
Majorana.
Marum Syriacum.
Marum, Maſtic. Gallorum.
Hedera terreſtris.
Lamium luteum variegatum.
Bugula.
Prunella.
Serpillum.
Ageratum.
Aloë Americana.
Aloë vulgaris.
Yuca.
Keiri Arabum ejuſque ſpecies.
Leucoium marinum incanum ejuſq; ſpecies.
Leucoium viridi folio odoratiſſimum.
Draba repens.
Naſturtium aquaticum.
Thlaſpi ſpinoſum luteũ.
Thlaſpi ſupinum luteum.
Cochlearia.
Anagallis aquatica.
Azarum Canadenſe.
Azarum vulgare.
Bellis.
Auricula urſi.
Primula veris.
Caryophyllus ejuſque ſpecies.
Caryophyllus Holoſteus tomentoſus.
Caryophyllus minor marinus Globoſus.
Armerius.
Lychnis Cotonaria .ſ. Coronaria.
Sedum vulgare.
Sedum Arboreſcens.
Sedum Roſaceum ſerratum ejuſque ſpecies.
Sedum Vermiculatum ejuſque ſpecies.
Telephium ejuſque ſpecies.
Tithymalus Caracias ejuſque ſpecies.
Tithymalus Myrſinites, &c.

Digitized by Google

Sanicula.
Pſyllium ſempervirens.
Chamæmelum nobile.
Coris cœrulea.
Cymbalaria Italica.
Eryngium marinum.
Ficus Indica } major.
.ſ. Opuntia. } minor.
Gentianella verna.
Hypericum perfoliatum & perforatum.
Fragaria.
Jacea luſitanica ſempervirens.
Scabioſa Cretica Arboreſcens.
Tritolium hæmorrhoïdale.
Antirrhinum.
Champhorata Monſpelienſium.
Carthamus cœruleus.
Capillares omnes.
Filices quædam.
Polypodium.

Plantæ ſuccum Lacteum fundentes.

Ficus ejuſque ſpecies.
Nerium .ſ. Laurus Roſea.
Hæ duæ Arbores ſuccum reddunt; hujus viridis illius lacteus, emanat.
Apocynum ejuſque ſpecies.
Aſclepias ejuſque ſpecies.
Convolvulus ejuſq; ſpecies.
Rapunculus ejuſque ſpecies.
Campanula, Trachelium .ſ. Cervicaria ejuſque ſpecies.
Viola pentagonia, Onobrychis dicta, ejuſque ſpecies.
Cichoreum Intybus ſeu Endivia

Endivia ejuſq; ſpecies.
Lactuca ejuſque ſpecies.
Sonchus ejuſque ſpecies.
Chondrilla ejuſque ſpecies.
Hieracium ejuſq; ſpecies.
Piloſella ejuſque ſpecies.
Dens Leonis ejuſq; ſpecies.
Tragopogon ejuſq; ſpecies.
Scorzonera ejuſque ſpecies.
Carduus Scolymus ſeu Eryngium luteum.
Seſeli paluſtre lacteſcens.
Thyſſelinum Dod.

Plurimæ umbellæ montanæ, ex radice inciſa, ſuccum fundunt lacteum.

Papaver ejuſque ſpecies.
Rhœas ejuſque ſpecies.
Argemone ejuſque ſpecies,
Tithymalus ejuſque ſpecies.
Eſula & ejus ſpecies.
Cataputia, peplus, peplis, Chamæſyce.

Hæ duæ Plantæ ſequentes ſuccum fundunt rubentem.

Beta rubra.
Lapathum ſanguineum.
Chelidonium majus ſuccum reddit aureum.

Plantæ

Plantæ tuberoſæ .ſ. tuberoſa radice prædítæ.

ANEMONE, omneſque ejus ſpecies.
Pulſatilla.
Ranunculus
- Aſiaticus.
- Illyricus.
- Montanus.
- Grumoſus, tuberoſus.
- Chelidonium minus.
- Aſphodeli radice.
- Rutæ folio, &c.

Geranium
- Triſte Cornuti.
- Tuberoſum.
- Batrachoïdes.
- Hæmatodes, &c.

Cotyledon
- radice rotunda.
- radice oblonga.

Rhodia radix.
Telephium ejuſq; ſpecies.
Saxifraga alba Chelidonides.
Bulbocaſtanum.
Orchis
- Teſticulata.
- Palmata, diverſæ earundem, omnes ſpecies.

Dens Caninus.
Dentaria.
Fumaria
- Canadenſis tuberoſa : Cornut.
- radice cava,
- radice non cava.

Biſtorta ejuſque ſpecies.
Doronicum radice
- Brachiata.
- Scorpioïde.

Chryſanthemum radice
- rotunda vide
- oblonga Battatas.

Aconitum ejuſque ſpecies & anthora.
Pimpinella ſanguiſorba.
Scrophularia.
Panax Coloni .ſ. Sideritis ſtrumoſa.
Cariophyllata.
Tormentilla.
Lathyrus v. Apios
- noſtras.
- Americana Corn.
- Sylv. repens Park.

Cycla-

Cyclamen ejuſque ſpecies.
Arum.
Ariſarum.
Dracuntium.
Symphytum tuberoſum.
Solanum tuberoſum vide Battatas.
Calceolus mariæ.
Orobanche ſeu Rapum Geniſtæ & Dentaria.
Tithymalus { Apios. Pyriformi-radice.
Polygonatum .ſ. ſigillum Salomonis.
Accorus verus.
Iris tuberoſa ejuſq; ſpecies.
Ariſtolochia { rotunda. longa.

Battatas { Indiæ Occidentalis .ſ. Inhame Orientalis luſitanorum, ſpecies Convolvuli lacteſcentis. Canadenſis, Chryſanthemi ſpecies Virginiana, Solani ſpecies.

Bryonia { alba. nigra.

Gramen { Nodoſum. Typhodes ſeu Perlarum inſtar.

Cyperus { edulis Thraſi. litoreus inodorus. aliæque ejus ſpecies.

Nymphæa lutea major.

Plantæ,

Digitized by Google

Plantæ, Carnoſa radice, prædítæ:

Coniformes ſeu Pyramidales.

Asphodelus { albus. luteus.
Alectorolophos alpina.
Rapuntium, Trachelium, Campanula eorumque ſpecies.
Napus.
Rapum radice { longa. rotunda.
Raphanus ſativus { ruber niger.
Raphanus ruſticanus & ſylveſtris.
Beta rubra radice Rapæ.
Lapathum ejuſque ſpecies.
Oxalis ejuſque ſpecies.
Dens Leonis.
Tragopogon.
Scorzonera.
Centaurium majus ejuſque ſpecies.
Carlina.
Eryngium.
Fœniculum.
Siſarum.
Ænanthe ejuſq; ſpecies.
Apium ejuſque ſpecies.
Paſtinaca ſativa, latifolia.
Paſtinaca ſativa tenuifolia, Carota.

Omnes Umbella montanæ, luteos gerentes flores, radicem habent craſſam, carnoſam, Coniformem ſ. Pyramidalem cum medulla duriore in medio: Lacteſcunt ut plurimùm.

Nardus radice { olivari. oblonga.
Circium ejuſque ſpecies.
Valeriana.
Filipendula.
Mentha tuberoſa.
Pœonia ejuſque ſpecies.
Mandragora.
Mirabilis peruviana.

Arbores

Digitized

Arbores Calidæ.

Ficus.
Nerium.

Baccifera Calidæ.

Terebinthus.
Lentiſcus.
Laurus domeſtica.
Laureola, ſempervirens.
Mezereon Germanicum.
Chamelæa tricoccos.
Thymelæa.
Rhamnus ſolutivus.

Coniferæ & Reſiniferæ Calidæ.

Abies.
Picea.
Pinus.
Pinaſter.
Cedrus Libani .ſ. Conifera.
Larix.
Thuya .ſ. Arbor vitæ.
Cupreſſus.
Cedrus Lycia; cupreſſi minor ſpecies.
Juniperus.
Sabina.
Styrax ſeu Storax.

Subfrutices Calidi.

Roſa ejuſque ſpecies.
Apocynum ejuſque ſpecies.
Aſclepias ejuſque ſpecies.
Clematis ejuſque ſpecies.
Lupulus.

Plantæ

Plantæ Calidæ.

Convolvulus ejuſq; ſpecies.
Naſturtium Indicum.
Glycyrrhiza vulgaris.
Melilotus ejuſq; ſpecies.
Fœnugræcum.
Iris tuberoſa.

Bulboſæ Calidæ.

Crocus autumnalis, ſaltem ejus ſtamina.
Cepa.
Allium.
Moly.
Porrum.
Scilla.
Lilium album.
Narciſſus.
Hyacinthus.
Rapunculus ſeu Campanula ejuſque ſpecies.
Trachelium ſeu Cervicaria ejuſque ſpecies.
Gratiola.
Hypericum.
Ruta.

Siliquoſæ tretrapetalæ Calidæ.

Braſſica.
Sinapi.
Eruca.
Naſturtium ſiliquoſum.
Draba ſiliquoſa.
Eryſimum ſeu Irio.
Napus.
Rapum.
Rapiſtrum.
Raphanus ſativus ſ. Radicula.
Leucoium.
Heſperis.
Dentaria.

Siliculoſæ

Digitized by Google

Siliculoſæ tetrapetalæ Calidæ.

Thlaſpi.
Naſturtium ſiliculoſum.
Burſa paſtoris.
Cochlearia.
Iberis.
Lepidium.
Draba ſiliculoſa.
Raphanus ſylveſtris.
Raphanus hortenſis ſeu Ruſticanus.
Myagrum.
Rapiſtrum Monoſpermon.
Coronopus Ruellii.
Veronica ejuſq; ſpecies.
Polygala.
Perſicaria urens.
Agrimonia.
Lactescentes Cichoraceæ & pappoſæ, per ſe omnes (quippe amaræ) ſunt Calidæ.

Corymbiferæ Calidæ.

Conyza.
Abrotanum.
Abſynthium.
Artemiſia.
Tanacetum.
Matricaria.
Chamæmelum.
Millefolium.
Ptarmica.
Ageratum.
Pimpinella & Agrimonoïdes Col.
Umbelliferæ omnes ſunt magis aut minùs Calidæ.

Verticillatæ Galeatæ, Calidæ.

Mentha.
Calamintha.
Clinopodium.
Pulegium.
Marrubium.
Dictamnus verus.
Pſeudodictamnus.
Bætonica.
Stachys.

Digitized by Google

Stachys.
Sideritis.
Lamium.
Meliſſa.
Tragoriganum.
Origanum.
Salvia.
Horminum.
Acinos.
Meliſſa peregrina.
Polium.
Chamædrys.
Chamæpitys.
Roſmarinum.
Hyſſopus.
Lavendula.
Stæchas.
Majorana.
Thymum.
Serpillum.
Satureia.
Ocimum.

Calidæ variarum claſſium.

Centaurium minus.
Pulſatilla.
Anemone ejuſque ſpecies.
Ranunculi omnes, exceptis Perpaucis.
Arum.
Ariſarum.
Dracunculus major.
Capſicum ejuſq; ſpecies.
Tithymali omnes, Eſulæ & Cataputia.
Sedum vermiculatum acre.

Digitized by Google

Purgantes omnes ſunt Calidæ: vide infra Purgantes.

Plantæ Frigidæ:

Pomiferæ Scandentes.

Melo.
Cucurbita.
Pepo
Citrullus ſ. Anguria.
Cucumis.

Dum ſucco alimentitio abundant, refrigerant; at ſemina (quippe oleo abundant) exſiccata, calefaciunt: Contra commune Dogma.

Beta.
Blitum.
Mercurialis.
Spinachia.
Lapathum.
Acetoſa.
Biſtorta.

Lactescentes Cichoraceæ & pappoſæ per accidens ſunt Frigidæ, dum ſucco alimentitio abundant: at quatenus amaræ Calefaciunt, & apperiunt.

Acanthus.
Plantago.
Holoſteum.
Coronopus, ſpecies Plantaginis.
Pſyllium.
Viola.
Malva.
Althæa.
Mercurialis.
Mandragora.
Hyoſcyamus,
Portulaca.
Sedum ejuſque ſpecies omnes, præter acre.
Telephium

Telephium.
Cotyledon.
Saxifraga { Chelidonides / Aurea.
Sanicula.
Papaver ejuſque ſpecies, per accidens dum ſucco alimentitio abundant; ut & Cichoraceæ & pappoſæ lacteſcentes refrigerant: de quibus alibi prolixiùs diſcutietur.

Arbores, frutices, ſubfrutices, & Plantæ purgantes.

Sorbus, ornus dicta, ex qua manna elicitur.
Tamarindus.
Caſſia ſolutiva.
Rhamnus Catharticus.
Malus Perſica ejus flos.
Senna.
Geniſta.
Roſa { pallida / alba Moſchata } utriuſque flos.
Laureola.
Mezereon Germanicum.
Chamelæa Tricoccos.
Tarton Rare.
Alypum montis Cete.

Plantæ purgantes ſucco.

Aloë vera.
Convolvulus ejuſq; ſpecies (*per excellentiam ſcammonia Syriaca ſ. Antiochena eſt hujus ſpecies; ex qua fit diagredium.*
Cucumis Aſininus *ex quo extrahitur elaterium.*
Tithymalus, Eſula & Cataputia: *quorum ſucco lacteo, adulteratur diagredium.*

Purgantes Radicæ.

Filix fœmina.
Polypodium.
Lapathum ejuſq; ſpecies: *per excellentiam Lapathum, Rhabarbarum eſt præſtantiſſimum.*
Helleborus niger & albus.

Purgantes, aut vomitum cientes foliorum infuſione, aut decoctione.

Sedum vermiculatum,
Narciſſus.
Gratiola.
Braſſica rubra, criſpa.
Centaurium minus.
Linum Catharticum.
Cnicus .ſ. Carthamus, *ſemine purgat.*

Plantæ Venenatæ.

Nerium.
Apocynum.
Aſclepias.
Lolium.
Colchicum.
Cicuta.
Pyrethrum.
Solanum lethale.
Aconitum.
Fungi lethales.

Plantæ

Digitized by Google

Plantæ Vulnerariæ.

Pimpinella.
Apium.
Cicuta.
Marrubium.
Betonica.
Stachys.
Syderitis.
Salvia.
Horminum.
Hedera terreſtris.
Chamædrys.
Chamæpitys.
Bugula, Brunella.
Bellis.
Milleſolium.
Verbena.
Polygonum.
Polygonatum.
Alchimilla.
Sanicula.
Caryophyllata.
Plantago.
Pſyllium.
Ariſtolochia.
Cynogloſſum.
Symphytum.
Pulmonaria.
Nicotiana.
Craſſula ſeu Telephium.
Sedum.
Scabioſa.
Conſolida regalis.
Virga Aurea, Doria, & Solidago Saracenica.

Plantæ Culinariæ.

Legumina, fructus horum inſerviunt Culinæ.

Phaſeolus.
Piſum.
Lathyrus.
Ochrus.
Lens.
Faba.
Lupinus.
Glycyrrhiza. *Hujus radix expeditur in Culina & Pharmacis.*

Digitized by

Plantæ Culinariæ.

Cærealea, horum grana inſerviunt Culinæ.

Triticum.
Secale.
Hordeum.
Oriza.
Avena.
Panicum.
Milium.
Panicum Indicum, Maiz dictum.

Plantæ Culinariæ.

Bulbi & folia, harum inſerviunt Culinæ, & Pharmacis:

Cepa.
Allium.
Moly.
Porrum.
Scilla.

Plantæ Culinariæ.

Radices harum inſerviunt Culinæ.

Rapuuculus ſeu Campanula.
Napus.
Rapum.
Raphanus { Sativus .ſ. Radicula. / Niger Icon. Lob.
Raphanus ruſticanus.
Beta rubra Romana, ſeu Rad. Rapæ.
Trogopogon.
Scorzonera.
Chryſanthemum tuberoſum, *vide* Battatas de Cannada.
Solanum Eſculentum, *vide* Battatas de Virginea.
Convolvulus, Indiæ Occidentalis Inhame, *vide* Battatas ſuprâ.
Siſarum.

Siſarum.
Paſtinaca latifolia.
Paſtinaca tenuifolia ſeu Carota.
Apium ſeu Petroſelinum.

Plantæ quarum folia inſerviunt Culinæ, cocta.

Braſſica & ejuſque ſpecies.
Thlaſpi Holeraceum.
Naſturtium hortenſe.
Eruca ſativa.
Beta vulgaris ſativa.
Bonus Hendricus.
Spinachia.
Lapathum:
Acetoſa.
Lactuca ejuſque ſpecies.
Cichoreum, Intybus ſeu Endivia.
Tragopogon.
Scorzonera.
Calendula, hujus etiam flores.
Urtica minor & major.
Apium ejuſque ſpecies.
Cerefolium.
Myrrhis odorata.
Tanacetum.
Mentha.
Majorana.
Roſmarinum.
Thymum.
Satureia.
Borrago.
Bugloſſum.
Portulaca.

Plantæ quarum folia uſurpantur in Acetariis.

Lactuca.
Acetoſa.
Cichoreum.
Pimpinella ſpicata minor.
Braſſica minor (Brocali Italis) dicta.
Valerianella ſeu Lactuca agnina.
Viola pentagonia.

Digitized by Google

Coronopus hortenſis.
Sedum vermiculatum inſipidum, ejuſq; ſpecies.
Naſturtium { hortenſe. aquaticum.
Ptarmica .ſ. Dracunculus hortenſ. Aſtragon dictus.
Cerefolium.
Myrrhis nova æquicolorum & odorata.
Apium.
Cinera aculeata Hiſpanica.
Apium Italicum ſeu Seleri.
Lactuca Romana.
Cichoreum criſpum.
Harum quatuor folia & coſtæ ſeu Nervi medii, dealbata, evadunt tenera.
Chrithmum { Umbelliferum. Chryſanthemũ.
Folia utriuſque ſale, & aceto condita, Menſæ inſerviunt, hyeme pro acetario: quemadmodum,
Flores inexpanſi Capparis, Geniſtæ, & Naſturtii Indici, eodem modo ſale, & aceto Macerati hyemalibus Embammatis inſerviunt ut & fungi Eſculenti.
Cucumis, *hujus fructus, rudimenta: ſale, aceto & ſemine Anethi, condiuntur & Menſæ inferuntur.*

Plantæ Culinariæ.

Quarum ſemina Culinæ inſerviunt.

Sinapi.
Eruca.
Fœniculum.
Cuminum.
Carui.
Aniſum.
Coriandrum.
Anethum.

Plantæ.

Digitized by Google

Plantæ Culinariæ.

Harum Turiones inſerviunt Culinæ.

Aſparagus.
Lupulus.
Sambucus.
Capſicum, *hujus ſiliquarum aut capſularum pelliculæ, loco Piperis in Gallaprovincia, Gallia Narbonenſi, aliiſq; locis, inſerviunt Culinæ. Quin & in Africa aceto, ſale & Saccharo condiuntur, & ab incolis avidè expetuntur pro Cordiali optimo.*
Cinera ſativa; *capita hujus, Culinæ inſerviunt.*

Plantæ Culinariæ.

Harum Fructus pulpoſi inſerviunt Culinæ, Menſæve.

Melo.
Pepo,
Cucumis.
Citrullus ſeu Anguria.
Cucurbita.

Arbores

Digitized by Google

Arbores Pomiferæ, quarum Fructus inſerviunt Culinæ, Menſæve.

Malus ejuſq; variæ ſpecies.
Pyrus ejuſque diverſæ ſpecies.
Malus { Aurantia. Limonia. Citria.
Malus { Cydonia. Punica.
Sorbus.
Ficus.
Meſpilus.
Morus.

Arbores, Oſſeam nucem, carne obductam, habentes; quarum Fructus exterior & interior inſerviunt Menſæ.

Malus { Perſica ejuſque diverſæ ſpecies. Armeniaca.
Prunus & ejus variæ ſpecies.
Ceraſus ejuſque variæ ſpecies.
Olea.
Palma.

Arbores Nuciferæ, harum fructus interior, inſervit Menſæ ſeu Culinæ.

Amygdalus.
Nux juglans.
Corylus.
Caſtanea.

Arbores

Arbores ſeu potius Frutices, quorum Fructus inſerviunt Culinæ, Menſæve.

Berberis.
Labruſca.
Vitis vinifera.
Rubus Idæus.
Rubus.
Ribes.
Uva criſpa.
Myrtillus Germanorum.
Fragaria, *hujus plantæ fragæ, inſerviunt Menſæ.*

FINIS.

Digitized by Google

Hallucinationes

CASPARI BAUHINI

IN

PINACE,

Tam in digerendis,

Quam denominandis Plantis:

ITEM

Animadverſiones, in tres Tomos, univerſalis Hiſtoriæ, Plantarum;

JOHANNIS BAUHINI.

Quibus acceſſit

Dialogus inter Socium Collegii REGII *LONDINENSIS, Greſham* dicti, & *Botanographum* REGIUM.

Præludiorum Botanicorum *pars altera.*

AUTHORE

ROBERTO MORISON Medico & Profeſſore *Botanico*, ſereniſſimo *CAROLO* ſecundo, magnæ *Britanniæ*, &c. REGI.

Doctis (in Arte) ſcribo:
Indoctos docebo.

Hallucinationes

CASPARI BAUHINI

IN

PINACE,

Tam in digerendis,

Quam denominandis Plantis.

Pag. 27 Eſamum & Eryſimum cœreale, malè inter cœrealia propriè dicta, ob uſum collocantur poſt Lapathum, Acetoſam, Biſtortam, quippe ſemine triangulari præditum Eryſimum cœreale, collocari debet. Ad ſiliquoſas Heteroclitas ſeu impropriè dictas, tranſmittatur Seſamum.

Pag.

Pag. 34. *Column. I.* Acorus adulterinus, malè à Bauhino, cum Acoro julifero ſeu vero, collocatur, cum adulterinus, Iridis florem gerit; ad Iridum latifoliarum claſſem, cum Ger. & Eyſt. relegandus, ut rationi conſentaneum videtur.

Pag. 40. *Col. II.* Iris tuberoſa amandetur ad Iridum tuberoſarum claſſem, nec aptè collocatur inter Iridem bulboſam & Sicyrinchium.

Pag. 47. *Col. II. Plant. III.* Hyacinthus Indicus tuberoſa radice, malè inter bulboſas propriè dictas collocatur.

Pag. 80. Lilium Aſphodeli radice, aptiùs (mea opinione) cum Aſphodelo & Phalangio, pag. 28, vel 29. collocaretur.

Pag. 89. Bauhin. collocat Rapum, & pagina 95. collocat Rapiſtrum Napum, & interjicit pag. 92, 93, 94. Campanulatas, Rapunculos ſcilicet, & Campanulas, Trachelium ve; ſiliquoſas à ſiliquoſis disjungit, quod non eſt faciendum. Inſuper Rapiſtri ſiliquæ ſunt craſſæ fungoſæ, articulatæ, ideò aptiùs poſt Raphanum ſativum; vel radiculam, collocaretur Rapiſtrum.

Pag. 91. *Col. I. Pl. VIII.* Batatas Camotes Hiſpanorum

Hiſpanorum Cluſ. hiſt. eſt verus & genuinus Convolvulus, quare malè cum Aloë inter ſiliquoſas collocatur, quamvis radices illius ſunt edules. De his quæ vidi & colui, judicium fero, de aliis exoticis in illo capite, mihi nunquam viſis: nihil dico.

Pag. 92. *Col. I Plant. XIII.* Campanula lutea Linifolia Ger. repetitur p. 214. col. I. pl IX. ſub Campanula lutea Linifolia Adv. Lob. Linum ſylveſtre luteum foliis ſubrotundis C. B. Pin. cujus vera eſt ſpecies & flore & ſemine; Campanula autem non eſt, quia nec lacteſcit nec ſemen minimum Lucidum (aliarum Campanularum more) fert.

Pag. 94. *Col. II. Pl. X.* Campanula foliis Echii C.B.Pin. & Trachelium θυρσοειδὲς Cluſ. ipſi Cam. & pag. 254. col.II. pl.VII Echium Alpinum luteum C. B. Pin. Trachelium θυρσοειδὲς Cluſ. Trachelium ſeu Campanula eſt, quippe lacteſcit non autem Echium; bis tamen repetitur.

Pag. 95 *Col.II.Pl.V.* Rapiſtrum Italicum ſiliquis longiſſimis C.B.Pin. ſylveſtre Irionis folio Col. Idem repetitur pag. 101. col.II. pl.V. Eryſimum anguſtifolium majus C. B. Pin. Rapiſtrum ſylveſtre Irionis folio Col.

Pag.95.Col.I.Pl.I. Rapiſtrum flore luteo C. B. Pin. eſt vera Sinapi ſpecies, (Rapiſtrum eſt utplurimùm ſiliqua articulata) ergò ad ſiliquoſas Tetrapetalas, Sinapi. Erucam, Naſturtium, Drabam cum reliquis ſequentibus, amandetur. Rapiſtrum Monoſpermum cum non ſit ſiliquoſa Tetrapetala planta; ſed, ſiliculoſa, ad calcem ſiliculoſarum, debet referri.

Pag.96.Col.II Pl.V.& pag.97. Raphanus ruſticanus & aquatici duo, non debent collocari inter Raphanos. Cum ſiliquas Raphani ſativi, ſeu radiculæ non ferant, ſed capſulas binas ſiliculaſve Lepidii aut Iberidis inſtár.

Pag.97. Col.I. Pl.III. Dentillaria (mea opinione) non ad Lepidium referri debet, cum nec florem nec capſulam ſeminalem, conformem Lepidii capſulis, aut ſiliculis habeat.

Ibidem Lepidium & Iberis malè collocantur inter ſiliquoſas cum capſulares ſeu ſiliculoſæ ſint, debent ad calcem Thlaſpios transferri.

Pag 98. Dracunculus, ejuſque ſpecies; malè coliocantur inter capſulares & ſiliquoſas

quoſas à C.B. in Pin. quare ad Corymboſas .ſ. Corymbiferas, veluti ageratum, Millefolium & hujuſmodi affines; relegentur Dracunculus hortenſis, & Ptarmicæ.

Pag.99.Col.I. Pl.X. Eruca maritima Italica, ſiliqua haſtæ cuſpidi ſimili C. B. Prodr. Cakile Serapionis Lob. duæ plantæ diverſæ,quoad ſemina confunduntur à Bauhino,una ſemen fert planum; altera,cuſpidi haſtæ ſimile: ut clarè videre licet figuram in Prodr. à Bauhino exhibitam, eamque ſi quis cum Lobelii figura in Iconibus comparet.

Pag. 100. Reſeda & luteola cum non ſint Tetrapetalæ ſiliquoſæ, malè diſponuntur inter Erucam & Eryſimum, Tetrapetalas ſiliquoſas.

*Pag.*101.*Col II.Pl.V.* Eryſimum anguſtifolium majus C. B. Pin. Rapiſtrum ſylveſtre Irionis folio Col. Bis repetitur,nempè ſuprâ pag 95.col.II.pl.V. & hic p.101.col.II.pl.V.

Pag. 101. Perſicaria ſeu Hydropiper malè inter Tetrapetalas ſiliquoſas colloca-tur cum ſit ſemine triangulari prædita, ad Lapathum, Biſtortam,Polygonum, aliaſque (ejus generis) referri debet.

Pag. 102. Piper Indicum cum non ſit planta Tetrapetala ſiliquoſa, malè hic collocatur à Bauhino, potiùs ad Solanum referatur; cujus, eſt ſpecies à flore & ſemine, ſeminiſque Thecis, ſeu involucris. Si objiciatur qualitas Urens; quæ non reperitur in ulla Solani ſpecie, facilè reſpondetur Ranunculos quoſdam, eſſe inſipidos ſeu dulces; quamvis maxima eorum pars, ſit cauſtica prædita facultate.

Pag. 103. Naſturtium hortenſe latifolium & criſpum malè illic diſponuntur, cum non ſint ſiliquoſæ, ut cœteræ illius ſectionis, ſed ſiliculoſæ, ergò ad ſiliculoſarum claſſem amandentur.

Pag. 104. *Col. II.* Naſturtium Petreum folio burſæ poſtoris; item Naſturtium pumilum vernum; Naſturtium Alpinum capſula Naſturtii hortenſis; & Naſturtium Oſyridis folio, cum ſint ſiliculoſæ plantæ, ad Thlaſpios, Drabæ, Cochleariæ Burſæ paſtoris calcem, aut claſſem, referantur; non autem confundendæ cum ſiliquoſis.

Pag. 106. *Col. II. Pl. V.* Thlaſpi umbellatum Naſturtii folio Monſpeliacum C. B. Pin. Narbonenſe umbellatum Tab. Ger. eadem planta

planta repetitur pag. 107. col. II. plant. III. Thlaſpi Alyſſum dictum maritimum C. B. Pin. Umbellatum Narbonenſe Ger.

Pag. 108, & 109. Burſæ paſtoris, ſimilis ſiliquoſa major & minor, non debent inter Burſæ paſtoris ſpecies reeenſeri cum non ſint ſiliculoſæ: amandentur ad ſiliquoſas. Ibidem Myagrum ſiliqua longa, ad ſiliquoſas referatur cum Gerardo: dicatur Eruca ſylveſtris anguſtiſolia.

Pag. 109. Drabæ ſiliculoſæ ad ſiliculoſas & Tetrapetalas plantas: ſiliquoſæ ad ſiliquoſas referantur, ut Tollatur confuſio generum.

Pag. 110. *Col. II Pl. VIII.* Draba flore cœruleo Galeato, malè denominatur Draba à Bauhino, cum nec ſit planta Tetrapetala, ſiliquoſa, nec ſiliculoſa. Rapunculi ſpecies eſt; Galeato flore violaceo: quemadmodùm flos Cardinalis colum. eſt Galeato Coccineo flore. Vide Rapuntium Urens Solonienſe in horto Regio Bleſenſi aucto, & inter plantas non ſcriptas à me exhibitas.

Pag. 110. Alliaria malè inter Drabam & Cochleariam collocatur: ad ſiliquoſas nempè Heſperides ſeu violam Matronalem referatur.

Pag. 113. Iſatis malè illic collocatur, inter Tetrapetalas ſiliquoſas, & multò ineptiùs, Anil ob uſum colorandi, cum ſit perfectè planta leguminoſa.

Pag. 114, *&* 115. Spinachia malè pro genuino Lapatho recenſetur cum non habeat ſemina triquetra, Lapathi inſtàr, ſubjungi debet Lapatho, quemadmodum bonus Hendricus.

Pag. 116. *Col. I. Pl. V.* Graviter Hallucinatum fuiſſe C. Bauhinum in Pinace conjicio, dum inter Lapatha collocat Apocynum Syriacum, ſeu Bedelzar Arabum, cum ne vel notulam Lapathorum generi communem, habeat: ſed Apocynum à lacte (dum rumpitur) manante, ſiliquiſque Sericea lanugine, refertis, ſeminibuſque. Apocynum ſeſe prodit, vel Medriocritèr in arte Botanica verſato; ad Apocyni claſſem, referri debere apparet. Hujus, lanugine Sericea, Togæ cubiculares pretioſiores apud Turcas componebantur calidæ & læves, quæ ab aliquot annis non pretioſa illa intertextura quæ Gallicè dicebatur Wat. ex ſeminalibus hujus Bedelzar involucris, præparata; ſed Goſſipio ſeu Xylo ex Americæ

inſulis

insulis adjacentibus: nunc communitèr fiunt. Prædictæ Togæ cubiculares & vili pretio væneunt.

Pag. 117. *Col. II. Pl. V.* Rhaponticum folio Helenii incano, malè hic disponitur cum Rhabarbaro (quod per excellentiam Lapathum est,) ergo reddi debet ad calcem plantarum capitatarum (quarum flores abeunt in pappos) ut sunt jacea Stœbe Cyanus, &c. Ibidem idem dicendum de Centaurio majore, ejusque speciebus quæ omnes debent amandari ad classem antedictam. Observa insuper Rhabarbarum esse per excellentiam optimum Lapathorum: à semine triangulari Lapathi simili.

Pag. 123. *Col. II. Pl. III, & IV.* Lactuca montana purpurocœrulea, major & minor, malè inter Lactucas collocantur, cum & semen & conformationem partium, cum Soncho habent.

Pag. 124. Lampsana non est tam Soncho affinis quam Cichoreo: quippe ut Cichoreum semina solida, angulosa, non autem in pappos abeuntia (ut Sonchus) producit. Ergò magis appositè dixisset C. Bauhinus Cichoreo affinis Lampsana domestica.

Pag. 122, & 131. Jacobæa & Senecio cum ſunt ſtellatæ & pappeſcentes debent ad ſtellatas & pappeſcentes, Aſterem, Conyzam, Doronicum, Helenium, &c. Non autem jungi cum lacteſcentibus, pappoſis.

Pag. 126. *Col.II. Pl.I.* Hieracium majus folio Sonchi, vel Hieracium Sonchites potiùs Sonchus dici debet quam Hieracium, nam Sonchus eſt utplurimùm altior, caulis fiſtuloſior folia læviora, flos ejus Hieracii flore citiùs evaneſcit: per terram (aliorum Sonchorum more) ejus radix repit.

Pag. 126. *Col.I. Pl.I,II,III.* Cichoreum ſylveſtre luteum & quæ ſequuntur ſunt Hieracia, quia pappeſcunt.

Pag. 127. Hieracia quatuor priora primæ Col. ſunt verè Sonchi à ſemine & tota facie & modo creſcendi.

Pag. 228. *Col.I. Pl.I & II.* Hieracium ſemine adunco C. Bauh. Hieracium falcatum col. I. pl. II. & Hieracium minimum nonum Cluſ. pag. 127. col. II. pl. VIII. & quæcunque alia reperiuntur Hieracia, quorum flores non abeunt in pappos, ut horum trium ad Cichoreum reducantur quia Cichoreum, à lacteſcentibus pappoſis hac ſola nota differt.

fert. Quod ejus flos non abit in pappos idem eſto judicium de Chondrilla verrucaria ſeu Zazintha pag. 130. col. I. pl. IX. cum non pappeſcit: non eſt Chondrilla. Sit cum Cluſio Cichoreum verrucarium idem dicendum de Lampſana, quæ eſt magis affinis Cichoreo, quam Soncho: propter ſemen Solidum anguloſum, Cichorei ſemini, ſimile.

Pag. 130. Chondrilla Verrucaria ad Cichoreum mandetur; propter prædictam rationem & jungatur prædictis Hieraciis. Eaque appellando ſic; Cichoreum falcatum, Cichoreum ſemine adunco; Cichoreum minimum, caule ad florem tubuloſo, mutato nomine Hieracii, in Cichoreum ut diximus anté.

Pag. 130. *Col. II. Pl VI.* Chondrilla cœrulea Cyani capitulo: cum hujus ſemen non abit in pappos non eſt Chondrilla, ergò ad Cichoreum cum prædictis reducendum judicarem; cum Lacteſcit, nec flos ejus abit in pappos, nec poteſt eſſe Cyanus, nec Chondrilla ut clariùs deducemus in Tabulis affinitatis noſtris.

Pag.

*Pag.*130. *Col.II. Pl.III.* Chondrilla rara purpurea cum non Lacteſcit per antedictam rationem, nec pappeſcit, non debet admitti pro Chondrilla. Ad Centaurium capitatum reducerem, & Centaurium capitatum ciliare annuum foliis laciniatis ſerratis appellari velim.

*Pag.*131. *Col.I.Pl.III.* Senecio pinguis incanus; Jacobææ eſt ſpecies, cum Jacobææ more, florem gerit ſtellatum multis petalis latis, diſcum luteum ex multis ſtaminulis conflatum, cingentibus. Nota inſuper hac p.131.malè collocari(ut alibi diximus) Jacobæam & Senecionem cum lacteſcentibus pappoſis, cum ad pappeſcentes non lacteſcentes pertineant: cumque eis collocari debent.

*Pag.*132.*Col.II.Pl.I,& II.* Tanacetum montanum inodorum flore majore & minore. Sunt Bellidis Leucanthemæ ſpecies: nec cum Tanaceto aliquid habent affinitatis niſi in diſpoſitione foliorum: ergò alibi aptiùs diſponi poterant, ſub nomine Bellidis Tanaceti folio: Deſumendo nomen à potiore & nobiliore parte, flore ſcilicet, ſeu modo florendi.

Pag.

Pag. 138. *Col. I.* Ambroſia Campeſtris repens, Coronopus Ruellii: Malè cum Ambroſia odorata collocatur, cum ſit planta Tetrapetala ſiliculoſa, ſubjungi debet ſiliculoſis Drabæ, Cochleariæ, Thlaſpi, Veronicæ, &c.

Pag. 138 *Colum. II. Pl. III.* Botrys Chamædryoïdes malè ad Botryos claſſem à Bauhino reducitur, cum ſit ex flore & ſemine ſeminumque capſulis, Chamædrys: ad hujus claſſem (cum Mattheolo) redigo. Nota quod Botrys Ambroſioïdes appoſitè quadrat; cum Atriplice, quoad diſpoſitionem florum & ſeminum, & ſi Atriplex odora diceretur per me non ſtabit. Quemadmodum fecimus nos in diſpoſitione Atriplicis in Tabulis cognationis noſtris.

Pag. 141. *Col. II. Pl. IV.* Millefolium aquaticum foliis Abrotoni, Ranunculi flore & capitulo C. B. Pin. Ranunculus Tricophyllon aquaticus medio-luteus Col. bis repetitur hic & pag. 180. col. II. pl VI. Ranunculus aquaticus capillaceus C. B. Pin. Ranunculus Tricophyllon aquaticus medio-luteus Col. Etiamſi Bauhinus bis citavit Columnam, debuit in poſteriori loco tantùm

tantùm citari. Sunt duæ, Ranunculi ſpecies, omninò diſtinctæ. Ut quilibet obſervare poſſit; cum intuendo figuram Columnæ & Icon. Lob. atque copiosè uterque hic Ranunculus in aquis paludoſis conſpicitur, vegetans.

Pag. 142. Conſolida Regalis & pag. 144, & 145. aquilegia & Nigella debent ad venenatas, & ſiliquiferas amandari ut clariùs demonſtrabimus poſteâ.

Pag. 144. Chelidonium majus cum ſit planta Tetrapetala, ſiliquoſa, ad Tetrapetalas ſiliquoſas, eſſet referendum.

Pag. 146. *Col. II. Pl. III.* Cuminum ſylveſtre capitulis globoſis C. B. Pin. non eſt Umbella propriè loquendo cum non habet bina ſemina ſimùl juncta, ſed cum folia Umbellæ in modum diſpoſita habet, ad calcem Umbellarum: cum Pimpinella amandetur.

Pag. 147. *Col. II. Pl. VII.* Chaa malè à Caſp. Bauh. in Pin. pro Fœniculi ſpecie exhibetur: cum ejus folia ex Japonia Sinenſiumque, finibus, ad nos delata, aqua macerata & extenſa, latiora ſunt; Bellidis foliis, declarantque hanc, non eſſe Umbellam, ferulaceis Fœniculaceiſve foliis prædítam.

Pag.

Pag. 147. Anethum ad umbelliferas, (quarum ſemina ſunt rotunda & foliacea) eſſet referendum: nulla habita ratione foliorum Fœniculo, accedentium.

A pag. 147, *ad pag.* 165. Adeò confuſus eſt ordo Umbellarum: Bauhini nec confuſio tolli poterit quoad Tabulas, cognationis affinitatiſve noſtras, publico, exhibuerimus.

Pag. 148. Pyrethrum flore Bellidis non obſtante cauſtica facultate, ad Corymboſas ſeu Corymbiferas eſt referendum: non inter Umbellas collocandum.

Pag 149. *Col II. Pl. III.* Peucedanum, eſt Umbella, Gramineis foliis, ſemine folioſo ſubrotundo prædita, parùm ergò rationi congruit; Peucedanum minus Caſ. B. Pin. eſſe Peucedanum: cum nec Graminea habet folia, nec ſemina foliacea, ſeu compreſſa, ſed ſtriata, & folia tenuitèr diviſa. Selinum montanum pumilum (cum Cluſio) aptiùs deſignabitur.

Pag. 150. Nota Bauhinum quatuor genera diſtincta Umbellarum, ſub uno & eodem genere (Dauco ſcilicet) comprehendiſſe. Quæ ſuis ſunt reddendæ ordinibus: ut ex doctrina noſtra, alibi tradenda (quin

&

& hic) cuilibet perſpicuum evadet.

Daucus foliis Fœniculi tenuiſſimis, ad Daucum ſeu Carotam vel Paſtinacam tenuifoliam, reduci poteſt, quippe ſemina villoſa, (Dauci vulgaris in modum) producit. Tres autem ſequentes Umbellæ; (ſub Dauci nomine) cum ſint foliis ferulaceis, & ſemine ſtriato minore donatæ; pro Dauco montano, ſeu Saxifraga tenuifolia haberi poſſunt. Quippe quod, Saxifragiæ tertiæ Matth. ſeu Dauci montani (Bauhino ſic dicti) habent, & folia & ſemina ſtriata minora. Sex autem ſequentes (incipiendo à Dauco montano, Apii folio majore) cum ſint ſemine compreſſo foliaceo, ſubrotundo, ut Peucedanum vel Oreoſelinum, ad Umbellas, foliis plurifariàm diviſis, ſemineque foliaceo rotundo, ditatas: ſunt religandæ. Ultimus ſeu 13. Daucus pratenſis, Millefolii paluſtris folio C. B. Pin. Daucus pratenſis Dalechampii hiſt. Lugd. ex radicum & ſeminum ſtriatorum ſtructura, declarat ſeſe eſſe Oenanthen. Unde nobis, potiore jure, Oenanthe Millefolii foliis (in pratis humidis vegetans, more aliarum ſuæ claſſis) denominabitur. Hæc diximus ut legentibus & profundiùs intuentibus

entibus, innueremus, malè distributam, hanc Dauci subalternam sectionem, à Casp. Bauhino : dum 4. distincta genera Umbellarum (ut suprâ clarè docuimus) sub uno & eodem militare vult signo.

Pag. 151. *Col. I. Pl. II.* Pastinaca tenuifolia sylvestris Umbella majore C. B. P. sylvestris Daucoïdes apula Col. Hæc planta ex semine oblongo aspero etiamsi folia gerit Pastinacæ tenuifoliæ: sese declarat potiùs, ad Myrrhidis classem pertinere.

Pag. 151. *Col. I. Pl. VI.* Pastinaca sylvestris angustifolia, fructu Echinato C. B. Pin. Pastinaca Echinophora apula Col. à semine debet hæc planta Umbellifera ad Crithmum maritimum, spinosum, sibi consimile; ad finem Umbellarum remitti: cum Pastinacæ marinæ seu Chrithmi marini spinosi habeat seminis involucrum, spinosum; semenque paritèr striatum, eódem debet referri.

Pag. 151. *Col. II. Pl. IV.* Gingidium, foliis Pastinacæ latifoliæ C. B. Pin. Gingidium 1. Dod. malè huic generi adaptatur, cum non sit semine striato minore (ut Gingidium) præditum, nec etiam caucalis dici debet.

Ut

Ut plurimi Neotericorum ſcriptorum dixêre: cum ſemen non habet Echinatum; caucalidum more. Ergò concludendum cum diſpoſitione ſeminis foliacei, ſubrotundi, in margine granulati (Tordylii apuli minimi Col. more) tum poſitione foliorum Lobatorum, ſub Tordylii ſigno militare debere: & magis appoſitè Tordilium Syriacum, ſemine majore granulato; deſignabitur.

Pag. 152. *Col.II. Pl. III.* Chærophyllo nonnihil ſimilis C. B. Pin. Perchepier Anglorum Lob. Cerefolium eſt Umbella, foliis plurifariàm diviſis, ſemineque roſtrato prædita. Ergò Perchepier Anglorum, Chærophyllo, nihil ſimilis, potiùs quam nonnihil ſimilis Bauhino. Non eſt Umbella, ergò longè arcenda, ſed cum in omnibus & per omnia Alchimillæ, illi conveniunt notæ eſſentiales: eò mandanda, cum doctiſſimo Fab. Columnâ.

Pag. 152. *Col.I.* Caucalis major Cluſ. hiſt. malè à Bauhino, pro Caucalidis ſpecie habetur, cum ejus ſemen non ſit (Caucalidum ritu) Echinatum: quod, eſt generica & eſſentialis Caucalidis nota. Aliò referen-

da

da nec aptiùs quam ad Tordylium (cum docto Jacobo Plateau, qui ejus ſemen ad cluſium miſit) referri poſſe autumno.

Pag. 152. *Col. II. Pl. XII.* Caucalis folio Peucedani C. B. Pin. Caucalis Anglica Lob. cum hæc Umbella ſit ſemine minore ſtriato donata. Dauci montani aut Saxifragiæ 3. Matth. in modum : ad ejuſdem ſpeciem amandari debet: neutiquàm ad Caucalidem, cujus non fert Symbolum.

Pag. 153. *Col. II. Pl. VI.* Apium montanum folio ampliori C. B. Pin. & montanum nigrum C. B. in Pin. Oreoſelinum Pariſienſium Lugd. & Apium ſylveſtre lacteo ſucco turgens, & Apium peregrinum Cicutæ folio C. B. Pin. omnes hæ quatuor Apii ſpecies, neutiquàm ſunt genuina Apia, quippè Apium eſt planta umbellifera, ſemine ſtriato minore, ornata ; at priores ſunt ſemine foliaceo ſubrotundo affectæ. Ergò aliò tranſmittendæ ; ut in Tabulis noſtris cognationis, præſtabimus. Nota inſuper ibidèm Bauhinum, duas diſtinctas conſtituere ſpecies: Oreoſelinum montanum folio ampliori, & Pariſienſium nigrum ; quæ, una & eadem eſt planta ; diverſa, tantùm natalibus.

Pag. 156. Poſt Imperatoriam admitti debet Imperatoria nigra, Tab. pagin. 186. col. I. pl. V. contenta, Aſtrantia nigra Lob. Icon. & Obſ. Elleborus niger Saniculæ folio, major & minor C. B. Pin. malè hæc planta in Ellebori claſſe collocatur, cum Umbellæ, omnes habeat conditiones requiſitas, flores in Umbella, Pentapetalos, quorum ſingulis evaneſcentibus, ſuccedunt, bina ſemina oblonga aſpera, quod clariùs intuentibus apparebit, quoniam è ſummo faſtigio Caulium exeunt; foliola ſubviridia, latiuſcula, ſtellatim cingentia baſin, radiorum ſemiuncialium, ſuſtinentium floſculos minimos, pentapetalos: Perſicæ florum coloris. Aſtrantiæ nomen ſortita eſt hæc planta, quoniam folia in 5. Lobos obſcuriùs virentes (Saniculæ inſtar) diviſa gerit. Saniculæ illi adaptarunt nomen ſed Imperatoria nigra, Aſtrantia dicta, Saniculæ folio, potiore jure denominabitur.

Pag. 157. *Col. I. Pl. II.* Libanotis major & altera ſeu vulgatior, cum utriuſque ſemina ex pluribus alis foliaceis, rotæ molendinæ fluviatilis inſtar; conſtent: eaſdem declarant, ex Laſerpitii familia oriundas; ſeu

genus

genus ſuum deſumere, quippe eadem, cum illis (quoad formam & figuram) ſunt ſemina.

Pag. 158. *Col. II. Pl. VI.* Panax Aſclepium ſemine folioſo C. B. Pin. Panax Aſclepium Apulum Col. cum ſemen habeat rotatìm, ſeu circularitèr ala foliacea cinctum, Thapſiæ more, ad Thapſiam redigatur: non autem ad Libanotidem, cujus ſemen, nulla ala membranacea pellucida cingitur.

Pag. 160. *Col. I. Pl. VI.* Pimpinella ſanguiſorba major, minor, hirſuta & minor lævis, & omnes quot fuerint ſanguiſorbæ Pimpinellæve, quum non ſunt Umbellæ (propriè loquendo) non debuere locari inter Umbellas(propter foliorum cum Pimpinella Saxifraga, ſimilitudinem) ſed ad calcem Umbellarum, cum aliis ejuſdem Farinæ, amandari: propter ſimilem foliorum, Lobatorum diſpoſitionem.

Pag. 161. *Col. II. Pl. VII.* Cicutaria paluſtris tenuifolia non habet bina ſemina, ſimùl juncta aliarum Umbellarum more, ergò non eſt Umbella propiè dicta.

Pag. 161. *Col. I.* Seſeli Creticum majus; Creticum minus; Creticum fructu majore;

Creticum minimum: Cum Seſelios, nota eſſentialis & Generica, ſit producere ſemen ſtriatum; atqui hæ quatuor plantæ umbelliferæ, ſemina producunt foliacea, ſeu membranacea, rotunda; ergò debent collocari cum congeneribus ſuis: Paſtinaca latifolia, Sphondylio, &c. ſub Tordylii nomine Generico.

Pag. 162. *Col. I. Pl. XI.* Seſeli paluſtre lacteſcens, propter præcedentem rationem, à Seſelios familia averruncandam, & ad Umbellas, ſemine foliaceo ſubrotundo foliis plurifariàm diviſis (cum Oreoſelino & Thyſſelino) revocandum eſſe judico.

Pag. 162. Oenanthe Apii folio Caſp. B. Pin. anguſtifolia Lob. nota quod C. Bauhinus pro Synonimo, attulit Cyminum bulboſum Plinii Col. quod eſt bulbocaſtanum aut ſaltem ejus ſpecies, nihil habens affinitatis cum Oenanthe, ut ait doctiſſimus Columna.

Pag. 163. *Col. I. Pl. V.* Oenanthe Hederæ folio C. B. Pin. Oenanthe Myconi hiſt. Lugd. non eſt Umbella, cum flores & ſemina Ranunculi producat. Optimè Ranunculus Thalictri folio, Aſphodeli radice, cum Parkinſono nominabitur. *Pag.*

Pag. 163. *Col. II. Pl. VII.* Filipendula vulgaris, cum non ſit Umbella, ſed tantum à foliorum cum Umbellis ſimilitudine; ad finem Umbellarum (tanquam affinis) debet collocari. Ibidèm Pedicularis ſeu Alectorolophos, cum ſit Galeata, ad finem Galeatarum, verticillatarum, collocanda. Ibidèm Barba Capræ, cum ſit tantum Umbellis affinis, à diſpoſitione foliorum, cum Thalictro, Filipendula, Pimpinella, &c. In fine Umbellarum addenda: Ergò inconſideratè, laborioſiſſimus Bauhinus, Filipendulam pedicularem & Barbam Capræ, interjecit inter Oenanthen & Valerianam, propriè dictas Umbellas: cum nil habent hæ plantæ, commune, cum propriè dictis Umbellis, niſi folia, quæ, non ſunt primò & principalitèr, partes eſſentiales, niſi per accidens tantùm; inſervientes fructificationi. Nota inſuper Plant. IX. & X. ejuſdem Columnæ ad caput ſequens, debere referri.

Pag. 164. *Col. II. Pl. VII.* Valeriana Græca; propter antedictam rationem, ab ipſo doctiſſimo & laborioſiſſimo Caſp. Bauhino, illic obſervatam: eſt aliò remittenda. Obſerva hanc Valerianam, nil habere

cum Valeriana propriè dicta, commune, præter folia; ergò ad finem itidèm Umbellarum, cum cœteris (propter conformitatem foliorum)referri debet.

Pag. 165. *Col. I,& II.* Valerionellæ omnes, non ſunt Umbellæ propriè loquendo, quia non habent bina ſemina (aliarum Umbellarum propriè ſic dictarum more) ſimùl juncta.

Pag. 168. *Col. II. Pl. I.* Solanifolia Circœa dicta,major & minor: quâ ſimilitudine Solanis annumerantur, non poſſum aſſequi. Semen fert verrucoſum; ergò ad Agrimoniam,Saniculam vulgarem ibidèm: (in locis Udis, ſylvoſis & umbroſis) habitantem, referatur. Ibidèm. Qua ratione mirabilia Peruviana, ad Solanum reduci poſſunt; non ſatis aſſequor. Eſt planta certè Heteroclita: quemadmodùm plurimæ aliæ exoticæ. Solano etenìm ſatis malè cohæret. Nos exactiùs exhibebimus (in Tabulis noſtris cognationis, univerſalis noſtræ methodi) hujus diſpoſitionem.

Pag. 172. Hypecoum malè inter Papaver & Argemonen collocatur: exilis cum ſit planta, foliis,Fumariæ adinſtàr, diſſectis; ſiliquas

ſiliquas edit longas articulatas, eſt Heteroclita planta hæc.

Pag. 176. *Col. II. Pl. V.* Anemone nemoroſa, flore majore Caſp. Bauh. Pin. à quinta Plant. ſecundæ Columnæ, pag. 176. ad Pl. X. ſecundæ Columnæ pag. 177. Ranunculi omnes ſunt. Cum, omnium flores abeunt in ſemina aculeata racematìm compacta, ſive orbicularitèr ſive ſpicatìm. Anemones propriæ dictæ, flos abit in pappos, ventoque aufertur. Ergò omnes contentæ à V. Pl. Col. II. pag. 176. ad X. Pl. Col. II. pag. 177. ſub Anemone, perperàm diſponuntur, debent conjungi cum Ranunculo nemoroſo luteo, pag. 178. Col. I.

Pag. 180. *Col. II. Plant. VI.* Ranunculus aquaticus medioluteus Col. bis repetitur hic & ſuprâ pag. 141. vide Millefolium aquaticum foliis Abrotani, Ranunculi flore & capitulo.

Pag. 180. *Col. I. Plant. VI.* Ranunculus aquaticus Cotyledonis folio, Caſp. B. Pin. Cotyledon aquatica Dod. cum ſemen non gerat Ranunculi, non eſt reputandus pro Ranunculo, ideò aliò remittendus; nec aptiùs quam ad aquaticas Heteroclitas plantas referri poteſt.

Pag. 179. *Colum. II. Pl. VI.* Ranunculus Aconiti folio, flore albo multiplici C. B. Pin. debet transferri ad Columnam primam pag. 182. & jungi Plantæ VII. nempé. Ranunculo montano Aconiti folio, albo flore majore C. B. Pin.

Pag. 182. *Colum. II. Pl. IX.* Ranunculus montanus Aconiti folio, flore globoſo C. B. Pin. eſt potiùs Aconiti ſpecies, cum ſiliculas ſeu Thecas ſeminales, membranaceas, Aconiti ritu habet, ſemenque inibi contentum. Nec ſemina racematìm in capitulum pungens compacta: ut Ranunculus *gerit.* Quare ad Aconiti claſſem retulimus appellandó Aconitum, Ranunculoïdes ſecundum, quemadmodùm Aconitum hyemale, à flore Ranunculi, Aconitum Ranunculoïdes primum ſeu hyemale; Thecas ſeminales. Utrumque Aconiti more habet.

Pag. 183. *Colum. II. Pl. XVI.* Aconitum racemoſum, Chriſtophoriana, dictum. Malè inter Aconita recenſetur, cum nec florem Cucullatum, nec Thecas membranaceas (aliorum Aconitorum more) gerit: ſed cum ſit planta baccifera; ad Solani potiùs genus referenda.

Pag.

Pag. 184. *Col. I. Pl. I.* Aconitum Pardaleanches ſeu Thora major & minor, ſunt Ranunculi ſpecies; ergò ad Ranunculi genus, montani pag. 182. pertinent, atque ibi locari debent.

Pag. 184. Doronicum cum ſit planta ſtellata, pappeſcens amandetur ad conſortes, Aſterem, Conyzam, Helenium, Tripolium, paginis 265, & 266.

Pag. 286. *Col. I. Pl. V.* Helleborus niger Saniculæ folio major & minor, debent amandari ad Imperatoriam, inter Umbellas, cujus ſunt ſpecies: vide pag. 156. ubi prolixiùs illic egimus.

Pag. 186. *Col. I. Pl. IX.* Helleborus niger tenuifolius Buphthalmi flore, ſub eodem genere militat cum Adonide; cujus eſt ſpecies, utraque, Ranunculorum claſſi accedere debet.

Pag. 197. Tuſſilago Petaſites & Cacalia, ad papeſcentes amandentur, ſcilicèt Eupatorium, virgam auream, Stœchadem Citrinam, Serratulam, 263. pag.

Pag. 201. *Col. II. Pl. I.* Leucoium Alyſſoïdes Clypeatum majus C. B. Pin. Alyſſum Dioſcoridis Dod. debet cum Lunaria Græca

ca collocari: item Leucoium incanum *siliquis* rotundis C. B. Pin. Eruca peregrina Cluſ. debent omnes hæ Leucoii ſpecies platolobi, ſubjungi Leucoiis ſiliquoſis.

Pag. 200, 201, 202. Malè collocantur Leucoium ſiliquoſum & viola matronalis híc. Cùm ad ſiliquoſas reliquas tetrapetalas debent transferri. Videlicèt Sinapi, Erucam, Irionem, ſive Eryſimum, &c. p. 99.

Pag. 210. *Col. I. Pl. VII.* Caryophyllus pratenſis flore laciniato ſimplici, ſeu flos Cuculi C.B. Pin. malè inter Caryophyllos collocatur, à Bauhino, cum omninò ſit Lychnis, à flore, ſemine & capſula Pyriformi.

Pag. 210. *Col. II.* Caryophyllus holoſtius ejuſque ſequaces, uſque ad Caryophyllum montanum. Alſines & Lychnidis Seriei, debent adjungi: Quum flores gerant quinis petalis, ad medium bifidis, conſtantes: capſulaſque Pyriformes. Non autem Caryophylli in modum Cylindriacas.

Pag. 211. *Col. I. Pl. I.* Linaria ſcoparia eſt planta ſui generis: nec præter folia, aliquam affinitatis notam habet cum Linaria. Ibidèm, idem dicendum de Oſyride fruteſcente Baccifera, C. B. Pin. ſeu Caſia Poëtica Monſpelienſium Lob.

Pag.

Pag. 213. *Col II. Pl. XV.* Linaria montana floſculis albicantibus C. B. Pin. Anonymos tenuifolia Cluſ. per eandem rationem antedictam, non eſt Linaria.

Pag. 213. *Col. I. Pl. VI, & VII.* Linaria folioſo capitulo luteo, major & minor: nil niſi folia Linariæ gerunt. Florem virgæ aureæ inſtàr, ex plurimis ſtaminulis, luteis conflatum, habent; qui poſtea abit in pappos: ergò jure merito, virga aurea folliis Linariæ appellabitur. Linariæ nota eſſentialis, eſt flos Calcari donatus, rictum compreſſus, habet flos. Semen, binis capſulis, Apum alvearia referentibus incluſum habet Linaria; ut in deſcriptionibus obſervavimus; & ut prolixiùs trademus, quando notas eſſentiales & genericas unicuique generi adaptabimus, in Tabulis noſtris cognationis. Linaria autem ſcoparia: Oſyris baccifera fruteſcens: Linaria montana ſeu Anonymos Cluſ. Linaria aurea Trag. cum nihil horum poſſident, alibi ſunt collocandæ.

Pag. 214. *Col. II. Pl. VI.* Linum minimum ſtellatum non eſt Linum: à nota eſſentiali Lino convenienti, cum nec florem, nec ſemen Lini habet. Ibidèm Col. I. Pl. IX.

Linum

Linum ſylveſtre luteum foliis ſubrotundis C. B. Pin. Campanula lutea linifolia Lob. bis repetitur, hic & ſuprâ pag. 93. Colum. I. Pl. XIII. nimirùm Rapunculus nemoroſus, anguſtifolius parvo flore C. B. Pin. Campanula lutea linifolia Lob. Ger. vide ſuprâ pag. 93.

Pag. 215. *Col. II. Pl. V.* Polygala repens nivea C. B. Pin. repens Lob. non debet flori Ambervali jungi ſed Polygono (Herniaria) dicto. Ibidèm Onobrychis Arvenſis ſeu Campanula Arvenſis erecta C. B. Pin. cum ſit Campanula (à flore) plantaque lacteſcens, florem, capſulam ſeminalem, triquetram, & ſemina pellucida (ut Campanula) gerit: Campanulis debuiſſet jungi.

Pag. 216. Stœchas Arrabica, Lavendula & pag. 217. Roſmarinum, malè claudunt ſectionem, cum ſequentem, verticillatarum galeatarum, ſectionem debeant incipere; quippè verticillatæ galeatæ, verticillatis galeatis (in eâdem ſectione) addendæ.

Pag. 218. *Col. II. Pl. XIII.* Hyſſopifolia C. B. Pin. cum nec ſit planta galeata nec galeatæ aut verticillatæ fert capſulam, parùm

rùm aut nihil cum Hyſſopo habet affinitatis aliò ergò releganda: Fortè ad Euphraſiam.

Pag. 221. *Col. I, & II. Pl. I, & II.* Ageratum foliis ſerratis & non ſerratis C. B. P. malè inter galeatas & verticillatas plantas collocantur, cum ſint Corymbiferæ .ſ. Corymboſæ. Adde quod odor & ſapor, non conſiderantur in diſponendis plantis. Ibidèm Ageratum ſerratum Alpinum C. B. Pin. Ageratum purpureum Lugd. malè etiam hic diſponitur: cum non ſit Galeata planta, nec capſulas, Galeatæ in modum: verticillatìm gerit. Aliò etiam releganda: eôdem cum Hyſſopifolia ad finem Euphraſiæ amandetur. Capſulas fert longas innumeris ſeminibus minutulis, ſcatentes ut Euphraſia, ſed nec hæc, nec Hyſſopifolia florem habent Galeatum, ſed quinis petalis diſtinctis conflatum.

Pag. 222. *Col. I. Pl. V.* Dictamnus albus vulgo, ſeu Fraxinella C. B. Pin. Fraxinella Dod. cum nil habeat affinitatis, nec in foliis nec in floribus, multò minùs in capſulis ſeminalibus, cum verticillatis Galeatis, aliò eſt remittenda, nec aptiùs (mea opinione) quam ad ſectionem venenatarum, ſiliquis membra-

membranaceis præditarum ; Aconitum (videlicèt) Delphinium, Staphydem-agriam, &c. (id genus) aliás.

Pag. 225. *Col. I. Pl. II.* Clinopodium arvenſe Ocymi facie C. B. Pin. planta eſt verticillata ; verticillis per Caules rariùs diſpoſitis, donata, per intervalla : at majus ſ. vulgare verticillos denſiùs diſpoſitos, per intervalla habet : Ergò ad Ocymi finem referri poteſt : Eſt enim Ocymi ſylveſtris ſpecies. Origanum Canadenſe fiſtuloſum Corn. in ejus locum ſufficio : tanquam plantam (inter odoriferas) magìs illi convenientem, eandem appellando, Clynopodium Canadenſe fiſtuloſum, & non Origanum, cum quo nil habet affine.

Pag. 225. *Col. II. Pl. V.* Clinopodium Alpinum hirſutum C. B. Pin. Clinopodium Alpinum Ponæ bis repetitur hìc, & p. 247. Col. II. Pl. VI. Teucrium Alpinum coma purpuro-cœrulea C. B. Pin. Clinopodium Alpinum Ponæ.

Pag. 226. *Col. I. Pl. II.* Mentha hortenſis Corymbifera C. B. Pin. coſtus hortorum Lob. malè à Bauhino ob odorem, Menthæ Seriei adjungitur, cum jure merito ad Corymboſas,

rymboſas, ſive Corymbiferas, videlicèt Tanacetum, Ageratum Milleſolium, &c. (id genus) alias plantas Corymboſas, debet amandari.

Pag. 229. *Col.* I. *Pl.* VI. Calamintha arvenſis verticillata C. B. Pin. malè inter Calaminthæ ſpecies recenſetur, quippè Calaminthæ, verticilli, rariùs diſponuntur, hùjus verticilli, denſiùs, per intervalla diſpoſiti ſunt. Ideò qui Mentham (Siſymbrium) hortenſem & ſylveſtrem, cum Mentha pulegii odore, & Mentha aquatica exigua & pulegio ſylveſtri Lugd. conjunxerit: aſſignando toti generi nomen Menthæ denſiùs verticillatæ, meliùs egerit.

Pag. 231. *Col.* I. *Pl.* V, *&* VI. Lamium maculatum C. B. Pin. & Lamium linea alba notatum C. B. Pin. pro una, duæ in medium adferuntur plantæ, cumque cuilibet propiùs conſideranti, una & eadem apparebit planta. Ineunte enim hyeme, albis maculis, folia conſperguntur. Exeunte autem hyeme, linea alba, per folii Nervum medium diſcurrente, notantur: æſtate una & altera, ſpoliata, folia habet viridia. Ibidèm Lamium foliis Caulium ambientibus

majus

majus & minus, pro una & eadem habeantur planta, cum ſpecifica, qua diſtinguantur, nulla aſſignatur nota.

Pag. 231. *Col. II. Pl. XI.* Lamium peregrinum .ſ. Scutellaria C. B. Pin. Aſtragaloides Cornuti. A forma capſulæ ſeminalis, apprimè referente imaginem aut formam tali vulgaris, qui mulierum calceis ſubjicitur: quæ exſiccata & apperta, Scutellam exprimit, unde ei nomen novum genus conſtituens Scutellaria; dictum, ad finem Galeatarum & verticillatarum, rejiciendum: qui conſtruxerit, gratum opus fuerit; quinque ſunt plantæ quæ ſub novo hoc genere militare poſſunt, ſub nomine Scutellariæ, gratiola latifolia Ger. Lyſimachia Galericulata Lob. Scordium Berardi ſeu Scorodotis Plinii Cluſ. Scutellaria Americana Malvæ folio. Et Lamium peregrinum ſeu Scutellaria C. B. Prodr. & ſi quæ aliæ verticillatæ Galeatæ hujus Farinæ detegantur, huc paritèr revocandæ ſunt.

Pag. 232. *Col. II. Pl. I.* Urtica aculeata foliis ſerratis, & altera foliis non ſerratis, malè Urticæ nominantur, cum ſint verticillatæ, Galeatæ propriè loquendo, ergò

ad

ad Sideritidis claſſem jure pertinent. Ibidèm Urtica urens maxima, minor, & pilulas ferens; cum non ſint galeatæ, ſed verticillatæ, aliquomodò racemi, verticillatim diſponuntur, ergò ad finem galeatarum, verticillatarum remittantur.

Pag. 232. Alyſſum verticillatum foliis profundè inciſis, malè disjungitur à Marrubio; cum ſit omnibus ſuis partibus, verum & genuinum Marrubium.

Pag 233. Euphraſia ejuſque ſpecies, cum ſint galeatæ, ſed non verticillatæ propriè loquendo, ad finem verticillatarum & galeatarum rejici debent: Ut clariùs demonſtrabimus in Tabulis noſtris cognationis.

Pag. 234. Nota Bauhinum inadvertentèr, confundere Euphraſiam majorem ſylveſtrem purpuream & latifoliam, cum ſunt duæ diſtinctæ plantæ; ut apparet ex duabus diverſis Iconibus Columnæ: Ut cuilibet intuenti Fab. Columnam; evidentèr apparet.

Ibid. Melampyrum ejuſq; ſpecies, cum ſint galeatæ plantæ, ſed non verticillatæ propriè dictæ; ad calcem galeatarum ſuerunt re-

jiciendæ; & non confusè (cum Bauhino) *in* medio sectionis collocandæ.

*Pag.*234. *Col.II. Pl.VI.* Melampyrum luteum minimum C.B P. Perpusillum luteum Lob. Adv. quam inconsideratè Bauhinus, Sequendo Lobellium, hanc plantam(quam depictam in adversariis exhibuit Lobellius) denominat. Melampyrum, legentium sequentia esto judicium. Melampyrum est planta galeata tritici referens grana. Hæc autem tenella planta,Ranunculi est species, à gustu, sapore, flore, quinis petalis, plurima staminula cingentibus, conflato. Cui deciduo, succedit spica coniformis, falcata gerens semina: Folia fert ferulacea. Nec apparentèr Lobellius, qui primus erravit, omnesque alios authores errare fecit: florentem hanc vidit plantam, multo minùs bene observavit spicam, (etiamsi pictam in adversariis exhibet) quæ neutiquàm refert Melampyri spicam; hactenùs de hac planta egimus, inter non scriptas à nobis exhibitas.

Pag. 235. Serratula malè disponitur hic in medio galeatarum,verticillatarum: cum ejus flos abit in pappos, plurimis staminulis fistulosis,

fiſtuloſis, virgæ aureæ modo, conſtat. Ergò ad virgam auream, ejuſque claſſem tranſponi ibique collocari debet. Ibidèm Scrophularia, cum nec ſit galeata nec verticillata planta, ſed bicapſularis; plurima minuta in ſe continens ſemina, ad claſſem bicapſularium, releganda floret : Nempè digitalem, verbaſcum, Blattariam, &c. (id genus) aliás.

Pag. 240. *Col. II. Pl. I, II, III.* Verbaſcum latis Salviæ foliis C. B. Pin. & anguſtis Salviæ foliis C. B. Pin. & acuto Salviæ folio C. B. Pin. perperàm hæ plantæ à Bauhino & authoribus aliis, ad Verbaſcum reducuntur, ab incana lanugine foliorum, Verbaſci foliis quodammodo accedentium ; cum Verbaſci omnis, flos eſt integer, in margine quinquefariàm diviſus, cui deciduo ſuccedit bicapſularis Theca, ſeminalis. Multa hinc indè continens ſemina minutula, ut videre licet in digitali, Blattaria, Scrophularia, Gratiola, Hyperico &, id genus, aliis ejuſdem claſſis. Sed more noſtro denominatio deſumi debet à nobiliore parte, flore ſeu florendi modo: Cum galeatum & verticillatìm diſpoſitum fert florem. Adinſtàr Salviæ,

capſulaſque ſeminales gerit Salviæ, quatuor ſemina oblonga, anguloſa. Verticillatarum omnium ferè ritu, Salvia fatua: cuilibet hæc noſtra dicta conſideranti apparet. Et ideò Salvia ſylveſtris Verbaſci latis foliis; & Salvia ſylveſtris Verbaſci anguſtis foliis; & Salvia ſylveſtris Verbaſci acuto folio; Impoſterùm à nobis deſignantur.

Nota quod Verbaſcum & Blattaria, pag. 239, & 240. malè inter verticillatas diſponuntur, cum ad capſulares bivalves debeant amandari; quod præſtabimus in generali noſtrâ hiſtoria plantarum, per Tabulas cognationis diſponendarum.

Pag. 241. Æthiopis, (ſine ratione) diſjungitur ab Horminis: Cum omnes, Hormini habeat partes; Horminum Æthiopicum incanum appelletur.

Pag. 245, & 246. Nota Lyſimachiæ caput, ſeu diſtributionem eſſe confuſiſſimam apud Bauhinum in Pinace. Primò, occurrit Lyſimachia lutea major; quæ florem quinis aut Senis petalis conſtantem fert; cui flaccido ſuccedit, capſula rotunda; Piſi medii magnitudine in qua continentur, minuta plurima rotunda ſemina. Ephemeron,

flore

flore & ſemine & capſula ſeminali, valdè huic Lyſimachiæ accedit. Secundò, occurrit Lyſimachia Virginiana, florem magnum, Unifolium ſeu Monopetalum convolvulorum more; cui deciduo, ſuccedunt Cornicula coniformia, maturo ſemine, in plurimas ſeſe findunt partes, & copiosè ſemina emittunt: ex loculamentis, adinſtàr Papaverum Corniculatorum. Tertio loco, occurrunt Lyſimachiæ propriè dictæ ſiliquoſæ, quarum flores exeunt in ſummitate filiquæ; quaternis, quinis petalis conſtantes; quibus Marceſcentibus, ſuccedunt ſiliquæ biunciales triunciales, ſeſe in quinque diſtincta loculamenta, apperientes, & ſemina plurima pappoſa (ut Salix) fundunt. Quarto loco, ſeſe offert nobis conſideranda, Lyſimachia ſpicata purpurea, florem fert galeatum rubrum; cui ſuccedit capſula ſeminalis (Euphraſiæ inſtar) continens plurima ſemina minuta; atque eô nomine haud abs re, eſſe judicarem; eandem Euphraſiam perennem aquaticam, flore purpureo, denominari. Quinto loco, nobis conſideranda exhibetur, Lyſimachia ſpicata cœrulea, quæ aliarum Veronicarum

more, florem fert tetrapetalum: cui ſuccedit ſilicula, bicapſularis, Burſæ paſtoris ſeu Veronicæ ſimilis; & eſt Veronica ſpicata cœrulea recta. Sextò, Lyſimachia cœrulea galericulata vel Gratiola cœrulea Caſp. B. Pin. Lyſimachia galericulata Lob. hæc eſt galeata verticillata, ſcutellata, per conſequens ad novum genus verticillatarum referri debet. Et ſcutellaria paluſtris cœrulea nobis dicenda: ita hæc diviſo Lyſimachiæ ad ſex diverſas claſſes diſtinctas, pertinet; variè à nobis per Tabulas cognationis, diſtribuenda.

Pag. 247. *Col. II. Pl. V.* Clinopodium Alpinum Ponæ bis repetitur, hìc & pag. 225. Col. II. Pl. V. vide ſuprâ paginam 225.

Pag. 247, 248, 249. Teucrium, Scordium, Chamædrys, Chamæpitys malè hic cum Lyſimachia veronica, inter veronicam & Alſinen, collocantur; cum inter galeatas, verticillatas, propriè dictas: eſſent collocandæ.

Pag. 248. *Col. I. Pl. V.* Chamædrys Alpina Ciſti flore C. B. Pin. Camædrys montana fruteſcens durior Ad. Lob. cum hæc planta nec florem galeatum, nec capſulam ſemina-

lem

lem verticillatìm diſpoſitam habeat; Chamædrys non poteſt dici: cum autem florem gerit, ſex petalis albis conflatum, cingentibus diſcum medium, cui evaneſcenti ſuccedunt globuli, hirſuti, Caryophyllatæ more, ergò appoſitè ſatis appellabitur Caryophyllata; Chamædryos, folio duriori.

Pag. 248. *Col. II. Pl. VIII.* Chamædrys unicaulis ſpicata Bauh. in Pin. Trixago Apula unicaulis Col. hæc florem gerit galeatum luteum, ſed capſulam habet & ſemina inibi contenta, Euphraſiæ: non eſt Chamædrys, quia non habet capſulas in margine quinquefariàm diviſas, continentes ſingulas, quaterna ſemina. Appelletur ergò Euphraſia Apula unicaulis magìs apté. Ibidèm Pl. XII. Chamædrys ſpuria major latifolia C. B. Pin. Veronica maxima Lugd. uſque ad Plantam XVIII. omnes hæ ad Veronicæ claſſem debent amandari; cum omnes, Veronicæ gerunt teſſeram, flores (ſcilicet) tetrapetalos, ſiliculas bicapſulates, Veronicæ more. Nec Chamædryos ſpecies, accenſeri poſſint: quippè nec galeatæ nec verticillatæ ſint.

Pag. 250. *Col. I. Pl. I, II, III, & IV.* Alsine Hederulæ folio Casp. B. Pin. Veronicæ folio floſculis Cauliculis adhærentibus C. B. Pin. Chamædrifolia floſculis oblongis pediculis inſidentibus; & Alsine Triphyllos cœrulea. Hæ omnes quatuor, notam eſſentialem Veronicæ habent; nempè flores tetrapetalos, & ſiliculas bicapſulares, ſepto medio membranaceo, duplicem ſeriem ſeminum diſterminante: ergò ſunt veræ & genuinæ Veronicæ annuæ. Sic nobis denominanda, Veronica Hederulæ folio; Veronica floſculis adhærentibus, cauliculis; Veronica Chamædrifolia floſculis pediculis longis inſidentibus; & Veronica Triphyllos cœrulea; ex dictis patet, omnes eſſe veras & genuinas Veronicas annuas; non autem ad Alsines, genus pertinent: Ut clarius alibi docebimus.

Pag. 150. *Col. II. Pl. VI.* Alsine ſcandens baccifera C. B. Pin. Cacubalus Plinii, eſt Alsine Heteroclita: Folia & flores habet Alsines, fructum ſeu baccas, Solani.

Pag. 151. *Col. I. Pl. XVI.* Alsine affinis Androſace dicta major C. B. Pin. Androſace

ſace altera Matth. nec florem nec capſulam ſeminalem, fert Alſines; nota eſſentialis Alſines eſt, habere florem quinis petalis conflatum; quorum ſingula ſunt bifida, Lychnidis more; cui evaneſcenti, ſuccedit capſula, Pyriformis continens plurima ſemina rotunda: Quod non conſpicere licet in hac planta, magis accedit ad auriculam Urſi. Multas alias plantas puſillas adjunxit Bauh. hìc quæ propter exilitatem (quamvis ſunt Heteroclitæ) aliò impræſentiarùm non remittemus. Ibidèm Col. II. Pl. VI. Alſine litoralis Portulacæ foliis C. B. Pin. Alſines Pelagicum genus Cluſ. nota quod ſit Rubiæ genus, ex ſemine.

Pag. 152. *Col. I. Pl. I.* Anagallis aquatica major, folio ſubrotundo Caſp. B. Pin. aquatica minor folio ſubrotundo; aquatica major folio oblongo; aquatica minor folio oblongo; aquatica folio rotundo non crenato; & aquatica anguſtifolia: Omnes hæ ſex plantæ, cum florem gerant tetrapetalum, ſiliculaſque bicapſulares, Veronicæ (procul dubio) ſunt veræ & genuinæ ſpecies. Ideo Veronicæ appellentur aquaticæ; ſicque denominentur. Veronica aquatica

major

major folio ſubrotundo; Veronica aquatica minor folio ſubrotundo; aquatica major folio oblongo; aquatica minor folio oblongo; aquatica folio rotundo non crenato; & Veronica aquatica anguſtifolia: & ſic tolletur gravis lapſus; Bauhini & cœterorum authorum. Quam habent affinitatem cum Anagallide videant oculati Botanici, & diligentèr examinent mecum.

Pag. 252, *&* *pag.* 253. Elatine ejuſque ſpecies: quum calcari & rictu hiante; & capſula ſeminali, duplici loculamento diſtincta, prædita ſint. Quæ eſt nota generica & eſſentialis Linariæ; Linariæ ſeſe veras & genuinas arguunt eſſe ſpecies; quod benè innotuit doctiſſimo Fabio Columnæ, dum Elatinen, Linariam Hederulæ folio appellat. Ibidèm pag. 253. Col. II. *Pl. II.* Elatine folio acuminato, in baſi auriculato C. B. Pin. Linaria Hederulæ folio Col. bis repetitur, hìc & pag. 306. Linaria Hederulæ folio Col. Cymbalaria C. B. Pin. Nota inſuper Cymbalariam eſſe veram & genuinam Linariæ ſpeciem. Quia Linariæ nota eſſentialis & generica, omninò ei quadrat.

Pag.

Pag. 254. *Col. II. Pl. VII.* Echium Alpinum luteum C. B. Pin. Trachelium θυρσοειδὲς Cluſ. & Campanula foliis Echii C. B. Pin. Trachelium θυρσοειδὲς Cluſ. ipſi Cam. Pag. 94. Col. II. Pl. X. bis repetitur. Campanula eſt: (quippè lacteſcit, ſemenque ruſum, pellucidum habet) non vero Echium.

Pag. 257. *Col. II. Pl. X.* Cynogloſſum minus C. B. Pin. Cynogloſſa minor montana ſerotina altera Plinii Col. & pag. eâdem Columna, eâdem Pl. VII. Cynogloſſum medium C. B. Pin. Cynogloſſa minor montana ſerotina altera Plinii Col. eâdem planta, bis repetitur.

Pag. 258. *Col. I.* Lithoſpermum arundinaceum forte Dioſcoridis & Plinii Lachryma Jobi Dod. frumentum eſt. Proindè ad cœrealia amandetur; quippe cum Lithoſpermo ordinario nil habet affine.

Ibidèm Pag. 258. *Col. II. Pl. VII.* Lithoſpermum arvenſe radice rubra C. B. Pin. Anchuſa degener facie Milii Solis Lob. ad Anchuſam referatur, cum nec florem nec ſemen fert Lithoſpermi.

Pag. 259. *Col. I. Pl. I.* Lithoſpermum Linariæ folio, Germanicum C. B. Pin. Li-

naria

naria altera botryoïdes montana Col. planta eſt Heteroclita: nil cum Lithoſpermo vulgari habet commune: niſi ſeminis duritiem. Eſtque ſui generis planta. Ibidem Col. II. Pl. V. Symphytum minus Borraginis facie C. B. Pin. Borrago minima herbariorum Lugd. malè à Bauhino inter Symphyta collocatur, cum ex flore, tum ex ſemine, ſit vera Borraginis ſpecies.

Pag. 260. Conſolida media ſive Bugula & conſolida minor ſive Brunella, malè diſjunguntur à verè galeatis & verticillatis, quum iis, nota eſſentialis & generica, galeatarum & verticillatarum, perfectè convenit. Ergo ad Sectionem quartam Bauhini pag. 217. cum cœteris ſuæ conditionis amandentur.

Pag. 262. Bellis malè hic collocatur: quare ad Corymboſas referatur leucanthemas; Chamæmelum, Cotulam, &c.

Ibidem. Piloſella ejuſque ſpecies malè hic collocantur, cum ſint plantæ lacteſcentes, earumque flores abeunt in pappos, cum Piloſella ſit omnibus notis parvum Hieracium: ad finem Hieracii pag. 129. relegenda. Nec cum Bellide aut Gnaphalio, quid habet affinitatis.

Pag.

Pag. 266. Aſteris omnis, flos debet abire in pappos; non minùs quam Conyzæ. Hæc eſt nota eſſentialis Aſteribus, & cœteris ſtellatis pappeſcentibus, communis. Aſt. quatuor à me obſervantur Aſteres, recitati à Bauhino in Pinace, quorum, flores non abeunt in pappos: ſuntque hi; Aſter foliis ad florem rigidis: Aſter Atticus foliis circa florem mollibus C. B. P. ſeu Cernuus Col. Aſter luteus folio glabro & crenato C. B.P. Atticus Dod. & Conyzoïdes Monſpelienſium. Ergò eô ipſo nomine quod, ſemina, ſolida gerant, à pappoſis, per conſequens ab Aſteribus arceantur: prædictæ quatuor plantæ: & ad Corymbiferas relegandæ.

Pag. 272. *Col. II. Pl. II, & III.* Jacea Oleæ folio, flore majore & minore, & Oleæ folio capitulis compactis; cum Jaceæ flos, abit in pappos; harum autem flores non abeunt in pappos, ſed ſemina producunt ſolida, anguloſa longa: Ergò non ſunt verè Jaceæ, aliò relegentur. Nec aptiùs (meâ opinione) quàm ad Corymbiferas ſeu Corymboſas ïanthemas; referri poſſe ipſas Autumno. Ibidèm Col. II. Pl. V. Jaceæ, Oleæ folio affinis C. B. Pin, Dorynium Dioſcoridis

dis ſorte Ponæ Italicè & Cneoro albo *affi*-nis C. B. Pin. Dorycnium forte Dioſcoridis Ponæ Italicè pag. 463. Col. II. Pl. III. bis repetitur, eadem planta pro duabus diverſis, duobus diverſis locis, collocata.

Pag. 274, & 275. Tragopogon & Scorzonera: cum ſunt plantæ lacteſcentes pappoſæ, ad congeneres, Lactucam, Sonchum, Chondrillam, Hieracium, Piloſellam & dentem Leonis, referantur; quæ continentur à pag. 123. uſque ad Paginam 131. Ibidèm Caltha & Helenium Indicum, ſeu flos Solis, malè hic diſponuntur cum ad Corymboſas pertineant. Helenium vulgare ad pappoſas ſtellatas, Aſterem, Conyzam, Doronicum, Crithmum, Chryſanthemum, mandari debet.

Pag. 276. *Col. I. Pl. IX.* Caltha paluſtris capſulas efficit ſeminales, ut Ranunculus globoſus Aconiti in modum: & ſemina ibidẽ contenta, paria; non autem ut Caltha vulgaris.

Pag. 276. *Col. I. Pl. I.* Helenium ad non lacteſcentes, quæ tamen abeunt in pappos; ut ad Aſterem, Conyzam, & ejus Farinæ pappeſcentes, (ut ſupra dictum) debet relegari. Helenium Indicum, ſeu Chryſanthemum

themum peruvianum, ad Corymbiferas ſeu Corymboſas, tranſmitti debet, ergò malè confunduntur. Helenium Indicum ſeu flos Solis, cum Helenio vulgari, cujus flos abit in pappos; ut antè dictum.

Pag. 277. Perfoliata & Bupleurum, cum ſint Umbellæ perfectæ; quia flores, quinis petalis conſtantes, producunt, in ſummis radiis caulium, quorum ſingulis evanidis, ſuccedunt bina ſemina, ſimùl juncta; ergò debebant inter Umbellas, ſuæ claſſis plantas; collocari. Quid quæſo connexionis habent Perfoliata & Bupleurum, cum Centaurio minore.

Pag. 278. Gratiola cum ſit planta in udis pratis, vegetans: Tubiformem florem galeatæ, aut potiùs parvæ digitalis inſtàr gerens; cui tamen ſuccedit Theca ſeminalis, bicapſularis, digitalis, omninò inſtàr; continens multa minuta ſemina: hac ratione, aptiùs cum ſuæ ſortis plantis, collocanda; nempe digitale, Verbaſco, Blattaria, Scrophularia, Hyperico, &c.

Pag. 280. *Col. II. Pl. II.* Coris cœrulea maritima C. B. Pin. Coris Monſpelica Ad. Lob. florem fert galeatum, ſed capſulam habet

hãbet peculiarem, differentem, ab omnibus galeatarum, verticillatarum, capſulis ſeminalibus. Ideô amandari ad calcem galeatarum, cum Scutellaria & Euphraſia, debet.

Pag. 281. Polygonum latifolium C.B.Pin. & polygonum maritimum latifolium C. B. Pin. cum ſemina gerant triquetra; ad ſuæ ſortis ſtirpes relegentur Perſicariam, frumentum Saracenicum, &c.

Pag. 281, & 282. Polygonum & anthyllis, proptèr exilitatem, plantarum harum, difficilè admodùm eſt, ipſas ad idem genus reducere, quia magna eſt earum differentia, cum in flore, tum in ſemine.

Pag. 284. *Col. I. Pl. X.* Sedum minus fruticoſum, C. B. Pin. Sedum minimum arboreſcens vermiculatum Lob. malè ab Authoribus ad ſedum refertur, cum nil niſi folia, cum ſedo minore longifolio commune habeat; florem & ſemen, Kali gerit, dicique debet Kali fruteſcens omnium maximum.

Pag. 284. *Col. II. Pl. XI.* Sedum Alpinum luteum C.B Pin. Phyllon thelygonon dalechampii in dioſcoridem Lugd. hæc planta quamvis folia gerat vermiculata hirta,

ſedi

ſedi montani minoris, more, florem tamen fert tetrapetalum cui ſuccedit ſiſicula oblonga Paronychiæ, Alſines foliis inſtar. Eſt enim hujus ſpecies, foliis craſſis hirtis: non autem ſedi.

Pag. 285. *Col. II. Pl. VI.* Cotyledon ſtellata C. B. Pin. Aizoon peregrinum Camhæc planta nec florem, nec involucrum ſeminum aliter gerit, quàm Sedum ſeu Sempervivum minus. Ergò pertinet ad Sedi claſſem, eſt enim perfectè Sedum annuum ſtellatum.

Pag. 285. *Col. I. Pl. II.* Cotyledon media foliis ſubrotundis C. B. Pin. Cotyledon altera quarta Cluſ. hiſt. Cotyledon media foliis oblongis ſerratis C. B. Pin. & Cotyledon media ſubhirſuta floribus punctatis C. B. Pin. Cotyledon altera 5. Cluſ. hiſt. Hæ tres non ſunt verè Cotyledonis ſ. Umbilici veneris ſpecies, nec à flore, nec à capſulis ſeminalibus. Florem gerunt quinis petalis conſtantem: cui ſuccedit involucrum ſeminis bifurcatum: quemadmodùm, Saniicula montana, Saxifraga Cheſidonides, generunt. Suntque omnes plantæ, ejuſdem generis.

Pag. 287. *Col. II. Pl. VI.* Telephium repens folio non deciduo C. B. Pin. ſeptimum ſive legitimum Imperati Cluſ. hiſt. malè hìc collocatur pro Telephio, cum neutiquàm iis accedit, nec flore nec ſemine, eſt potiùs Polygoni ſpecies.

Pag. 287. Scorpioïdes Bupleuri folio, ejuſque ſpecies, item Portulacæ folio, cum florem edunt Papiliaceum, leguminum more; folliculos, per articulos (ſuo, quoque modo) incurvos, & in ſingulis articulationibus ſemina: quæ manſa, perfectè leguminum præbent ſaporem. Ideò ad leguminum claſſem, ſiliquis articulatis, præditam, remitto.

Pag. 288. Cepæa Matth. Sedi vera eſt ſpecies: Proindè Sedis jungenda, nec hìc per ſe collocanda. Ibidèm Crithmum maritimum majus & minus C. B. Pin. & maritimum ſpinoſum. Cum ſunt perfectæ Umbellæ, malè ab ipſis Umbellis diſjunguntur, & hìc collocantur: Item Crithmum maritimum flore Aſteris Attici, malè hìc jungitur duobus Crithmis Umbellis. Quarè amandetur ad Aſterum claſſem, & pappeſcentes ſtellatas plantas.

Pag.

Pag. 294. *Col. I, & II.* Scammonia Monſpeliaca foliis rotundioribus C. B. Pin. & Monſpeliacæ affinis, foliis acutioribus, quum flores & ſiliquas lanigeras; Apocyni in modum, gerant: ad Apocyni claſſem debent amandari: Item Scammonia Syriaca, cum ſit perfectè Convolvulus, ad Convolvuli ſeriem releganda.

Pag. 295. *Col. I. Pl. IV.* Convolvulus minor ſemine triangulo C. B. Pin. Helxine Ciſſampelos Lugd. malè inſeritur inter Convolvulos. Convolvulus enim eſt planta lacteſcens, florem fert Monopetalum unifolium ſeu integrum, capſulas ſubrotundas, ſemina anguloſa nigra, continentes. At hæc, nec lacteſcit, nec florem habet Monopetalum unifolium, quin & ſemen fert triangulare. Frumenti Saracenici, eſt vera ſpecies: Nil commune cum Convolvulo habet; niſi quod vicina quælibet amplectatur & ſcandat.

Pag. 295. Soldanella vel Braſſica maritima major & minor. Cum lacteſcunt & florem, & capſulam ſeminalem & ſemina, Convolvulorum habent: Convolvulis adſcribantur. Ibidèm Soldanella Alpina rotundifolia

tundifolia major & minor, nil cum Soldanella præcedente, habent affine: niſi folia (quod multis ſemper impoſuit) cum ſint ſiliquoſæ, aut Corniculatæ: alibi ad finem ſiliquoſarum, (impropriè ſic dictarum) collocandæ eſſent.

Pag. 303. & 304. Polygonatum, Lilium convallium, Monophyllum, Laurus Alexandrina: ad nervoſas bacciferas debent relegari.

Pag. 306. Hedera terreſtris vulgaris quum ſit galeata, verticillata, ad claſſem galeatarum verticillatarum debet remitti. Quare Bauhinus hîc collocavit poſt Hederam Corymboſam; nullam ego reperio congruentiam niſi in ſerpendo, ſimilitudinem: Ita infinitæ congeneres (etiamſi natura diverſæ) à naſcendi modò diſponerentur. Ibidem pag. 306. Cymbalaria, cum florem calcari donatum habet, qui compreſſus rictum edit, & capſulam duplici loculamento, conſtantem; ſepto membranaceo medio, idem diremente producit. Ut cæteræ omnes Linariæ; porro ſemina minuta ſuffuſca, ejuſdem formæ cum Linariæ ſemine ſunt. Igitur debet collocari inter Linarias:

Linarias: cum doctiſſimo Fabio Columna. Bis repetitur, hîc & pag. 253. ut ſupra, loco allegato, oſtendimus.

Pag. 306. Balſamina rotundifolia repens, ſivè mas C. B. Pin. Balſamina Cucumerina punicea Adv. Lob. mas Ger. cum ſit Cucumis, ad Cucumeris claſſem debet referi. Ibidem Balſamina fœmina C. B. Pin. fœmina Perſicæfolia Lob. & Balſamina lutea ſeu noli me tangere C B. Pin. cum utraque hæc planta, florem gerat calcari, ſeu Corniculo donatum, rictum habet peramplum paritèr utraque: utriuſque flori deciduo, ſuccedit ſiliqua, quinis loculamentis donata, quæ (maturò ſemine) tacta ſeſe contrahit, & ſemina vi ſpargit. Ad finem ſiliquoſarum, inter ſiliquoſas minùs propriè dictas collocari debent.

Pag. 309. Gramen parnaſſi, albo ſimplici, & pleno flore ad Pyrolam referantur. Ibidèm Saxifraga rotundifolia alba C. B. Pin. alba Chelidonides Lob. hæc ad Saniculam montanam, ut ante dictum, cum qua omninò quadrat: & (flore & ſemine) remittatur.

florem, capſulamque ſeminalem diverſimodè factam.

Ibidèm. Potentilla, cum florem & fructum, capſulamque ſeminalem, ſemen, granulatìm continentem: ad eundem modum, cum Fragaria ſterili, Pentaphyllo, Pentaphylloïde habet: Eiſdem jungi debet, nec per ſe hìc locari.

Pag. 322. Dentaria cum ſit planta tetrapetala & ſiliquoſa, malè hìc à Bauhino locatur, inter Caryophyllatam & Epimedium, à quibus toto Cœlo differt: ergò ſiliquoſis congeneribus ſuis, Leucoio, Heſperidi aſſociari debet. Vide pag. 201, 202.

Pag. 323, & 324. Pæonia & Staphis Agria, ad venenatas ſiliquiferas, Aconitum, Delphinium, &c. amandentur.

Pag. 324. Leontopetalon, cum ſiliquoſa ſit planta, tranſmittatur ad calcem ſiliquoſarum venenatarum. Ibidem Staphis Agria, cum capſulas ſeminales, accedentes ad Aconitum habet, cum ipſis, diſponatur verſus finem ejus ſectionis.

Pag. 326. *Col. I.* Quinquefolium fragiferum C. B. Pin. 5. fragiferum Cluſ. Quinquefolium fragifero affine C. B. Pin. Pentaphyllum

phyllum ſupinum Potentillæ facie Lobel. Icon. cum utraque hæc planta coſtas longas habet, quibus quinque interdùm plura, folia adhærent, è contrario, inter ſe foliorum exortu, naſcentia : uno extremum claudente. Potentillæ ſeu Argentinæ ritu : Ego (cum florem & ſemen, aliorum Pentaphyllorum more, perficiunt, folia autem aliter diſpoſita habent) Pentaphylloïdes nuncupari velim ; novum conſtituendo genus.

Pag. 326. *Col. II. Pl. VII.* Quinquefolium Lupini folio C. B. Pin. Pentaphyllon πεμπόφυλλον Thalii eſt planta ſiliquoſa non leguminoſa, proinde hinc averruncanda. Lupinus peregrinus Pentaphyllos C. B. Pin. Lupinus Arrabicus ſive Pentaphyllum peregrinum Ponæ Italicè, pag. 348. Colum. II. Pl. V. eadem planta duobus his ſuprâ dictis locis repetita, non poteſt pro Quinquefolio propriè loquendo accenſeri; Pentaphyllum eſſe fateor ſed Heteroclitum ; multò minùs eſt Lupinus, cum nec florem Papileaceum, nec ſiliquam leguminoſam fert.

Pag. 326. *Col. I. Pl. III.* Tormentilla Alpina folio Sericeo C. B. Pin. Candida Dalechampii hiſt. Lugd. cum omnibus ſuis

partibus

partibus eſſentialibus, flore & ſemine, Alchimillam æmulatur. Rectè concluditur (cum doctiſſimo Fabio Columna) eſſe Alchimilla.

Pag. 327. *Col. I. Pl. II.* Trifolium paluſtre Caſp. Bauh. Paludoſum Lob. cum Trifoliis leguminoſis nil habet affine; niſi quod folia gerat terna: ergò remittendum ad aquaticas plantas, tanquam Heteroclitum.

Pag. 327. Trifolium, Medica, Lotus, Melilotus quæ continentur à pag. 327. uſque ad pag. 331. ad finem leguminum pag. 353. debent collocari: cum omnia Trifolia ſint legumina Trifoliata, aut Trifolia leguminoſa, ex florum ſtructura & ſeminum ſapore, cuilibet guſtanti & propiùs conſideranti innoteſcet. Nos prolixiùs alibi (de horum Sympathia ſeu Identitate, quoad notam Genericam & eſſentialem) tractabimus.

Pag. 328. *Col. I. Pl. VI.* Trifolium montanum Lupulinum III. in Prodr. non debet disjungi à Lupulino vulgari, quamvis montanum: Nam locus non debet conſiderari ubi ſpicæ ſeu capitis ſimilitudo datur.

Pag. 328. *Col. I. Pl. I.* Trifolium pratenſe luteum, capitulo Breviori C. B. Pin.

Meleloti

Meleloti exigua vel minor ſpecies. Thal. floret ut Melilotus ſpicatìm, ſemina rugoſa dura Meliloti inſtàr, producit; rectè igitur pro Meliloto habenda; cum Trago.

Pag. 329. *Col. I. Pl. X.* Trifolium album anguſtifolium, floribus veluti in capitulum congeſtis C. B. Pin. Dorycnium Monſpelienſium Adv. Lob. cum hæc planta ſit Lotus ex appoſitione foliorum, nempè terna habet in extremo pediculo, bina ad appoſitionem Caulis. Ut cernere licet, quemadmodùm, in aliis, qua nota diſtinguitur Lotus. Addenda eſſet Loto Lybicæ, Dalechampii hiſt. Lugd. pag. 332. Col. II. Pl. X. & Loto Pentaphyllo ſiliquoſo villoſo C. B. Pin. Loto Hæmorrhoïdali majori ſeu Trifolio Hæmorrhoïdali majori Park. immediatè ante Pl. IX. in Col. I. paginæ 232.

Pag. 330. *Col. I.* Trifolium acetoſum, etiamque Hepaticum, malè interjiciuntur hîc inter Trifolia leguminoſa (quia nil ſimile niſi folia, habent terna communia, cum Trifolio leguminoſo) videlicèt inter Trifolium & Melilotum.

Pag. 331. *Col. II. Pl. VI.* Lotus hortenſis odora C. B. Pin. Melilotus vera Tab. ſingularis

ſingularis Alpini, Ponæ: eſt vera & genuina Meliloti ſpecies, à ſpicæ florum, & ſeminum diſpoſitione, juncto odore: Meliloto, inter Trifolia leguminoſa, ſemper debito.

Pag. 332. *Col. I. Pl. VI.* Lotus Pentaphyllos veſicaria C. B. Pin. Trifolium Halicacabum ſive veſicarium Cam. Omnis Lotus conſtat quinis foliis, tribus in extremo, pediculi; & duobus parvis, juxta appoſitionem, ad caulem. Hujus foliorum nervus, medius ornatur, binis terniſve foliolis ex adverſo ſitis, magno uno & longo extremum, claudente, Anthyllidis, vulgaris etiam ritu. Ergò aptius Loto affinis, Anthyllis Bætica veſicaria dicenda; quam Lotus, propter rationem antè dictam.

Pag. 333. *Col. I. Pl. XII, & XIII.* Rubia latifolia ſpicata & anguſtifolia ſpicata: cum Rubiæ propriè dictæ, nota Generica ſit, baccas aut ſemina rotunda, binatìm teſticulorum inſtár; ferre. Legentium judicium, penes ſit fides; num ſpicata major & minor (niſi à diſpoſitione foliorum ſtellatìm radiantium caulem) Rubiæ nomen mereantur.

Pag. 336. Ruta, Thalictrum malè inter ſtellatas

ſtellatas (à foliis) Gallium, cruciatam & legumina diſponuntur. Cùm Ruta, aptiùs collocabitur poſt bicapſulares cum, ſit Tetralobos ſeu Quadricapſularis. Thalictrum amandetur, ad Umbellarum finem, inter alias affines: propter ſimilitudinem foliorum cum quamplurimis, Umbellarum, cœterâ nil habent affine quoad partes eſſentiales, flores, & ſemina, binatim ſemper juncta, quod contingit in omnibus & ſingulis Umbellis, cujuſcunque, ſint ſemina figuræ, ut clariùs patefiet in Tabulis noſtris cognationis, in diviſione Umbellarum.

Pag. 337. Faba malè recencetur, pro legumine Capreolis donata: ergò ad legumina, ſiliquas erectas habentia; nobiſcum referatur, Lupinum, Glycyrrhizam, Galegam.

Pag. 338. *Col. II.* Faba ſylveſtris fructu rotundo atro C.B. Pin. ſylveſtris Dod. nulla Fabarum, Capreolis donatur; hæc Capreolis ſcatet, proinde rectiùs vicia omnium maxima diceretur (mea opinione:) adde quod nec ſiliquam ſeu folliculum; nec fructum fert Fabæ in modum: ſed Viciæ omninò æmulum.

Pag.

Pag. 343. *Col. I. Pl. X.* Piſum cordatum Adv. Lob. Veſicarium fructu nigro, alba macula notato C. B. Pin. malè inter leguminа Papiliacea & ſiliquoſa ſeu folliculacea collocatur; cum,leguminibus,nec flore nec ſiliqua quadrat; quippè veſicam, nucis juglandis magnitudine trifariàm diviſam, terna in ſe continentem ſemina producit, rotunda nigra, cor album lateralitèr, referentia; nec Maſticata, leguminis habent ſaporem: collocetur poſt Convolvulos cum Naſturtiio Indico. Heteroclita etiam eſt planta, hoc ſufficit impræſentiarùm.

Pag. 348. *Col. II. Pl. V.* Lupinus peregrinus Pentaphyllos,bis repetitur. Vide ſuprâ pag. 326. Col. II. Pl. VII. Quinquefolium Lupini folio C. B. Pin.

Pag. 348. Lupinus malè illic diſponitur, cum debeat adjungi Fabæ; quippè quod ſiliquas habet erectas; per intervalla, tumidiuſculas ubi ſunt ſemina, depreſſas in interſtitiis. Ibidèm malè dirigitur,ſeu locatur, Fœnugræcum, cum ſit Trifolium,ergò amandetur ad Trifolium cujus eſt ſpecies.

Pag. 349. *Col. I. Pl. II.* Aſtragalus Monſpelianus Cluſ.hiſt. Securidaca lutea minor, Corniculis

Corniculis recurvis C. B. Pin. hæc eſt planta annua, florem fert Ochroleucum ſeu pallidè luteum. Secundus eſt Aſtragalus Monſpeſſulanus Johan. Bauh. purpureus, hiſt. Lugd. pag. 351. Col. II. Pl. V. an Aſtragalus parvis foliis Hedyſari, ſeu Securidacæ hiſt. Lugd. in Appendice. Hic, eſt valde reſtibilis, in multos annos durat: flore eſt elegantèr purpurocœruleo. Tertius eſt Aſtragalus erectus annuus nobis; Vicia Seſamacea Apula Col. Ornithopodio, affinis hirſuta, fructu ſtellato C. B. Pin. pag. 350. Col. II. Pl. V. Hæc flore eſt etiam pallidè luteo, ſiliquas fert triplò majores ſiliquis, priorum. Nota ſtellam leguminoſam & Viciam Seſamaceam Apulam Col. eſſe duo diſtincta & diviſa legumina, etiamſi pro uno & eodem à Bauhino ponantur. Quartus Aſtragalus eſt perennis, flore luteo pallido nobis: Glycyrrhiza ſylveſtris floribus luteo palleſcentibus C. B. Pin. pag. 552. Col. II. Pl. III. Aſtragalus Monſpeliacus Joh. Bauh. citatur à Caſp. Bauhino in Pinace, pag. 351. Col. II. Pl. V. Aſtragalus villoſus floribus globoſis C. B. Pin. purpureus hiſt. Lugd. non eſt idem, cum Aſtragalo Monſpelliano

Cluſ.

Cluſ. ex ſupra dictis. Hæc quatuor legumina, cum ſub eodem vexillo militent, ad idem & novum conſtituendum genus elegimus; ex longe diverſis (ut obſervare licet in Bauhino) locis coegimus, à forma ſeminum, Officula Seſamacea, articulis digitorum pedum, ſubſtrata, referentium. Legumina Seſamacea ſeu Aſtragaloïdea, ſiliqua Gemella: Seu breviùs, Aſtragalus ſiliqua Gemella dici poſſunt. Nec norim, in tota familia leguminum, folliculaceorum caterva: præter hæc quatuor, quæ ſiliquas triangulares curvas & inflexas, falcis inſtàr, duplice Serie junctas, duplici membranacea Tunica, duplicem dirimente ſeriem ſeminum ſeu fructuum. Hiſce addi poteſt, ſed alterius figuræ, Hedyſarum peregrinum, ſeu Securidaca peregrina Cluſ. hiſt. quatenùs etiam folliculum profert, Spadæ Piſcis, Roſtri inſtar, aut Serræ utrinque Dentatæ, ſimilem: duplicem etiam ſeminum Seriem, planorum fert. Sicque conſtituto novum genus.

1. Aſtragalus Monſpeſſulanus perennis, flore purpurocœruleo: nobis. Purpureus hiſt. Lugd. vide pag. 351. Col. I. Pl. V.

2. Aſtra-

2. Aſtragalus Monſpelianus annuus, flore luteo palleſcenti: nobis. Monſpelianus Cluſ. hiſt. vide pag. 349. Col. I. Pl. II.

3. Aſtragalus erectus annuus, flore luteo palleſcente: nobis. Vicia ſeſamacea Apula Col. vide pag. 350. Col. II. Pl. V.

4. Aſtragalus reptans perennis, flore Ochroleuco ſeu luteo-pallido: nobis. Glycyrrhiza ſylveſtris ſive Glaux vulgaris. Vide pag. 352. Col. II Pl. III.

5. Aſtragalus, ſiliquis Serræ ſimilibus: nobis. Securidaca peregrina Cluſ. hiſt. vide pag. 349 Col. I. Pl. III.

Proprium eſt & leguminoſæ illi plantæ (quæ Tragacantha dicitur) ſiliquas ſeu ſoliculos, ſemiunciales, inter folia & aculeos, (poſt flores à longo tempore deciduos) producere. Duplici Serie ſeminum & ideò prioribus quo ad ſiliquarum, & ſeminum inibi contentorum, ſimilitudinem: valdè accedit. Nec unquàm memini hujus Farinæ, ſeptimum legumen obſervaſſe.

Pag. 349. Securidaca vera: folliculum longè diverſum habet ab Aſtraguli Monſ-

peliani Cluſ. folliculo; nota, ab utrâque: ſecuridacam dumetorum, ſiliquis ſuis articulatis, multùm differe. Nos quemadmodum ſuperiore obſervatione novum conſtituimus genus Aſtragalorum, ab identitate ſeu ſimilitudine non ſolum florum, (qui in omnibus quaſi leguminibus ſunt pares) ſed & ſiliquarum Gemellarum. Et hìc idem intendimus legumina ſiliquis articulatis, longis rotundis, ſtriatis, prædita, eodem revocare & communi nomine Baptizare, etiamſi ſub diverſis, apud Authores nominibus, diverſis in locis reperiantur.

1. Securidaca dumetorum major flore vario C. B. Pin. vide pag. 349. Col. I. Pl. IV.
2. Securidaca Enneaphyllos, ſiliquis in ſummitate: nobis. Lotus malè Dalechampio; nec benè ferrum equinum. C. B. Pin. ibidem Col. II. Pl. IV. cum ſequenti.
3. Securidaca Valentina: nobis. Polygala Valentina; 1. Cluſ. vid. 349. Col. II. Pl. II.
4. Securidaca Scorpioïdes major: nobis. colutea Scorpioïdes Ad. Lob. vide pag. 397. Col. I. Pl. II.

Omnes

Digitized by Google

Omnes hæ plantæ (quamvis ab Authoribus variis in locis ut à Bauhino variè denominentur) leguminoſæ ſunt, habentque pro nota eſſentiali & Generica, hanc, quòd omnium & ſingularum, ſub ipſarum capitulis contentarum ſiliquæ, exſiccatæ, per articulos rumpuntur, in ſingulis interſtitiis, ſingula ſemina oblonga, Cylindriaca continentur.

Pag. 350. Tribulus terreſtris, malè hìc inter legumina propriè dicta collocatur, cum nec florem, nec ſiliquam, nec ſemen, habet conformia, ipſis. Ideò propter diſpoſitionem foliorum, Rachi ſeu nervo medio adhærentium, uno extremum Claudente, (leguminum multorum more) ipſis adjungatur, Tribulus terreſtris, ſub finem.

Pag. 350. *Col. II. Pl. V.* Stella leguminoſa Lob. non eſt eadem cum Vicia Seſamacea Apula Col. ut cernere licet plantas creſcentes: imò & Icones Cluſ. & Columnæ differentiam facilè detegunt.

Pag. 360. Malè collocavit Bauhinus, Herbam minoſam poſt Polypodium, inter idem & Muſcum, cum ſit legumen, ad legumina eſſet referenda.

Pag. 376. Cardui cum ſunt plantæ capi-

tatæ perfectæ, quarum flores abeunt in pappos, Spinis ut plurimùm armantur. Ergò malè locantur poſt Muſcum, Fucum & Fungum (imperfectas plantas;) debebant collocari immediatè poſt capitatas pappaſcentes, non Spinoſas ut Jaceam, Stœben, Cyanum, Centaurium majus, &c. Sola hæc eſt differentia, quòd maxima pars Carduorum ſit Spinis armata, paſsim, exceptis quibuſdam. Quin et ſunt quædam genera Carduorum, quorum capita non abeunt in pappos, ſed ſolida efficiunt ſemina; & idcirco ſunt Cardui minùs propriè dicti. Ut clariùs poſteà innoteſcet.

Pag. 382. Bauhinus, inter Carduos Sphærocephalos (ſaltèm in eâdem ſectione) Collocat Carduum Eriocephalum; ſed Carduus Sphærocephalus non eſt Carduus propriè dictus (ut clariſſimè docebimus, in doctrinâ de Carduis, in Tabulis noſtris cognationis) ergò malè, his adjungitur Carduus Eriocephalus (cum ſit perfectè Carduus) Carduo Sphærocephalo, ejuſq; generi, quia ei deeſt ſecunda conditio ad conſtituendum Carduum propriè dictum.

Pag. 383. Acanthus ſativus & aculeatus non

non ſunt Cardui, nec à flore nec à ſemine nec ab aliarum partium, ſuarum diſpoſitione, alterutèr eſt Carduus. Ideò malè in medio Carduorum diſponuntur : ſubſtitui poſſunt ad finem Carduorum, quia Acanthus aculeatus eſt ſpinoſus. Sunt enim à tota facie, florum, ſeminum & capſularum ſeminalium formâ & figurâ plantæ peculiares & heteroclitæ.

Pag. 386. *Col. II. Pl. IX.* Eryngium arvenſe foliis Serræ ſimilibus C. B. P. Eryngium quartum Dod. malè hic inter Eryngia diſponitur, cum ne vel minimam notulam fert ejuſdem. Perfecta eſt umbellifera planta, & rectè ab Oculatiſſimo Dalechampio in hiſtoria ſua Lugd. Ammi quorundam nuncupatur. Nota inſuper Eryngium, omneſque ejus ſpecies non eſſe Carduos propriè dictos, quia eorum flores non abeunt in pappos, quæ eſt ſecunda conditio requiſita ad Carduos propriè dictos conſtituendos. Ut prolixiùs poſteâ ſignificabimus.

Pag. 386. Camphorata malè inter Ericam & Sabinam collocatur, cum ſit herba, perennis, una in agris Galliæ Narbonenſis & Gal-

lo-provinciæ frequens reperitur. Altera annua eſt,& paſsìm in Vinetis cernitur.

Pag. 387. *Col. I. Pl. VI.* Carduus Stellatus luteus Cyani fŏlĭis C. B. Pin. Spina Solſtitialis Dod. confundit hìc Bauhinus, Spinam Solſtitialem Dodonæi, cum Carduo Stellato minore Col. qui,& Spina Solſtitialis mitior Apula, eidem nuncupatur, & planè eſt omninò diverſa à priore, quoàd alas magìs undulatas, capita habet minora, & ſemina pappoſa paritèr minora, utrumque colui, ad aliquot annos.

Pag. 338. Tragacantha eſt planta leguminoſa: ſiliqua ſeu folliculo brevi Gemella prædita, duplicem Seriem ſeminum, continente,proindè ad legumina Aſtragaloïdea, ſeu Aſtragalos ſiliquâ Gemellâ, referri debet. Vide Aſtragalos, ſuprâ; non autem ad Carduos pertinet, nihil enim conforme habet cum ipſis: Summitates Ramorum ſeu Nervorum ſuſtinentium, multa folia hinc indè è Regione diſpoſita: Exſiccatæ, pungunt. Quaproptèr Bauhinus hìc collocavit, non video: remittatur ad ſuas congeneres plantas, (legumina videlicèt Seſamoïdea .ſ. Aſtragalos ſiliquâ Gemellâ.) novum genus facientes,

ſacientes, ex nobis. Ibidèm Col. II. Pl. II. Poterio affinis ſpinoſa foliis Pimpinellæ C. B. Pin. Pimpinella ſpinoſa Cam. χαμαῖον Anguillaræ Poterium Dalechamp. in hiſt. Cluſ. Hæc eſt vera Pimpinella, ut cuilibet intuenti cernere licet, ex diſpoſitione foliorum, modo florendi; ſemina globatìm rotundà fert. Ejus Nervi exſiccati pungunt: non tamen Carduis debet aſſociari, ſed cum Pimpinellâ diſponi.

Ibidèm. Anonis haud appoſitè adjungitur Carduis, propter Nervos ad extremum paritèr exſiccatos (vulgaris ſaltem Anonidis) pungentes. Sed cum Anonis ſit Trifolium, ab omnibus partibus eſſentialibus, debet Trifoliis annumerari, ut clariùs demonſtrabimus: cum Trifoliorum diverſa ſubalterna genera, methodicè per Tabulas cognationis exhibuerimus.

Pag. 389. Cytiſus Maranthæ Lugd. incanus ſiliquis falcatis C. B. Pin. Cytiſus omnis vulgò ſic dictus, ſiliquas profert longas, compreſſas, planas, aut hirſutas, ſeu hirſutie pubeſcentes, continentes in ſe ſemina compreſſa tumidiuſcula: Geniſtæ vulgaris penè ritu. Atqui hic ſiliquam producit parvam,

curvam ſeu falcatam, quæ ex Libro naturæ, propiùs accedit, ad Medicam falcatam: quod me impellit ad judicandum faciliùs & certiùs, Medicæ ſtuteſcentis eſſe ſpeciem, quam Cytiſi, utraque & Cytiſus & Medica eſt Trifolium; ergò cui magis accedit in ſiliquis ei eſt magis affinis ex doctrina noſtra. Et ſi, Medica, frutex incana perpetua fronde virens appellaretur: per me non ſtabit. Judicent; qui hæc legerint, an rationi conſentaneum ſit, quod dico.

Cytiſus incanus Creticus C. B. Pin. Ebenus Cretica Proſp. Alpin. exot. Ebenus ſecundus floribus purpureis Joh. Bauh. à quo figuratur. Ex antè dictis de Cytiſo Maranthæ. Hæc planta non eſt Cytiſus: ſed ſpecies elegantiſſima Lagopi maximi; & poteſt dici Lagopus Creticus fruteſcens, folio candicante. Semen collêgi apud Dominum Morinum natu minimum, atque ſatum ſemine, ex Italia ad me miſſo è terra pullulavit, ſed rigore hyemis ſequentis, periit; vidi ad aliquot annos, apud Dominum Morinum natu minimum, habitantem in ſuburbio Divi Germani.

Pag. 520. *Col. II. in appendice* C. Bauh. *ad Pin.* Doronicum Americanum laciniato folio C. B. Pinacis appendice: Cum omnis Doronici (propriè dicti) flos abit in pappos. Quæ eſt nota Generica & eſſentialis Doronici. Hujus autem plantæ ſemina ſunt ſolida, quadrangula Chryſanthemi adinſtar. Quarè potiore jure, Chryſanthemum Americanum majus, laciniatis foliis appellabitur. Multò minùs accedit ad Aconiti claſſem: quippè Aconitum florem fert Cucullatum, cui ſuccedunt capſulæ plurimæ membranaceæ: minuta ſemina, aſpera, continentes, ideò non poteſt dici, Aconitum Helianthemum Canadenſe: ut volunt alii.

Animad-

Digitized by Google

Animadverſiones IN *Tres TOMOS Hiſtoriæ* UNIVERSALIS PLANTARUM, *JOHANNIS BAUHINI.*

Animadverſiones in Tomum primum, Hiſtoriæ univerſalis Plantarum, Johannis Bauhini.

Pag. 374. *Partis Secundæ.* CYTISI facie, Alyſſum fruticans Lobel. quorundam. Malè inter Cytiſos collocatur à Johanne Bauhino: cum ſit ſiliculoſa planta, non verò leguminoſa. Ibidèm: Trifolium fruticans

fruticans: malè collocatur inter leguminoſas plantas, cum ſit Baccifera planta Trifolia: ad Jaſminum referri debet.

Pag. 520. *Partis primæ.* Vitis Idæa anguloſa deſcribitur, appingitur autem, Echii aut Anchuſæ Icon.

Pag. 460. *Part. pr.* Polygonum marinum anguſtifolium Dalechampii; malè inter Arbores collocatur, cum ſit ſuffrutex tantùm.

Pag. 553. *Part. pr.* Sambucus Roſea deſcribitur; apponitur autem Sambuci aquaticæ Icon. eâdem figurâ, ad *pag.* 552. tranſpoſitâ, & Roſeæ, figurâ ad *pag.* 553. tranſlatâ, tolletur lapſus.

Pag. 598. *Part. pr.* Dorycnium Imperati & Alypum Monſpelianum, malè collocantur hìc, cum ſint tantùm plantæ (ut & plurimæ aliæ ſuffrutices illìc collocatæ) inter Arbores.

Animad-

Animadverſiones in Tomum Secundum Hiſtoriæ univerſalis Plantarum; Johannis Bauhini.

Pag. 12. CIstus Ledum IX. Cluſ. & Ledum Thymi foliis Cluſ. nec à flore nec à capſulâ ſeminali aut ſemine, ſunt habendi Ciſti.

Pag. 23. Roſmarinus ſylveſtris quorundam; malè ſic nominatur, cum à flore & capſulâ ſeminali, ſit perfectus Ciſtus. Idem dicendum de duobus ſequentibus *pag.* 24, & 25. nil enim habent eſſentialitèr, cum Roremarino, nec in flore nec in capſulâ ſeminali, commune. Ibidèm: Libanotidis ſpecies Rondeletii, eſt Anchuſa lignoſior Lobelii, quarè ad Anchuſam amandetur, nullâ habitâ ratione conformitatis in foliis.

Pag. 354. Lotus Corniculata glabra, & hirſuta minor, deſcribuntur: duæ itidèm apponuntur figuræ, quarum prior eſt vitioſa, (quippè Medicam cochleatam, hiſpidam refert) poſterior eſt Loti glabræ, vera figura.

Pag.

Pag. 391. Epimedium quorundam, Columnâ primâ, ejuſdem paginæ, deſcribitur: appingitur autem Anonis ſeu Reſta bovis: & Columnâ ſecundâ ejuſdem paginæ, Anonis deſcribitur. Appingitur Epimedium: hæ duæ figuræ transferantur, hujus eò, illius huc, tranſlatâ figurâ: tolletur lapſus.

Pag. 398. Penultimâ paginâ adverſa Libri XVII. Caryophyllata vulgaris flore luteo parvo deſcribitur: appingitur Hepatica Trifolia.

Nota à pag. 398. *Sedecim paginas ad finem iſtius Libri, eodem numero continuò notari.*

Pag. 459. Gramen repens cum paniculâ Graminis Mannæ deſcribitur. Securidacæ appingitur figura.

Pag. 446. Gramen verſicolor deſcribitur: & pag. 476. Gramen ſtriis pictum deſcribitur. Eadem figura, duabus diverſis deſcriptionibus, appingitur.

Pag. 491. Gramen majus aquaticum Lob. deſcribitur. Appingitur autem Potamogeton.

Pag. 508. Gramen Triglochin: & Hyacinthi parvi facie, Gramen Triglochin. Uni-

nius

nius ejuſdemque plantæ, Graminis Triglochin videlicèt florentis, & ſemen ferentis, figura: pro duabus, toto Cœlo differentibus plantis, ponitur; duabus diverſis aſſignatis deſcriptionibus.

Pag. 590. Leucoii Hexaphylli, duæ apponuntur Icones: ultima vitiata eſt, loco floris appingitur bulbus.

Pag. 611. Deſcribitur Paucratium Monſpeſſulanum; appingitur Phalangium non ramoſum.

Pag. 803. Campanula Lychnidea: Non eſt Lychnis, quia non habet florem quinis petalis conflatum; ſed integrum ſeu Monopetalum, ut Convolvulus. Et ſi lacteſcit, eſt Convolvulus, mea opinione, nec florem producit Campanulæ in modum, quia, omnis Campanulæ .ſ. Trachelii Campanulati flos, in margine eſt quinquepartitus.

Pag. 810. Deſcribitur Rapunculus flore globoſo purpureo. Appingitur Campanula minor rotundifolia Alpina Bauh. in Pin.

Pag. 814, & 815. Jaſminum Mexicanum ſive flos Mexicanus multis. Malè collocatur poſt bicapſulares plantas: Cum ſit planta ſui generis: ſive Heteroclita.

Pag.

Digitized by Google

Pag. 816. Campanula Serpilli folio *Joh.* Bauh. malè etiam hìc collocatur, poſt digitalem ſeu bicapſulares, quum cum Campanula *p.* 796, & 797. in initio Libri XX^mi. debet poni.

Pag. 817. Campanula Linifolia lutea: cum in hallucinationibus Caſp. Bauh in Pin. à nobis exhibitis, ſuprâ, clarè ſatis docuimus, eſſe Linum; non poteſt eſſe Campanula, cum non lacteſcit, nec capſulam habet ſeminalem, nec ſemen Campanulæ, ſed Lini; ergò ut ſuperiùs diximus, ad Linum eſſet referenda hæc planta.

Ibidèm. Quo jure Soldanella montana, hìc collocatur à Joh. Bauh. non ſatìs comprehendo; cum ad calcem ſiliquoſarum aptiùs reduceretur.

Pag. 837. Turritis pulchra nova: non habet florem tetrapetalum, nec eſt ſiliquoſa planta: ergò malè inter ſiliquoſas collocatur.

Pag. 898, & 899. Telephium ſcorpioïdes, & ſcorpioïdes ex ſententiâ veterum; cum flores gerant Papiliaceos, ut cœtera legumina, ſeminaque articulatìm oblonga, perfectum leguminum ſaporem Maſticata, ſuggerentia:

Digitized by Google

ſuggerentia : Amandari debent ad finem plantarum ſiliquoſarum Papiliaceārum ; leguminum ſcilicét. Non autem ad finem ſiliquoſarum plantarum, tetrapetalarum. Ibidèm, deſcribitur Scorpioïdes montanum, appingitur Graminis alicujus figura. Ibidèm : Hypecoon ſiliquoſum deſcribitur. Papaveris malè picti, ſativi, appingitur figura.

Pag. 902. Coralloïdes minor bulbifera deſcribitur. Apponitur Rhodia Radix ex Trago.

Pag. 903. Lyſimachia lutea deſcribitur. Appingitur autem Naſturtii figura.

Pag. 927. Naſturtium vel Thaſpi maritimum deſcribitur. Graminis Sparti autem apponitur figura.

Pag. 939. Draba multis, flore albo deſcribitur. Duæ autem figuræ, diverſæ apponuntur. Drabæ videlicèt luteæ, ſiliquis ſtrictiſſimis Cam. & Naſturtii Babylonici Lob. quarum prior (quia ſiliquoſa) eſt rejicienda. Altera (quia ſiliculoſa) eſt vera & genuina.

Pag. 941. Dentillaria Rondeletii ; cum non ſit planta ſiliculoſa (quamvis cauſtica) alibi debet poni.

Pag. 982. Melochia, ſive Corchorus; cum ſit planta ſiliquoſa, Heteroclita: Ad finem plantarum ſiliquoſarum tetrapetalarum, debet poni. Non immediatè ante Lapathum propter uſum, quia eſt Olus Judaicum.

Pag. 984. Lapathum vulgare, folio obtuſo deſcribitur; appingitur Lapathi minimi Tab. figura ſeu Icon.

Animadverſiones in Tomum Tertium Hiſtoriæ univerſalis Plantarum: Johannis Bauhini.

Pars Prima.

Pag. 6. SCABIOSA peregrina rubra, quibuſdam Indica deſcribitur: At figuratur Circium.

Pag. 12. Scabioſa globularis, quam ovinam vocant: Hæc inter Scabioſas non debet locari; cum omnes conditiones, Rapunculo eſſentiales poſſideat (nempè lacteſcit, florem

florem habet Corniculatum, multorum aliorum Rapunculorum more: ſemenque producit ſubruſum, pellucidum) ad Rapunculorum claſſem, releganda eſſet hæc planta.

Pag. 13. Aphyllantes ſeu Globularia Bellidi ſimilis, & Aphyllanti affinis, Scabioſa quinta Cluſ. Bellidi affinis. Hæ duæ, cum non habent notam Genericam, Scabioſæ; ſed omninò accedunt Bellidum claſſi, aptiùs eò relegantur.

Pag. 25. Xeranthemum ſeu Ptarmica quorundam, & pag. 26. Xeranthemi fortè ſpecies & Catanance Dalechampii, flore Cyani, folio Coronopi, & pag. 27. Catanance affinis, Seſamoïdes parvum, flore magìs completo Cam. Omnes hæ plantæ, cum non pappeſcunt, non fuiſſent hìc collocandæ; Xeranthemi ſpecies ad Corymboſas, ſolida producentes ſemina amandentur. Catanance utraque, ad Cichorii genus (quippè lacteſcit, & Solida producit ſemina, non autem pappoſa) cujus, ex noſtrâ doctrinâ ſuperiùs traditâ in hallucinationibus Caſpari Bauhini, eſt vera & genuina ſpecies.

Pag. 30. Scabioſa ſquammata argentea. Malè à Johanne Bauhino, pro Scabioſa re-

cenſetur, cum Scabioſæ caput non abit in pappos, ſed in Solida ſemina. Hujus autem abit in pappos, ergò eſt aut Jaceæ at Stœbes ſpecies.

Pag. 71, & 72. Spina alba, quibuſdam, capite Echinato & Echinopus minor, vulgò Authoribus, Cardui Sphærocephali dicuntur. Itèm *pag.* 73, 74, & 75. Dipſaci ſpecies malè hìc collocantur, in medio Carduorum propriè ſic dictorum: quia non abeunt in pappos eorum capita, ut omnis Cardui perfecti caput: Suntque minùs propriè Cardui, ut & Eryngium, Scolymus: quod alibi clariùs deteximus. Idem eſto judicium de Spina lutea ſeu Scolymo, *pag.* 84. & de Eryngio *paginis* 85, 86, 87, & 88. Amandentur ad calcem Carduorum propriè dictorum. Nota inſupèr *pag.* 86. Eryngium cœruleum deſcribitur à Johan. Bauhino, & Plomos Lychnitis appingitur.

Pag. 106. Crithmum marinum tertium, Mattheolo flore luteo Buphthalmi; cum ſit planta Stellata pappeſcens, ad Stellatas pappeſcentes debet relegari, non autem hìc in medio plantarum Corymbiferarum .ſ. Corymboſarum collocari: quæ Solida anguloſa producunt ſemina. *Pag.*

Pag. 108. Helenium ſeu Enula Campana (per antedictam rationem) ad pappeſcentes congeneres debet amandari, non autem hìc locari; ut alibi in hallucinationibus Caſp. Bauh. obſervavimus.

Pag. 113. Bellis flore viridi deſcribitur. Thlaſpios, Myagri aut potiùs Erucæ peregrinæ Cluſ. appingitur figura ſeu Icon.

Pag. 144. Ageratum purpureum Dalechampii; cum non ſit corymboſa planta, malè hìc collocatur. Ad finem Euphraſiæ cum Hyſſopifolia Bauh. in Prodr. amandetur.

Pag. 151. Heliochryſos Tragi ſive Linaria tertia; cum hæc planta pappeſcit & florum tubulos, virgæ aureæ in modum producit; virgæ aureæ ſit ſpecies, Linariæ foliis. Sumendo denominationem à potiore & nobiliore parte, ſcilicèt florum ſeu florendi modo. Nec Linaria debet appellari, ut prolixè ſatìs in hallucinationibus ſuprâ exhibitis in capite Bauh. in Pin. de Linariâ docuimus.

Pag. 163. Piloſella Phlomodis exotica; cum lacteſcit & pappeſcit, debet ad ſuæ ſortis familiam relegari: nempè Chondril-

Digitized by Google

lam, Hieracium, Piloſellam, Dentem Leonis, &c. lacteſcentes, pappeſcentes.

Pag. 195. *& indè ad pag.* 200. Quâ ratione, ductus Joh. Bauhinus, ſubjungit Rutam Corymboſis, non capio; cum Rutæ omnis, involucrum ſit quadricapſulare. Multò magis ipſum miror, dum *pag.* 201. uſquè ad *pag.* 206. collocat Fumariam. Quin & in hujus Serie diſponit Ranunculi ſpeciem, Muſcatellinam', à denticulatione, (junctâ quâdam cavitate) radicis, nullâ habitâ ratione floris & ſeminum, nempè notæ eſſentialis. Deindè, quarè ſubjungit fini Libri XXVI. Melanthium, & conſolidam regalem, non video; cum ſint aliò à nobis aptiùs relegandæ.

Tomi tertii pars Secunda.

Pag. 6, & 7. ANETHUM & Anetho ſimilis planta, ſemine Lato laciniato; cùm ſint Umbellæ, ſemine membranaceo ſubrotundo prædítæ: Malè inter Fœniculum & ſequentes Umbellas (quippè ſemine ſtriato donatæ ſunt) collocantur.

Pag.

Pag. 8. Millefolium aquaticum, Fœniculo ſimile umbelliferum; non eſt perfecta Umbella. Millefolii aquatici ſpecies eſt: ergò ad aquaticas eſſet mandanda.

Pag. 9. Deſcribitur Daucus pratenſis Dalechamp. foliis Millefolii: appingitur Filipendulæ figura ſeu Icon.

Pag. 23. Cuminum ſylveſtre primum valdè odoratum, globuloſum, non eſt Umbella propriè dicta, (quia non habet bina ſemina ſimul juncta, quæ eſt nota Generica & eſſentialis toti Umbellarum Serei conveniens) ut in hallucinationibus Caſp. Bauhini, ſuprâ ſatìs dilucidè docuimus.

Pag. 33. Seſeli Maſſilienſe nuperorum folio aliquatenùs ſimili viſnagæ; cum ſemina habet membranacea, oblonga; ad ſui generis Umbellas (non autem inter Umbellas ſemine ſtriato tumidiuſculo ditatas) referri debet. ut clariùs in univerſali noſtrâ Umbellarum diſtributione deducemus.

Pag. 36. Peucedanum malè itidèm, inter Umbellas, ſemine ſtriato tumidiuſculo donatas, locatur: ex dictis ſuprâ. Deniquè adeò confusè Umbellas diſponit Joh. Bauhinus: ut impoſſibile ſit ipſas corrigere

perfectè, quoàd Tabulas noſtras cognationis, ſeu affinitatis exhibuerimus.

Pag. 40, *&* 41. Libanotis Cachryophoros, quibuſdam floribus luteis & Libanotis Cachrys Hiſpanica; cum ſemen gerant fungoſum exteriùs ad nucis parvæ Myriſticæ magnitudinem, & figuram ovalem: ex quâ notâ peculiari ego creo novum genus Umbellarum; quod Cachrys ſeu Libanotis Cachryophera appellabitur. Ut clariùs eluceſcet, ex diviſione Umbellarũ, in decẽ claſſes diſtribuendarum; ſic à naturâ notatarum, ut apparebit in Tabulis noſtris cognationis.

Pag. 43. Nota Joh. Bauhinum à *pag.* 43. ad *pag.* 53. duo genera Umbellarum inadvertentèr confudiſſe, quorum unius ſemen eſt compreſſum, membranaceum, ſubrotundum ut ſerularum omnium. Alterius ſemen eſt alâ foliaceâ membranaceâ cinctum: ut Thapſiæ ejuſque ſpecierum omnium.

Pag. 64. Paſtinacæ ſativæ, ſive Carotæ rubræ, luteæ & albæ ultima Icon. ex tribus; non indigitat Paſtinacam tenuifoliam; Paſtinacam potiùs latifoliam, ſeu Elaphoboſcum Matth. repreſentat. Ideò malè pro Paſtinaca tenuifolia, hìc ponitur: quarè

ad

ad *pag.* 151. debet referri; ubi eadem Icon. cum alterâ ejuſdem ſpeciei conſpicitur.

Pag. 74. Percipier Anglorum quibuſdam; cum ſit Alchimillæ ſpecies (ut in hallucinationibus Caſp. Bauhin. clarè ſatìs, ſuprâ docuimus) non debet inter Umbellas collocari.

Pag. 79. Gyngidium Matth. cum ſit ſemine minore ſtriato, foliis tenuiſſimè diſſectis, præditum. Malè à Joh. Bauhino inter Umbellas roſtratas Scandicem, Myrridem; & Umbellas aculeatas, Caucalidem ſcilicèt ſeu Lappulam (ut cum Johanne Bauhino loquar) diſponitur.

Pag. 83. Caucalis Hiſpanica ſemine rugoſo, & ab illâ paginâ prædictâ, uſquè ad *pag.* 86. Umbellæ omnes illic contentæ: (cum nota Caucalidis ſeu Lappulæ: eſt, ſemine aculeato, ſeu hiſpido donari, ut prolixiùs alibi trademus, ex Libro naturæ inſtructi) ſemine foliaceo, ſeu membranaceo ſubrotundo ditantur, ſub eodem vexillo cum congeneribus ſuis debent Militare. Videlicèt Sphondylio, Paſtinaca ſativa latifolia, Panace peregrino & cœteris (id genus) aliis cognatis, à naturâ ſignatis: Ut magìs

magìs dilucidè offeremus, in Tabulis affinitatis noſtris.

Pag. 91. Coriandrum: Coriandrum minus odorum, & Coriandrum ſylveſtre; cum genus Umbellarum per ſe conſtituunt, nempè ſemine rotundo teſticulato præditum: Malè hìc collocantur cum per ſe debent locari. Nota Umbellas ſequentes uſque ad *paginam* 113. eſſe omnes ſemine ſtriato minore præditas; quæ non debebant disjungi à conſortibus præcedentibus, ſemine itidèm ſtriato donatis.

Pag. 113, *uſque ad pag.* 120. Malè Sanguiſorba ejuſque ſpecies hìc collocantur; cum non ſint Umbellæ propriè dictæ, ſed plantæ ſpicatæ, aliquam in foliorum, conformationem cum Pimpinellâ, Saxifragâ umbellata habent. Ideò rejiciendæ eſſent ad calcem Umbellarum propriè ſic dictarum (propter ſimilitudinem foliorum) cum Filipendulâ, Barbâ Capræ, Valerianâ Græcâ, Thalictro, &c. ejuſdem Farinæ. Omninò, non fuerunt diſponendæ à Joh. Bauhino inter Umbellas ſemine minore ſtriato ditatas.

Pag. 137. Laſerpitium, proprium conſtituit genus Umbellarum: Quippè ejus ſemen, plu-

ribus

ribus alis, foliaceis ſeu membranaceis, alæ molendinæ fluviatilis inſtàr, (quandò binatìm jungitur ante exſiccationem) conſtat. Quâ notâ eſſentiali & Generica, indigitatur peculiare hoc genus Umbellarum.

Pag. 149, 150, & 151. Paſtinaca ſativa latifolia: cum ſuæ ſortis Umbellis lobatis, ſemine foliaceo, compreſſo, ſubrotundo prædìtis (ut ſuprâ dixi) diſponi debuit. Non inter Umbellas ſemine ſtriato minore ornatas; ut Siſarum poſt ſe, & Leviſticum ante ſe.

Pag. 164, & 165. Libanotis Theophraſti, Seſeli Æthiopicum Mattheolo; & Libanotidi Theophraſti quorundam valdè ſimilis; Coſtus amara dicta ſemine criſpo: Hæ duæ Umbellæ, ex Libro naturæ, Laſerpitii, gerunt teſſeram (id eſt ſemina ſunt ejuſdem formæ) ergò jure merito, à nobis recenſebuntur Laſerpitia.

Pag. 183. Phellandrium, vel Cicutaria aquatica quorundam; non eſt Umbella propriè loquendo: Quia omnes Umbellæ, habent bina ſemina ſimùl juncta, cujuſcunque figuræ aut formæ ſit ſemen; atque propter defectum hujus præcipuæ, univerſalis

ſalis & eſſentialis conditionis; minùs propriè dici debet Umbella: In cœteris omnibus, eſt Umbellæ æmula, hæc Cicutaria aquatica.

Pag. 186. Thapſia latifolia & Carotæ facie, ſemen gerunt ſtriatum, oblongum; alâ membranaceâ cinctum; Quâ notâ peculiari diſtinguitur, ab omnibus ſubalternis generibus Umbellarum: & ideò nos ſic diſtinguemus in Tabulis (exhibendo Genericam & eſſentialem hanc notam;) noſtris diſtributivis, Umbellarum: confusè omninò ab Authoribus, hucuſque exhibitarum. Nota inſupèr *pag.* 50. Thapſiæ majoris ſeu Turbith Garganici, ſemen latiſſimum benè repræſentari: Sed vitiosè Sileris aut Cachryos, Peucedani foliis, appingitur folium.

Pag. 189. Filipendula deſcribitur; Dauci autem pratenſis apponitur figura.

Pag. 193. Oenanthe Myconi: Malè hìc, inter Umbelliferas, diſponitur; cum ſit Ranunculi, Aſphodeli radice ſpecies; ut ex flore & ſemine, totaque facie apparet. Quarè ad Librum XXX. inter congeneres relegenda eſt hæc planta.

Pag.

Digitized by Google

Pag. 282, 285, & 286. *usque ad pag.* 287, 288, 289, 290. Sive ſub Teucrio, ſive Chamædre, omnes hæ plantæ nominentur, ſunt veræ & genuinæ Veronicæ ſpecies; quippè ſunt tetrapetalæ ſiliculoſæ, ſepto medio membranaceo, diſcriminante duplicem Seriem ſeminum, ut videre licet in ſiliculoſis tetrapetalis; ut Burſâ paſtoris, Thlaſpi, Cochlearia, Naſturtio hortenſi, Drabâ ſiliculoſa, Lepidio, Iberide, &c.

Pag. 290. Chamædrys Alpina, flore Fragariæ albo deſcribitur: Hormini appingitur figura.

Pag. 291. Scutellaria Teucrii facie & *pag.* 293. Scorodotis, ſeu Scordium folio Salviæ. Nota has verticillatas & Galeatas, ad finem verticillatarum & galeatarum rejici debere. Quia capſulas ſeminales non more aliarum, quinqueſariàm in margine diviſas gerunt; ſed Scutellæ in modum: undè novum genus creavimus ex pluribus hìnc illìnc, ex Authoribus variis collectum. Vide, & conſule hallucinationes Caſp. Bauh. à nobis exhibitas.

Pag. 307. Verbaſcum ſylveſtre Monſpelienſium flore luteo hiante; deſcribitur.

At

At umbelliſera appingitur planta, fortè Scandix.

Pag. 321. Deſcribitur Galeopſis, ſive Urtica iners, &c. & *pag.* ſequenti appingitur Balſaminæ Perſicæfoliæ (fortè) figura florentis.

Pag. 323, & 324. Locuſta herba, & Locuſta altera foliis ſerratis: Malè hìc committuntur, ad finem galeatarum, verticillatarum, cum quibus nil planè habent affine; remittantur ad *pag.* 212. & claudant Sectionem Umbellarum; etiamſi non ſint Umbellæ perfectè. Quia non habent bina ſemina ſimùl juncta, quæ eſt nota eſſentialis & Generica omnibus & ſingulis Umbellis, perfectè ſic dictis conveniens. Tamen florent quodammodò Umbellarum more & earundam folia, aliquatenùs Valerianæ Alpinæ accedunt.

Pag. 348. Flos Cuculi, flore multiplici purpureo, & altero albo deſcribitur. appingitur Alſines minimæ ſpecies.

Pag. 362. Alſine major, repens, perennis deſcribitur: Apponitur Polygalæ luteæ Icon.

Pag. 364. Alſine minima deſcribitur:

Appingitur

Appingitur odontidis flore pleno Icon. hujus figura ad *pag.* 348. illius ad *paginam* 364. transferatur, & tolletur lapsus.

Pag. 366. Alsinæ fœtidæ quodammodò similis laciniata; delineatur seu describitur: Phaseoli autem apponitur figura seu Icon.

Pag. 367. Alsine serrato folio hirsutiori, floribus & loculis Cauliculis adhærentibus; Alsine oblongo folio serrato, flore cœruleo. Et *pag.* 368. Alsine folio profundè secto, flore purpureo seu violaceo; Item Alsines genus, Fuchsio folio Hæderulæ hirsuto: Omnes hæ quatuor plantulæ; cum singulæ singulos edunt floseulos (cujuscunque coloris) quatuor petalis constantes, singulis earum floribus succedunt siliculæ bicapsulares, Veronicæ in modum. Ut satis clarè in hallucinationibus Casp. Bauhini, in Pin. nobis solummodò observatis; ostendimus. Inadvertentèr omnes quasi Authores, hucusque has; Alsines species constituêre. At pro Alsines speciebus, non sunt reputandæ; cum Alsines flos constat quinis petalis, ad medium bifidis; cui deciduo succedit capsula coni instar. Ergò intuendo Librum naturæ Veronicæ species sunt annuæ: Consule hallucinationes

lucinationes Caſp. Bauh. à nobis ſuprâ editas. Muta Alſines nomen, & denominando ipſas, Veronicam ; habes naturæ intentum.

Pag. 375. Polygoni; duæ appinguntur Icones, quarum poſterior eſt vitioſa.

Pag. 381. *uſque ad pag.* 385. In quibus omnibus continentur Androſæmum, Aſcyrum, Hypericum; quæ omnia malè illic conſtituuntur inter Polygonum, & Polygalam: Cum ſint bicapſulares, Pyramidales ſeu Coniformes (quoad involucra ſeminum) eſſent relegandæ ad digitalem, Verbaſcum, Blattariam, Gratiolam, &c. ut ſuprâ docuimus ; in hallucinationibus Caſp. Bauh.

Pag. 382. Hypericum Aſcyrum dictum; Caule quadrangulo deſcribitur: appingitur Ocymoïdes non ſpecioſum, ſeu *Lychnis* ſylveſtris, alba noctiflora Cam. hujus Icon. ad *pag.* 344. & illius ad *pag.* 382 tranſmittatur, & ſic tolletur error Joh. Bauh.

Pag. 386. Onobrychis lutea Daleehamp. deſcribitur; at appingitur Alſines majoris repentis, perennis, Icon. hujus Icon. ad *pag.* 362. illius, ad prædictam 386. transferatur; & inadvertentia Joh. Bauhini auferetur. Nota ibidèm inſuper Joh. Bauh. haud appoſitè

positè Polyganon pro Polygala scripsisse.

Pag. 410. Pulsatilla alba describitur. Appingitur autem alterius plantæ Icon.

Pag. 414. Ranunculus cum flore in medio folio, radice tuberosa; cum non producat semina Ranunculi in modum, sed Aconiti: Unde à siliquis Aconiti & flore Ranunculi, Aconitum Ranunculoïdes 1. appellamus in hallucinationibus Casp. Bauh. sicut per eandem rationem Ranunculus flore globoso quibusdam, trollus flos *pag.* 419. à siliquis longis membranaceis Aconitum Ranunculoïdes 2. appellamus: Consule hallucinationes Casp. Bauh. à nobis superiùs editas.

Pag. 421, & 422. Scrophularia maxima radice fibrosa, describitur; appingitur autem Sideritis Monspeliaca. Nota insuper, malè Scrophulariam inter galeatas verticillatas collocari, cum sit planta bicapsularis ut alibi docuimus.

Pag. 426. Sideritis Monspeliaca describitur; appingitur autem Scrophularia maxima radice fibrosa. Hujus Icon. ad *p.* 422. & illius ad *pag.* 426. transferatur, & corrigetur error.

Nota insuper, quæ stirpes continentur à

pag. 421. *Libri XXX. uſque ad pag.* 468. *ejuſdem Libri, debent referri ad Libros XXVIII, & XXIX. videlicèt, quæ continentur à pag.* 421. *ubi intuenti ſeſe offert Scrophularia vulgaris major, uſque ad pag.* 449. *ubi legitur Bangue; debent amandari ad Lib. XXVIII. pag.* 323. *ubi legitur Galeopſis. Quippè quod verticillatæ & galeatæ, verticillatis & galeatis eſſent addendæ Quæ autem continentur à pag.* 450. *ubi in conſpectum ſeſe dat Linum: uſque ad pag.* 467. *ubi legitur Reſeda: Debent referri ad Libri XXIX. pag.* 371. *ubi legitur Numularia.*

Pag. 434. Gratiola, malè à Joh. Bauhino (cum ſit bicapſularis Coniformis planta, ad Scrophulariam, Verbaſcum, Blattariam, &c. referenda foret) inter verticillatas & galeatas plantas diſponitur.

Pag. 446. Urtica minor annua, hujus Icon eſt vitioſa.

Pag. 455. Alſine verna glabra, malè hìc à Joh. Bauhino collocatur, inter Lini ſpecies: Quia eſt perfecta Alſine à flore, quinis petalis ad medium bifidis & capſulâ ſeminali, Alſinarum more, factis donata.

Pag. 456. Paſſerina Tragi, & *pag.* 455. Paſſerina

Paſſerina Lobelii neutra harum eſt Linaria, nec à flore, nec à ſemine: Nil habent commune cum Linaria niſi folia; ſunt plantæ ſui generis & Heteroclitæ.

Pag. 461. Linariæ ſimilis; duæ adferuntur in medium figuræ, quarum neutra eſt Linaria, prior refert Polygalam, Altera eſt Anonymos Cluſ. ſeu Heteroclita cujus folia quodammodò Linariæ foliis ſunt ſimilia. Idem dicendum venit, de Linaria belvidere.

Pag. 463. Antirrhinum vulgare, hujus tres apponuntur Icones: quarum, prior eſt Junci cum utriculis: (Caſp. Bauh. in Prodr.) Icon.

Pag. 468. Scrophularia minor Lib. XXX. debet amandari, ad *pag.* 420. ubi legitur Ranunculus. Poſt admiſſos alios ejuſdem generis, in appendice ad Tomum tertium, contentos.

Pag. 471. Caltha paluſtris flore multiplicato: Cum ſiliquas plures oblongas, membranaceas, ſemina in ſe continentes nigra; producit Caltha paluſtris: Benè mea opinione, ſubjungi debet Aconito flore Ranunculi in medio folio: quod continetur *pag.* 414.

Digitized by Google

Pag. 482, & 483. Chelidonia major & eadem folio laciniatomalè, collocantur hìc; cum ad ſiliquoſas tetrapetalas plantas debent referri. Quæ continentur in Tomi 2. Lib. XXI. tetrapetalæ ſiliquoſæ, tetrapetalis ſiliquoſis jungendæ ſunt: cum in Tomi 3. Libro XXX. nil habent affinitatis. Nec cum genere antecedente Ranunculo, ſcilicèt, nec cum genere ſequente aquilegia: Sed ſunt tetrapetalarum ſiliquoſarum veræ & genuinæ ſpecies; ut cuilibet conſideranti, cernere licet.

Pag. 484, & 485. Aquilegia cum ſit planta ſiliquis membranaceis plurimis, Aconiti & Delphinii adinſtàr, debuit iiſdem ſubjungi.

Pag. 486, & 487. Thalictrum ejuſque ſpecies; à ſimilitudine foliorum, in ipſorum diſpoſitione, cum plurimis Umbellis conveniunt. Idem contingit Filipendulæ Ulmariæ, ſeu Barbæ Capræ, aliiſque; jure accidentali, poſſunt ad finem Umbellarum diſponi: ut ſuperiùs clarè ſatis oſtenſum fuit.

Pag. 489. Leontopetalon cum ſit planta Heteroclita, ad finem Ranunculorum poteſt amandari.

Pæonia

Digitized by Google

Pæonia ejuſque ſpecies, adjungantur iis quæ plurimas ſiliquas membranaceas habent; ut Delphinio, Aconito, Aquilegiæ, Staphidi Agriæ, cœteriſque illius Farinæ: ut nos concluſimus.

Pag. 496. Primula veris, caulifera, prolifera, odorata deſcribitur. Appingitur confusè cum Primula veris floribus plenis ex ſingularibus pediculis.

Pag. 497. Primula veris floribus plenis ex ſingularibus pediculis, deſcribitur. At appingitur Umbellifera Primula veris confusè cum prædictâ.

Pag. 521. Gentiana folio convoluto, eſt à flore & capſulâ ſeminali etiamque ſemine Lychnidis (Saponariæ dictæ) ſpecies genuina; neutiquam eſt Gentiana. Ergò malè hìc cum Gentianæ ſpeciebus locatur; relegari debet ad Lychnidum claſſem. Saponariam nempè, eandem cum Caſp. Bauhino Saponariam concavam nuncupando.

Pag 524. Gentianæ ſpecies, Calathiana quibuſdam radice perpetua, ſive paluſtris deſcribitur: duæ autem Gentianæ, omninò diverſæ Icones apponuntur; prior vera, alterius eſt Gentianæ punctato flore Cluſ. Icon.

Digitized by Google

Pag. 533. Bifolium majus, ſive Ophrys major quibuſdam: Malè hic locatur, cum meliùs congruit cum Helleborine; cujus eſt multò magis æmula, quoàd omnes ſuas partes.

Pag. 538. Biſtorta, cum ſit ſemine triangulari donata; magis appoſitè diſponeretur poſt Lapathum acetoſam, &c.

Pag. 563. Tuſſilago, & à *pag.* 563. uſque ad *pag.* 569. in quibus continentur Tuſſilago, Petaſites, Cacalia; quæ omnes, cum pappeſcunt & non lacteſcunt, debent pappeſcentibus, non lacteſcentibus ſubjungi: Videlicèt Jacobææ, Senecioni, Gnaphalio, Stœchadi Citrinæ, &c.

Pag. 579. Bugloſſum vulgare, majus & minus, utrumque flore duplici: malè figuram exhibet Joh. Bauhinus Bugloſſi vulgaris, flore ſimplici.

Pag. 588. Echium tenuifolium deſcribitur. Appingitur Bugloſſi ſpecies, latiore folio: obſerva hoc diſcriminis intercedere inter Bugloſſum, ſeu Borraginem, & Echium ſive Lycopſin, Bugloſſi ſeu Borraginis flos conflatur ex quinis petalis acutangulis, cingentibus ad baſin fimbriam villoſam; in cujus

jus medio totidem prominent Apices conjunctim, aculeum referentes; sed Echii seu Lycopsios flos, est corniculi aut tubuli infra, oblongus, in superiore floris parte longius porrectus; in inferiore margine breviori utriusque (floribus discussis) in vasculis, tria vel quatuor semina conjunctim posita sunt, oblonga, rugosa: veluti semina Consolidæ majoris.

Pag. 590. Cynoglossa fortè Topiaria Plinio, seu Echium lappulatum quibuldam, describitur; sed perperàm appingitur Cynoglossa, folio virenti Joh. Bauhini.

Pag. 592. Lithospermum majus Dodonæi flore purpureo, semine Anchusæ describitur. Sed autem aliqua appingitur species; nempè Sedum Alpinum minimum foliis cinereis, perenne candido, descriptum in *pag.* 692. hujus Icon. ad *pag.* 692. illius autem Icon. ad *pag.* 596. transferatur, facilè emendabitur error.

Pag. 598. Cynoglossum vulgare malè exhibetur; quia Icon. est vitiosa.

Pag. 600. Cynoglossa folio virenti Joh. Bauhini describitur. Appingitur autem Cynoglossa fortè Topiaria Plinio, seu Echium

Bibliotheca Carolina

lappulatum quibuſdam, hujus Icon. *pag.* 600. contenta, ad *pag.* 590. amandetur: & illius huc transferatur, & ſic corrigetur lapſus.

Pag. 637. Buphthalmum Dodonæi aliis Conſiligo tenuifolia, cum ſit planta Corymboſa, ſemine Adonidis Emulo prædita, ad Corymboſas debet amandari.

Pag. 638. Sanicula fœmina quibuſdam, aliis helleborus niger; cum ſit planta umbellifer, aut clarè ſatis docuimus in hallucinationibus Caſp. Bauh. ad Umbellas eſſet releganda, cum Tabernemontano, eamque appellando Imperatoriam nigram.

Pag. 641. Staphis Agria, cum ſit planta gerens plures ſiliquas, membranaceas continentes ſemina dura, lucida, nigra; relegetur ad Aconitum, Delphinium, Aquilegiam, &c.

Pag. 643. Ricinus cum ſit planta tricoccos, & ſimili purgandi virtute prædita, cum Tithymalo, ejuſque ſpeciebus; debet ipſis ſubjungi.

Pag. 650, & 651. Thora folio Cyclaminis, & Thora Apula folio Alchimillæ; cum ſint Ranunculi, à flore & ſemine; non eſt dubitandum quin debeant relegari ad Ranunculorum claſſem.

Pag.

Digitized by Google

Pag. 660. Aconitum racemoſum, actæa quibuſdam; cum hæc planta ſit baccifera, Solano magìs accedit (mea opinione) quàm Aconito, ut ſuprâ innuimus.

Pag. 678. Portulaca hortenſis, latifolia deſcribitur. Appingitur autem Menianthes, ſeu Trifolium paludoſum.

Pag. 680. Sedum Echinatum vel ſtellatum flore albo deſcribitur; vitioſa appingitur Icon. fortè alicujus Polygoni.

Pag. 684. Cotyledor, altera olim Mattheolo; non eſt Cotyledonis propriè dictæ ſpecies; ſed florem fert Sedi aut Saniculæ montanæ, capſulamque paritèr in ſuperiore parte bifurcatam, in qua plurima continentur ſemina tenuiſſima.

Pag. 685. Cymbalaria floſculis purpuraſcentibus, neutiquàm huc pertinet, ſed debet jungi Linariæ familiæ. Cujus fert teſſeram ſeu notam Genericam: ut ſuprâ in hallucinationibus Caſp. Bauh. ſatis evidentèr à nobis deducitur.

Pag. 692. Sedum Alpinum minimum foliis cinereis, flore candido deſcribitur. Appingitur autem Lithoſpermon majus Dod. flore purp. ſemine Anchuſæ. Vide *pag.* 592.

fiat

fiat tranſpoſitio Iconum,& tolletur lapſus.

Pag. 710, & 711. Lunaria Botrytis ejuſque ſpecies, malè hic digeruntur; cum ſint perfectæ filices, inter filices collocentur.

Pag. 722. Spergulæ tres Icones,malè *illic* diſponuntur inter Rubiæ ſpecies. Quia Spergulæ, à flore & capſulâ ſeminali, ſunt Alſines perfectæ ſpecies. Ergò ad Alſines claſſem debent relegari; nullâ habitâ ratione foliorum, per intervalla ſtellatìm poſitorum.

Pag. 728. Drypis Theophraſti, cum ſit Carduus,malè hìc inter Aſparagum & Equiſetum digeritur; & à ſuæ ſortis familiâ diſjungitur: Ergò ad Carduorum ſeriem debet jure mandari.

Pag. 754. Trichomanes,ſive Polytrichum deſcribitur. Appingitur autem Cardui alicujus Icon.

Pag. 762. Phyllon, Thelygonon Dalechampio, cum ſiliculam habet bicapſularem, ſepto membranaceo, duplicem diſterminante ſeriem ſeminum; Paronychiæ muralis adinſtàr, aptè (meâ ſententiâ) Paronychia foliis craſſis, hirtis diceretur. Ibidèm Tridactylites Alpina: florem & capſulam

ſemina-

Digitized by Google

ſeminalem habet; ut Sedum Tridactylites tectorum: Ergò ad Sedi claſſem eſſet releganda, & non hìc collocanda hæc plantula.

Pag.868.in appendice ad Tomum tertium. Auricula urſi orbiculata, ſerrata coloris rubicundi deſcribitur. Appingitur autem vitioſa Icon: fruticis ſcilicèt alicujus, ramus.

Ad

Digitized by Google

Ad virum insigniſſimum, Doctiſſimum & vitæ integerrimum, Digniſſimumque Dom. *Johannem Fell*, in ſacroſanctâ Theologiâ Doctorem, Collegii ædis *Chriſti* Decanum & Univerſitatis *Oxonienſis* Procancellarium vigilantiſſimum S. P. D. Author.

Clariſſime Vir,

POST *reditum meum*, Oxoniâ Londinum, *amicorum quorundam Precibus ſolicitatus: ut præludia* Botanica *hæc Mea, typis committerem: quæ jam tandem prelo exceſſêre. Quæque viam, ad methodum noſtram novam (cujus tibi exemplar oſtendi) præſternunt: novam appello quia nemini hucuſque cognita: antiqua tamen quia à tertio die creationis mundi, eadem ſemper fuit, quæ hodie eſt. Reverende Vir, non ita pridèm (ut noſti) inviſebam ſplendida veſtra & magnifica ædificia; Collegia, Scholas, Bibliothecam,*

Bibliothecam, Theatrum, publicis actibus de-ſignatum, vix adhuc finitum. Liberalitate & ſumptibus, Reverendiſſimi in Chriſto Patris, Archiepiſcopi Cantuarienſis; Gilberti Shelden, *ejuſdémque Univerſitatis Cancellarii dignißimi; à fundamentis erectum. In Symbolum gratitudinis ſuæ, erga almam matrem ſuam: cui præſentem ingenii Culturam debet. Magnificum hoc Theatrum (omnium in* Europâ *facilè primum) ut & omnes graviores curæ Univerſitatis; vigilantiâ, labore & inſpectione tuâ, (nec auribus tuis jam do) diriguntur. Dum fabricam & ſtructuram omnium* (inquam) *horum publicorum ædificiorum, animo volvo: Nihil unquam apud exteras nationes, ipſis comparandum, audactèr aſſerere audeo. Sed dum numeroſam Doctißimorum, in omni literarum genere, hominum illic Congregationem penitiùs conſidero: Dumque omnes eâdem ſuſpendo bilance, Dialogum hunc, inter Socium* Londinenſis *Collegii Regii, & Me: nulli aptiùs, quam tibi inſcribere; ratum & ſtatutum habeo. Solutiones quarundam quæſtionum propoſitarum; inter prandendum, Domi tuæ, ſub idem tempus; invenies. Cum de methodo Authorum, in rem* Botanicam,

Botanicam, *hucusque scribentium*; *tum*, *de Plantis imperfectis agit*: *Estque hic totus ferè scopus totius Dialogi. Accipe igitur*, *hilari animo*, *hæc nostra alterna colloquia* Botanica. *Tu qui me benignè recepisti*, *omni humanitatis officio*, *me affeciſti*: *dum apud vos conversarer. Utinàm vobis aut vestris inservire possim commodis*: *Me invenietis paratissimum*; *Vale. Et ut diu vivas*, *Deum optimum*, *maximum precor*:

Vir Reverende

&

Doctissime,

Tuus obsequentissimus

&

Observantissimus servus,

ROBERTUS MORISON.

Digitized by Google

DIALOGUS

INTER

SOCIUM,

COLLEGII REGII

LONDINENSIS,

GRESHAM dicti:

ET

BOTANOGRAPHUM

REGIUM.

Socius. Botanographus.

Socius. FANDO ad meas pervenit aures, imo jam cum in memoriam revoco, ex tuo ore hauſi: Te excogitaſſe, aut ſaltem incidiſſe in methodum generalem omnium ſtirpium, per Tabulas quaſdam cogna-

cognationis, ſeu affinitatis; ſed quid ſibi velint hæ Tabulæ, nondùm aſſequor. Quarè velim, ut paucis privatìm inter nos de eiſdem conferamus. *Botanographus.* Methodum me obſervaſſe fateor: Eſtque, omnium, quæ unquam adhuc fuerunt exhibitæ: præſtantiſſimæ & certiſſimæ: eſt quippe à naturâ data, à me ſolummodò (citra jactantiam) primum obſervata, quamque publici juris facere brevi, in me ſuſcepi. Si Mœcenates faverint, noſtris conatibus & lucubrationibus per aliquot annos à nobis exantlatis. *Soc.* Volumina multa magna evolvi, de re Botanicâ: Sed methodum ſtirpium hucuſque deſidero; quæ multùm conduceret ad citiùs & certiùs vaſtum illum Oceanum, plantarum ſeu Stirpium addiſcendum. Inſupèr Calcaria adderet multis nobilibus, doctiſque hominibus, plantas è terrâ pullulantes avidiùs contemplari & divinum artificium ſummoperè mirari: ſed quid quæſo tu dicis, qui tres modios ſalis in Galliâ comediſti: de hiſtoriâ univerſali Plantarum, Lugdunenſi; à Dalechampio editas. *Botan.* Fateor me fœlicitèr in Galliâ vixiſſe multaque illìc evolviſſe volumina, cum

cum magna, tum parva, de arte (aut scientiâ si mavis) Botanicâ sub auspiciis serenissimi principis Gastonis Franciæ: qui in re Botanicâ fuit peritissimus, omnium scriptum, diligentissimus scrutator, & oculatissimus observator. Sed ut tibi satisfiat, redeo ad Gallum nostrum Botanicum, Dalechampium: fateor Dalechampium fuisse Medicum insignem, & perdoctum Botanicum. Quin & opera ejus, Botanicis nostrûm profuêre: sed quod utplurimùm mortalibus contingit, ipsæ præmaturâ morte abreptus, chartas suas & Icones à Mattheoli, Lobelii, Dodonæi, & Clusii hæredibus; suis sumptibus emptas, & collectas: in mufæo suo, post se reliquit. *Sor.* Quomodò ergo prodiit in lucem historia illa Lugdunensis? *Botan.* Paucis id tibi perstringam: Hæredes ejus aliquot post mortem annos, cum Typographis lucrum faciendi avidi; ipsis pretio & chartas & Icones prædictorum Authorum vendidêre. Typographi autem, rem totam Medico oudam (illius oppidi) dicto Molinæo, commisêre. Hic cum Typographis, aut saltem cum erumenâ consultans, talem, qualem præmanibus habemus, historiam Lugdunensem

 edidit.

Digitized by Google

edidit. Primò Latinè, deindè Gallicè, & in duos Tomos, in folio diſtinxit: Utraque editio eſt vitioſa & depravata, ut colligere poteſt quivis; ex animadverſionibus Caſpari Bauhini in hiſtoriam Lugdunenſem, ubi quingentæ plus minuſve plantæ, bis terve repetuntur. *Soc.* Animadverſiones, Caſp. Bauh. in hiſtoriam Lugdunenſem vidi, ipſamque hiſtoriam perluſtravi; & gravitèr Authorem, Molinæum ſic à te deſignatum, hallucinatum eſſe obſervavi: ſed ad propoſitum redeamus. Quid dicis quæſo, de methodo Dalechampii? *Botan.* Eſt nulla, aut ſaltem parùm idonea, ad methodicè plantas diſcendas. *Soc.* At ego (ni fallor) aliquam obſervavi methodum, & ut puto à locis natalibus, ſeu natalitiis deſumptam. *Botan.* Æquum dicis. Oblivioni tradideram; ſed jam revoco in memoriam. *Soc.* Sed quid tu de iſta methodo ſentis? *Botan.* Eſt planè ex meâ ſententiâ rejicienda. *Soc.* Quare? *Botan.* Quia plantas per natales (utplurimùm) ſeu natalitia loca diſponit: Et ego libentèr à te peterem; quid quæſo debet impellere vel mediocritèr Botanicæ rei ſcientem, disjungere Ranunculum à Ranunculo: an quia quidam creſcunt

in

in ſummis Alpium jugis & Pyrenæis montibus, alii in ſylvis, ſive annoſis, ſive cœduis, alii deſcendunt ad prata & loca campeſtria; novalia & arva ſcilicèt; alii deniquè in paluſtribus & aquoſis locis proveniunt: Sed cum omnes habent teſſeram ſeu ſymbolum Ranunculi, debent ſimul diſponi in unâ aut alterâ Tabulâ cognationis. *Soc.* Quid tu vis, per teſſeram .ſ. ſymbolum? *Botan.* Ego per teſſeram intelligo notam Genericam, & eſſentialem; toti Seriei Ranunculorum convenientem, ubicunque locorum creſcentium. *Soc.* Quænam eſt nota eſſentialis Ranunculorum. *Botan* Eſt habere florem ex quinis, utplurimùm (Aſiaticos excipio, qui à luxuriante naturâ ſex, ſeptem, octo; nunc plura, nunc pauciora producunt) petalis conflatum; eidem flori, ſuccedit, racematìm, aliquando orbicularitèr, aliquando longitudinalitèr, ſpica; ex ſeminibus compreſſis, acutis compacta. *Soc.* Jam intelligo quid ſit nota Generica hujus longè patentis generis: ego nunquam hoc obſervavi ante hanc tuam deductionem; ſed in pratis & ſylvis, imò in hortis ipſis ſuburbanis, diligentiùs impoſterùm obſervabo. Unus mihi reſtat ſcrupulus, qui

me malè habet, de temperamento Ranunculorum. Suntne omnes calidi, & cauſticæ facultatis, quod nobis innotuit primò (niſi me animus fallit) ex Mendicis publicis, qui Ranunculorum foliis, tibiis applicatis, exuunt cuticulam ſeu epiderma; imò nonnunquam cutis ipſius & muſculorum magnam partem, eamque ulcerant, ad prætereuntium movendam commiſerationem, & ad vilem ſtipem ſeu Eleemoſynam obtinendam? *Botan.* Hoc lippis & tonſoribus notum: hoc præſtant viles animæ, Mendici: Confiteor maximam Ranunculorum partem, eſſe cauſticâ facultate prædítam, ſed non omnes. *Soc.* Qui ſunt illi, qui non ſunt calidi? *Botan.* Ipſæ Authorum denominationes hoc tibi indicant, ut Ranunculus Holeraceus, Ranunculus dulcis Tragi, Ranunculus Muſchatella, dictus, &c. advocetur guſtus in opem. *Soc.* Ego conſtantèr aſſeveravi, cauſticam facultatem fuiſſe de eſſentiâ omnium Ranunculorum. *Botan.* Contrarium vides, & ego Anthoram, quamvis contrariam habeat virtutem (ſi credendum eſt veteribus ſcriptoribus) ex Aconitorum genere, à flore Cucullato, & Thecis membranaceis, ſeminibus Angulosis

guloſis, ſubrotundis, aſperis (aliorum Aconitorum more) eſſe concludo. Quin & inſuper, ſi bene obſervaveris, Delphinium, aut Conſolidam regalem, per eandem rationem, Aconiti ferre teſſeram, percipies: & ideò ego ſic Tabulas cognationis .ſ. affinitatis conſtruendas ſuſcepi, ſequendo ductum naturæ, ſolâ naturâ monſtrante mihi methodum & viam; à fine ultimo petitam. Videlicèt fructificationis ſimilitudine, ut prolixiùs alibi ex me audies, ut plenè ſatisfiat tibi: ſecundam, adferam in medium inſtantiam. An tu liberè disjungeres Lapathum aquaticum, ſeu Hydrolapathum, à Lapatho diverſi generis ſuburbano, aut montanum, ſeu Lapathum Monachorum, ab utroque prædicto genere, Lapathorum? Quia in montibus celſis Alpium provenit: ſicut Rhabarbarum in Thraciæ montibus altis. *Soc.* Quid, tu Rhabarbarum ſtatuis Lapathum itidèm? *Botan.* Nil aliud eſt κατ' ἐξοχὴν, quàm omnium Lapathorum optimum. *Soc.* At Caſp. Bauh. disjungit & per ſe collocat, nec in medium producit aliquem Authorum .ſ. veterum .ſ. Neotericorum, Lapathum appellantem: Sed ab omnibus Rha, & Rhaponticum ſeu Rhabarbarum dicitur.

Digitized by Google

Botan. Ego nunquam eſſentiam plantarum, deſumendam ab Authorum denominatione putavi, ſed à florum formâ, & ſeminum conformatione; vide *pag.* 117. *Col. II. Pl. V. in* hallucinationibus Caſp. Bauhini à nobis exhibitis, præludiorum Botanicorum noſtrorum, parte ſecundâ: Et videbis quam confusè Caſp. Bauh. digerit capita illa duo de Lapatho, & Rha ejuſque ſpeciebus. *Soc.* Ergò tu concludis, Rhabarbarum eſſe Lapathum, à fine ultimo, id eſt, ſimilitudine ſeminum triangularium, ut ſunt omnium Lapathorum, ſemina triangularia ſ. triquetra, rubra aut nigra. *Botan.* Quid ni? noli quæſo de hoc dubitare. *Soc.* Poteſne alias dare inſtantias? *Botan.* Centum, imò ſexcentas ſi velis. *Soc.* Aliam audire libentèr velim. *Botan.* Noſtine vulgarem & putidam illam plantam (ejuſq; ſpecies varias) quæ Atriplex dicitur? *Soc.* Nihil meliùs. *Botan.* Nondumne percipiſti à ſemine; duo eſſe ejus genera; ab Authoribus nunquam (ante me) obſervata? *Soc.* Hoc me latet. *Botan.* Ego paucis tibi apperiam: primò, ſi obſervaſti Atriplicem hortenſem, rubram & albam; ſemina ſpicatim producit, in ſummis caulium, ſingula, binis

nis foliolis incluſa. *Soc.* Quid tùm ſequetur? *Botan.* Obſerva in ſimetis & ſterculinis foſſiſque ſuburbanis, Londinum circumjacentibus, eſſe plures hujus generis ſpecies ſubalternas; quæ habent ſemina ſingula binis foliolis incluſa, ut Atriplex folio haſtato ſ. Deltoïde in horto Regio Bleſenſi aucto; etiam Atriplex Polygoni folio, Halimus ſeu Portulaca Marina, ſic Authoribus denominatæ; Sunt veræ & genuinæ Atriplicis ſpecies, quemadmodùm Atriplex Halimi folio: quia hæ noviſſimæ Atriplicis ſpecies ad oram maritimam, aliæ prædictæ in ſimetis & Stercoratis ſuburbanis locis, aliæ priùs dictæ in hortis cultis proveniunt; nonne debent ibidèm digeri ſeu collocari, quondoquidem eandem gerunt teſſeram, ſeu notam eſſentialem? *Soc.* Do manus, jam abundè mihi ſatisfactum eſſe puto de methodo depravatâ Dalechampii; atque uno verbo velim aliquid mihi dicas de methodo Parkinſoni noſtri, qui ſatis ingentem farraginem plantarum, omnium Authorum antè ſe ſcribentium, in magnum volumem congeſſit compilavitque. *Botan.* Parkinſonum laborioſum fuiſſe ſenem, multaque tranſcripſiſſe fateor,

fateor, quin & numerum auxit; ſed magis confundit genera plantarum quàm quivis, dum novam vult (ex proprio cerebro) cudere methodum. *Soc.* Quomodò hoc quæſo demonſtras? *Botan.* Obſervaſne ipſum à plantarum ſeu ſtirpium viribus vel facultatibus, claſſes ſuas deſumpſiſſe. *Soc.* Atqui hæc videtur mihi optima methodus. *Botan.* Eſt Medicis aliquomodò utilis & neceſſaria, ſed non eſt à naturâ deſumpta; non eſt de eſſentiâ primò & principalitèr plantæ; purgare, necare, .ſ. occidere, calefacere, aut frigefacere, &c. ſed ſecundariò & accidentalitèr, ut elariùs tibi alibì oſtendemus. *Soc.* Ego omnes purgantes, ſimùl collocatas, optimam ſemper putavi methodum, pro iſtâ ſaltem claſſe. *Botan.* Ne multis verbis tempus teramus; Vis ut uno verbo dicam? quod ſentio repugnat methodo naturæ & per conſequens noſtræ, per Tabulas cognationis digerendæ, & ut capias faciliùs quod propono, libenter à te peterem, quam affinitatem habet Rhabarbarum, ſeu Rhaponticum quod Lapathum ſuprâ (ſi memor es) à fine ultimo, ſcilicèt fructificatione, clarè ſatis fuiſſe demonſtravi: quid inquam affinitatis habet Rhabarbarum, cum ſcammo-

niâ

niâ Antiochena, quæ eſt Convolvulus? Ob uſum ſucci lacteï, qui ex eodem manat. Quique diagredium poſt præparationem dicitur: Eſt per excellentiam, omnium Convolvulorum præſtantiſſimus: aut dic mihi. Cucumis Aſininus, ex quo Elaterium Medicorum antiquorum extrahitur; ſecundum methodũ naturæ, nil habet affine cum prædictis ſtirpibus? Fateor quod cum Colocynthide, & à naturâ, & à virtute diſponi debet: Deindè vide differentiam Tithymalorum, Eſulæ, & cœt. (id genus) aliarum, tricoccarum lactescentium, à Jalapio, Senna, Tamarindo, Caſia. Quæ cũ prædictis & poſtea nominandis, nullam habent inter ſe connexionem, ſi exacta & omnium perfectiſſima methodus naturæ conſideretur: purgare enim hiſce nominatis, & multis aliis in obſervationibus noſtris generalioribus, ex horto Regio Bleſenſi aucto collectis; hiſce inquam purgare accidentalitèr datur à naturâ. Ita ſecundariò .ſ. accidentalitèr à purgante facultate poſſunt omnes hæ ad unam claſſem reduci; ſed ex Libro naturæ, habitâ ratione finis ultimi, ſcilicèt fructificationis; toto Cœlo differunt. Quod non te jam poteſt latêre; poſt tàm prolixam, de hac re,

re, ratiocinationem. *Soc.* Video & concedo novam illam Parkinſoni, non veram & genuinam (ſecundum naturæ intentum) eſſe methodum : properemus ad alios; ne dialogus in immenſum excreſcat. *Soc.* Quid *dicis* quæſo de magno illo volumine duobus Tomis immenſis, diſtincto? Quod, hort. Eyſt. nuncupatur. *Bot.* Laudo liberalitatem iſtius Epiſcopi magnoperè, qui tantum impenſarum liberè fecit; ut publico prodeſſet. Nullus inquam, Icones omnes, æri tàm affabrè & elegantèr inſculptas (heu nimis tamen magnas) tam gloriosè & ſumptuosè edidit. *Soc.* Sed ad rem : Quid dicis de methodo ipſius Beſleri; qui Librum compilavit? *Botan.* Vis ut apertè loquar? Quod ſentio, noſti (ſi benè meminiſti) Beſlerum compilatorem, ſeu directorem, prædicti Libri; per anni quatuor tempeſtates, plantas ſuas digeſſiſſe : ut acupictoribus, prædictum Librum evolventibus & intuentibus plus prodeſſet, quam Botanicæ ſcientiæ, ſtudioſis. *Soc.* Certè Icones iſtius Libri, ſeu hort. Eyſt. (ſi mavis) elegantèr ſunt deſignatæ & ſculptæ, & pro proportione plantarum quarundum minorum : Sæpè reperiuntur nimis magnæ. Inſuper fateor Librum

brum illum principum & magnatum muſæis, & acupictorum ſeu acupictricum, cubiculis valdè eſſe neceſſarium. *Botan.* Utinam haberem ego talem patronum, ego methodum naturæ, de quâ ſuprâ diximus ſuppeditarem & patronus deſignatores, & ſculptores conduceret & pretium perſolveret. *Soc.* Ego id cordicitùs optarem, & optare debent omnes quotquot in Europâ extant Botanicæ rei amatores. Sed quid dicis de Matth. Medico Italico, nullamne ille propoſuit methodum? *Botan.* Ille & multi alii ſuo tempore, Coætanei Botanici, ſimilitudinem foliorum .ſ. partium accidentalium ritè, & attentè obſervantes, plantas multas ad eandem claſſem reduxêre, quæ aliò à nobis ſunt relegandæ. *Soc.* Tu appellas folia, partes accidentales ſtirpium? *Botan.* Sunt tantum; quemadmodum radix, Fibræ .ſ. Nervi, caules: cur non folia? Hoc ego paucis deducam, radix ex terræ ſuperficie, tanquam matrice, attrahit nutrimentum ſeu alimentum, caules vehunt ad omnes partes, per ductus illos ſive canales, qui Fibræ .ſ. Nervi dicuntur; folia arcent calorem & frigus: ergò per accidens omnes inſerviunt productioni ſeminis; ejuſdemque perfo-

perfectioni. *Soc.* Cui uſui data ſunt folia, præter ad arcendum frigus & calorem, & ad Dod. ductum .ſ. vehiculum Nervorum per omnes partes plantæ. *Botan.* Inſerviunt ad ſubalternas ſpecies conſtituendas etiàm, aliquando multa genera intermedia deſignantur ab accidentali à ſimilitudine foliorum. *Soc.* Certè ego, certum indicium Genericum à ſimilitudine foliorum peti exiſtimavi. *Botan.* Et ego hoc obſervavi in multis Authoribus ſcribentibus aliquot abhinc annis, imò non ita pridèm, in Libro quodam ab Authore, veſtri Collegii Socio edito, in quo exhibetur methodus, per Tabulas, diſponentes claſſes plantarum, à ſimilitudine foliorum, ſumptarum. Ego tantum confuſum Chaos: illìc, de plantis legi, nec quicquam didici, ut monſtrabo tibi & lapſus & confuſionem, aliás. Quia degrediendum nunc, non eſt. *Soc.* Ego vix clarè percipio quid tibi vis dum folia non vis, inſervire ad digeſtionem plantarum? *Botan.* Ego nunquam hoc dixi; ſed audi attentè non inficias eo, folia multùm conferre ad ſpecies intermedias denominandas, ſed non ad Genericam, & eſſentialem aſſignandam notam. *Soc.* Clariùs velim te temetipſum

sum exprimere. *Botan.* Quandoquidem exempla clarè docent, quædam asseram, in medium: quæ te convincent, plantas, non denominari debere à primo intuitu, seu conformatione foliorum ad seinvicem. *Soc.* Adfer quæso in medium exempla illa tua. *Botan.* Audi ergo attentè. Observatine in citatione Authorum apud Cæsp. Bauh. complurima Ranunculorum genera subalterna, ab accidentalibus (foliis scilicet) partibus, desumpta, ut exempli gratiâ, Ranunculus Gramineis foliis, apparet Botanicis non mediocriter veritas in arte: Gramen sub pedibus: sed fimulac florem Mense Junio produxerit luteum, quinis petalis conflatum, & semina Ranunculi in modum; Nemo dubitabit ipsâm illam plantam esse Ranunculi speciem, ipsamque nominare Ranunculum; & tot multa paucis comprehendam, observa mecum: Ranunculus Plantaginis folio, Ranunculus Rutæ folio, Ranunculus Beupleuri folio, Ranunculus Thalictri folio, Ranunculus Aconiti folio, rotundo folio & diffecto folio. Antequam maxima pars horum floreat femenque produxerit, non debent nominari à primo intuitu foliorum, per terram stratorum.

Digitized by Google

rum, ne primus pro Gramine, ſecundus pro Plantagine, tertius pro Ruta, quartus pro Thalictro habeatur, & ità deinceps: ſed ſecundariò à ſimilitudine foliorum, ad ſpecies ſubalternas determinandas; ergò clariùs, jam luce meridiana, vides notam Generícam nunquam deſumi à ſimilitudine foliorum, ſed (ut antè à me dictum) diverſas ſpecies ejuſdem generis conſtituit, diverſa foliorum ſimilitudo ſeu facies. *Soc.* Video clarè & perſpectè in hoc genere, videlicèt Ranunculorum: Folia tantum per accidens & ſecundariò (quoàd formam & figuram) inſervire ad ſpecies deſignandas; ſed an hoc contingit paſsìm in omnibus, plantarum generibus? *Botan.* Paſsìm, ubi numeroſæ ſunt ſpecies ut hìc. *Soc.* Sed danturne plures ſpecies intermediæ diverſæ, ſub diverſis omninò generibus, quæ folia habent conformia, & à ſeinvicèm vix cognoſcibilia. *Botan.* Auſculta paucis, attentiſſimè obſervabis mecum. Milleſolium Tanaceti foliis; flore purp & alb. & Erucam Tanaceti foliis: Utraque notatur à me non ſcripta, in horto Regio Bleſenſi aucto, Tanacetum inodorum Bauh. in Pin Bellis Tanaceti ſolio nobis: Hæ tres plantæ, notâ

Gene-

Generica & eſſentiali, toto Cœlo diſſerunt, à ſine ultimo petitâ. Sed hæ tres plantæ, dum primo è terrâ pullulant, & antequam cauleſcant: intuendo, acutâ acie oculorum, propter conformitatem (alarum ſucchi ſeu Nervo medio adhærentium & diſſectarum) in foliis, non poſſunt facilè diſtingui; ſed paulatim cauleſcente primâ plantâ, & florente, ſemenque poſtquam ad maturitatem produxerit eadem: Millefolium, Tanaceti foliis ſeſe declarat, ſimendo denominationem, à nobiliori parte, finc ultimo fcilicet, & folia data diſſecta, Tanaceti adinſtar delignant, pro qui ſpecie Millefolii, haberi debet. Secunda pariter planta, ad flores vergens, ſeſe tetrapetalam & ſiliquoſam prodit: addeacrem & calidam qualitatem, ſi Maſticetur, guſtuſque vocetur in opem; Eruca ſtatim apparet: folia Tanaceti inſtar inſervitum ad ſpecificam hanc Erucæ ſpeciem delignandam. Tertia florens, Bellidis adinſtar, diſcum fert luteum, quem cinguut, circulariter ſeu ſtellatim, lata alba folia; unde Bellis Tanaceti folio dicitur: eam denominando à parte eſſentiali; ſiquæ videlicet, aptius ad genus Bellidum à me reducitur; & folia Ta-

Digitized by Google

Tanaceti in modum, conſtituunt ſpecificam ejus Bellidis notam. *Soc.* Jam tandem percipio folia non eſſe de eſſentiâ plantarum, ex exemplis ſatis ingeniosè à te allatis. *Botan.* Gaudeo me nullam poſticam dubio *tuo* reliquiſſe. *Soc.* Quid dicis de methodo Joh. Bauhini, qui hiſtoriam univerſalem, plantarum ſeu Stirpium in tres Tomos digeſtam, ſcripſit? *Botan.* Vis certè ut ingenuè fatear, quod res eſt; omnium Authorum, ante ſe ſcribentium, optimam reliquit methodum. *Soc.* Nihilne corrigendum aliis reliquit? *Botan.* Imò multa, ut videbis ex animadverſionibus meis in hiſtoriam univerſalem ejuſdem Joh. Bauh. ibì multa tranſpono, diruo genera; alia de novo creo & conſtituo; denique omninò novam condam & componam methodum, meliorem, facilioremque. Quâ, plus addiſcet methodicè. Botanicus diſcendi cupidus; biennii ſpatio, quam ego ad decennium didicerim. *Soc.* Utinam eandem præ manibus haberemus, & optarem tibi publicam Cathedram, ex quâ, tam vivâ voce; indoctos præſentes doceres, quàm ſcriptis è longinquò doctis ſcriberes. *Botan.* Ego me ſemper divinæ providentiæ me commiſi,

&

& adhuc idem præſto. *Soc.* Quid tibi videtur de prolixitate Johannis Bauh. in deſcribendis aut potiùs tranſcribendis, aliorum Authorum plantis. *Botan.* Tedioſum ipſum puto in hoc, ipſumque dum dicta & ſcripta aliorum, in medium producit; citando dicta, & allegando ſcripta Mattheoli, Fuſch. Lob. Dod. Cluſ. & cœterorum tam, veterum quam Neotericorum ſcriptorum, in tria immenſa volumina ſeu (ſi mavis) tres Tomos, ejus Hiſtoria univerſalis accrevit. *Soc.* Sed quid quæſo emolumenti advenit, Reipublicæ Botanicæ; dum tantam molem ſeu farraginem, dictorum Authorum in in tam vaſtam hiſtoriam congeſſit. *Botan.* More Germanorum & Teutonum, ut ſe laborioſum & diligentem indefeſſumque collectorem, ſeu compilatorem poſteris monſtraret ſeu exhiberet. *Soc.* Te verum dicere paſſim obſervo & in hoc tibi aſſentior: ſed tu qui exactiori trutinâ, omnes Botanicos perpendiſti; quot quæſo addidit plantas, nondùm deſcriptas, aut non ſculptas ab aliis Authoribus Botanicis, ſcribentibus ante ſe. *Botan.* Ego nondùm calculum exactè inivi. *Soc.* Proculdubio con-

jecturam veram facere potes; & fortaſſe exactum, etiam plantarum ab ipſo edditarum numerum, nobis exhibere potes. Vide Schedulas tuas de hoc recollectas. *Botan.* Niſi me memoria fallit plantæ à ſolo Johanne Bauhino in Tomo ſecundo ſuo, & tertio, figuratæ, non excedunt centum, ſeptuaginta tres. *Soc.* Exhibeas quæſo eaſdem nominatîm cum pag. ipſius Tomi primi, ſive ſecundi, ſive tertii. *Botan.* Quid hoc proderit Reipublicæ Botanicæ; & quod non amo, in aliis, ſed vitupero; committam ipſe: Hoc eſt ut appertis verbis tibi dicam, tranſcriberem. *Soc.* Eſt notare tantum quot acceſſêre ab ipſo Bauhino tanto operi, quod quinquaginta annorum fuiſſe, ipſe aſſerit. *Botan.* Forte morem geremus amicis tuis precibus. *Soc.* Sed ſi recte impræſentiarùm recordor; dum cederemus ſermones Botanicos, nos, inter nos: Tu certiorem me feciſti, te habuiſſe plantas non ſcriptas; cuiquam Authorum; ducentas quinquaginta. Quarum nobis deſtinaveras ſuccinctam, & curtam delineationem. *Botan.* Hoc hactenùs; in præludiis meis Botameis præſtiti.

præſtiti. *Soc.* Ego plurimi facio, Enchiridium tuum Botanicum, Manuale continens præludiorum Botanicorûm tuorum, partem primam, & alteram. *Botan.* Fortè plus emolumenti & voluptatis ex lectione hujus Libelli, Botanicæ rei amatoribus & reverâ ſtudioſis proveniet. Quam ex tedioſorum prædictorum ingentium voluminum contemplatione & evolutione, quæ nauſeam etiam ingenioſis lectoribus (expertus loquor) pariunt. *Soc.* Omittamus prædictos omnes prolixos tranſcriptores, & accedamus ad Caſparum Bauhinum, quæ eſt tua ſententia de Pinace ſuo. *Botan.* Plus ille laboravit quàm omnes ſuprâ citati Authores. Nomina omnium Authorum, ſtirpes diverſimodè nominantium, in medium producit; ipſos concordare facit. Quod non minimum præſtitit: ad evitandam confuſionem ex multiplici Nomenclatura Authorum, enatam. *Soc.* Sed de ejus methodo, in digerendis plantis: tuum judicium tua exiſtimatio valebit. *Botan.* Vide hallucinationes Caſp. Bauh. in Pin. in præludiorum Botanicorum meorum parte poſteriore: Ibi obſervo ipſius halluci-

nationes in Pin. tam in denominandis, quam digerendis plantis, ubi diruo, ædifico genera; muto genera cum generibus, de his conſule locum citatum. *Soc.* Utinam haberemus ad manum tuam methodum promptam; deſignatam & impreſſam quâ inſtructi, methodicè poſſimus plantarum ſeu ſtirpium, addiſcere affinitates ſeu cognationes. *Botan.* Brevi, ſi Deus dederit & Mœcenas ſeſe obtulerit, (per me non ſtabit) habebitis. *Soc.* Unde tu hanc methodum didiciſti, quæ non reperitur apud omnes prædictos Authores. *Botan.* Ex Libro naturæ. *Soc.* Quid tu appellas Librum naturæ. *Botan.* Ego appello Librum naturæ, id quod omnibus innoteſcet. Quam facillimè poſtquam præludia mea diligentèr evolverint, & hiſtoriam univerſalem plantarum ſeu ſtirpium meam methodicè diſpoſitam; viderint. *Soc.* Quid intelligis per methodum univerſalem. *Botan.* Ego intelligo novam methodum per Tabulas cognationis, ſeu affinitatis diſponendam. *Soc.* Quid tibi vis per Tabulas cognationis ſeu affinitatis; nunquam ego legi de *tali* methodo apud Authores. *Botan.* Nec

ego:

Digitized by Google

ego: ergò non eſſet nova; etiamſi fuit ab incunabulis mundi (ut impoſterùm audies) ſed nemini adhuc detecta ante me: & neutiquàm dubito, quin illa methodus, & tibi & aliis omnibus φιλοβότροις; (ubicunque locorum viventibus) ſatis grata futura ſit. *Soc.* Valdè cuperem eandem illam methodum videre, & optarem, ut ex tuâ Pharetrâ illa ſagitta, à tot & toties expetita, exiret. Nam ſæpiùs tecum inivi colloquium, de arte noſtrâ Botanicâ; ſatis adhuc ut video confuſa: Aliquid ex te aliàs hauſi, utinam & mihi & Publico ubique terrarum, brevì proficias; ſed dum recordor, & ne mente excedat, Mattheolus Italus ille Medicus, qui confuſam edidit plantarum hiſtoriam, multa appinxit falſa arboribus ut ipſas congruere faceret, cum antiquorum deſcriptionibus. *Botan.* Hoc non me latet, ſed quod magis miror: Imperatorem, Reges & Principes, & magnates omnes, totius penè Europæ, contribuentes habuit, hæc ipſe jactitat in Præfatione Libri ſui. *Soc.* Utinam Reges & magnates hujus ſæculi, Chalcographorum, & Typographorum labores, tuoſque com-

penſarent; ut methodo illâ novâ, & inauditâ; à naturâ datâ: Et à te ſolummodo (citrâ jactantiam) obſervatâ: frueretur, totus terrarum orbis; ut in arte *Botanicâ*, quondàm Italia, Germania, Gallia gloriatæ ſunt; ſine methodo. Itidèm nobiſcum, cum methodo, gloriari Britannia noſtra poſſit: Jam tandem unam à te ſciſcitarer quæſtionem; antequam decedas: plantæ poſſuntne ſine ſemine; aut ſatione ſeminis, provenire, ſponte è viſceribus terræ, tanquam ex matrice. *Botan.* Jam grandem mihi proponis quæſtionem quæ anxios, & ad ſolutionem tardos reddere poteſt, & ſocios Pariſienſes & Londinenſes; imò vexaret ſortaſſis, ſocios Lyncæos Romæ olìm viventes. *Soc.* à te hanc nunc expoſtulo, tu qui tot ſemina collegiſti, ſeminaſti, recondidiſti in promptuario tuo ſeminario; ex horto Regio Bleſenſi: totius Europæ, non ita pridèm, ſub obitum principis, Mœcenatis ipſius; refertiſſimo: Earundem omnium ſtirpium (quotquot illìc ſemina; tulêre) dicis te habere bis Mille quingentarum vegetantium ſemina. Iterùm, jam ad rem; putaſne

tu

tu tellurem, tanquam ex propria matrice, plantas sine semine aut satione producere? *Botan.* Certè lupum auribus teneo, an laterem lavabo, an oleum & operam perdam mihi adhuc incertum. dicam tamen quod sentio. Primò exordiamur à divinis oraculis, ubi affirmant in Genesios cap. I. versibus 11, & 12. producat terra Gramina, & herbas reddentes semina sua: & Arbores ferentes fructus secundum carnem genere eu species, continentes in se semina, vidit Deus quod hoc fuit bonum, &c. *Soc.* Ex hac citatione divinæ Scripturæ innuere videris: omnia vegetantia seu floripes seu Arbores ex proprio semine, & Deo, tertio die creata produci, in terris, seculorum, ad finem mundi: Ex diversum tam genorum, quàm specierum, diversis seminibus à Deo datis, provenientium. *Botan.* Hoc ego ità intelligo & longâ experientiâ edoctus, observavi, à minimâ plantulâ, seu Graminis herbâ (ut cum Virgilio loquar) usque ad Libani Cedros. *Soc.* Sed multæ conspiciuntur plantæ, quæ sponte provenunt sine flore, capsulâ seminali. *Botan.* Ego fateor plurima elle genera & infinitas

finitas ſpecies (ſi ita loqui licet) ſub ipſis contentas, quæ ex mixturâ ſalis, & ſulphuris è terra aquave producuntur, ſponte & ſine ſemine; ego autem, has omnes imperfectas appello plantas vegetantes. *Soc.* Quas tu appellas perfectas; quas imperfectas. *Botan.* Perfectas appello quæ florent; capſulas producunt ſeminales, & ſemina inibi contenta. *Soc.* Jam capio quid velis, ſed unum à te peterem. Capillares omnes, ut Phyllitis, ſive Scolopendria, aſplenium, Coterach, Ruta muraria, Polypodium & Filices omnes, cœteræque (id genus) aliæ: ſuntne perfectæ plantæ? *Botan.* Imò ſunt, quia habent flores (qui fugiunt quaſi obtutum) & ſemina, quaſi pulviſculum in dorſo alarum. *Soc.* Hoc tu appellas ſemina: Capillarium, & earundem familias. *Botan.* Quidni appellarem, & quemadmodum non omnis fert omnia tellus. Idem hoc ſemen ſi aquoſis & umbroſis ſylvis, vetuſtis muris aqua riguis commitatur, pullulat, producitque ſui generis plantam: Si propriam ſortiatur matricem. *Soc.* Quid tibi videtur de junco, an eſt planta perfecta. *Botan.* Perfecta certè

certè omninò, nam floret & semina (utplurimum) triangularia producit. *Socius.* Collegistine unquam junci semen. *Botan.* vigesiès, & habeo penes me in promptuario meo seminali, dena aut duodena distincta, juncorum seminum genera; aut si mavis species appellare, per me licet. *Socius.* Hoc ego nondum adverteram. *Botan.* Hoc clarè tibi demonstrabitur quando velis. *Soc.* quas quæso, appellas tu imperfectas plantas? *Botan.* quæ (ut suprà dixi) sponte sine semine proveniunt: ex quadam commixturâ salis .s. fixi .s. volatilis, sulphuris; junctâ terrâ ruderosa, stercorali denique, aut aquâ stagnante, vel fluente; & in motu perpetuo ut marina, vel fluviatili: seu dulci fontali. *Soc.* Recense quæso quædam imperfectarum plantarum genera, si lubet. *Botan.* In superficie terræ maxima pars omnium muscorum, sive in sylvis annosis, sive densis & umbrosis, sive in vetustis arboribus, muris antiquis, casularum tegumentis aut Lapidum saxorumve superficiebus crescentium. *Soc.* Ergò, generalitèr (ut percipio ex te) omnes musci sunt plantæ imperfectæ. *Botan.* Ego

hoc

hoc non affero. *Soc.* An dantur muſci qui florent & ſemen ſuum perficiunt. *Botan.* Etiam plurimi ut polytrichum utrumque, majus & minus Apulei. Muſcus clavatus, muſcus abietinus & multi alii, quos in ſylvis annoſis & umbroſis obſervare licet: lichen piliatus, ſtellatus itidém. *Soc.* Ego hoc obſervabo exactiús. *Botan.* Fac quæſo. *Soc.* Quid dicis de fungorum Cluſii ingenti catervâ. *Botan.* Sunt excreſcentiæ terræ: (ſive ſunt edules ſive lethales) nec ſemen nec florem habent ſed ſponte noctè unâ aut alterâ è terra emergunt; ex quadam commixtione Salis, ſulphuris, junctâ terræ pinguedine ex ſtercore quadrupedum orta. *Soc.* Ego hæc omnia vera eſſe perſuaſum mihi ſatis habeo: & eſt conſentaneum rationi; Sed quid infers de tuberibus terræ. *Botan.* Tubera terræ in gremio terræ ſuperficiali producuntur ſponte: ſine radice, caule, fibris, aut foliis. *Soc.* Quomodo augentur aut ad talem magnitudinem ereſcunt. *Botan.* Per juxtapoſitionem: ut mineria, & lapides terræ. *Socius.* Sit ita: Non reperiuntur hic in Anglia. *Botan.* Neutiquàm, nec in borealibus

bus Galliæ tractibus. *Soc.* Nunquamnè reperisti in meridionalibus tuis peregrinationibus, sub auspiciis sereniſſimi principis Gastonis Franciæ. *Botan.* Etiam in agro pictavienſi, Burgundia, quin & Gallopro-vincia, & Gallia Narbonenſi reperiuntur, frequentér. *Soc.* Quomodo deteguntur? *Botan.* A porcis ſive ſuibus. *Soc.* Participem, me fac hujus venationis tuberum. *Botan.* Paucis tibi dicam, inſtinctu quodam naturæ, porci ſive ſues odoratu ſuperficialem terræ partem, detegunt; & ſtatìm terram evolvunt quoad appropinquent ad tuber ipſum, & ſtatìm ſubulcus accurrit, & abigit pedo aut baculo ſuo paſtorali, porcum; & lævisſimo labore colligit tubera: Ecce habes quod vidi, in agro pictavienſi & habui ex relatione ſubulcorum aliorum. *Soc.* At ut ego audivi valdè avide expetuntur, à magnatibus Galliæ, Italiæ, qui Embammatis variis, gulam irritantibus utuntur. *Botan.* Ita eſt. *Soc.* Sed quid tibi videtur Corallum & Pſeudocorallum & ejus generis aliæ complurimæ, maritimæ in fundo maris vegetantes, dum in loco nativo ſunt; adhuc

ſunt

ſunt molles; ſed extra mare dureſcunt & colorem mutant. *Botan.* Idem mihi dicendum videtur, quod de diverſitate numeroſa, fucorum marinorum; (ſi mavis algæ marinæ) qui omnes creſcunt in *fundo* maris, & tempeſtatibus, propriis ſedibus avulſi: Ad oram maritimam ejiciuntur multiformes, & diverſarum proportionum. Præcipuè hyemali tempore, verſus oram maritimam, Borealium regionum. *Soc.* Ego multas facie diverſas fucorum marinorum ſpecies, nunc penes me, in muſæo meo habeo; tibi in hoc accedo: puto *certo* certius, has omnes ſive algæ, ſive fucorum marinorum ſpecies, eſſe plantas maritimas imperfectas: Quandoquidem, nec florent nec ſemen producunt. *Botan.* Si non longè fallor; ſunt & plantæ imperfectæ in fluviis & fontibus aliiſque rivulis. *Soc.* Benè obſervas ut conferva, lenticula & ejus generis aliæ multæ aquaticæ, aquis innatantes ſine flore, ſine ſemine: Jam abundè mihi ſatisfactum puto; de imperfectis plantis, tam terreſtribus, quam aquis marinis, ſive dulcibus, ſtagnantibuſve innaſcentibus; ſed quoniam adveſperaſcit:

&

& tù longè hinc diſtas, putaſne plantas perfectas, ſpontè & ſine ſemine produci poſſe, in locis ubi ſemen nunquam fuit ſatum? *Botan.* Suprà ſi ſatis meminiſti, hanc quæſtiunculam examinavimus, & firmitèr credidi, quod tibi hactenus aſſeveravi, & certè ſæpè ſæpiùs affirmavi, conſtantèr: ſereniſſimo principi Gaſtoni Franciæ Domino, Domino meo (dum in vivis erat) colendiſſimo: Et proh dolor! dum jam eſt mortuus, nunquam ſatìs mihi deplorandus. Vivus, ſæpe ſæpius, nos colligere fecit plantas omnes, quæ è rimis lapidum, quibus tegebatur palatium, Regia Chambort, dictum. Multæ fuerunt pappeſcentes illìc vegetantes; ut Senecio, Jacobæa, Conyza acris cœrulea, Lactuca ſylveſtris, & complures ſpecies Hieracii, Dens Leonis, &c. (id genus) aliæ pappoſæ multæ, quas omnes non mirabar, illuc à ſemine per aera volitante, dilatas fuiſſe: Sed & plantas complures, ſolida gerentes ſemina, illìc promiſcuè reperimus, ut Lapathum, Cerafum arbuſculam, Prunum, rubum utrumque, & Roſam ſylveſtrem. *Soc.* At undè venêre ad tantam altitudinem illa ſolida ſemi-

ſemina? concedo pappeſcentia ſemina, per aerem volitare, & in altum attolli; & longè latéque portari poſſe: ſed quomodo ſolida illa & quodammodo gravia, ſemina; potuerint illuc volitare, nondum capio. *Botan.* Ego paucis meam opinionem, notam tibi faciam: Pluriès obſervavi, in altis & vetuſtis turribus, Campanilibuſque Templorum, arbuſculas imò Arbores, è fiſſuris ſive rimis lapidum creſcentes. Ego puto nucleos (ſi fuerunt nuciferæ) glandes (ſi fuerunt glandiferæ) aliaque ſuprà citata, ſolida ſemina ab avibus talia *loca* frequentantibus, fuiſſe tranſlata, in rimas lapidum fortè fortuna cadentia, aut ab avibus illic dormientibus, & noctu quieſcentibus evacuata, germine intacto, ſeu inconcocto pullularunt: atque hoc de aliis ſeminibus omnibus, in talibus provenientibus locis, aſſerere audactèr, tibi audeo. *Soc.* Habet umbram rationis, ſed non plenè mihi ſatisfacit, illa tua ratio: Progrediamur ad illam relationem tuam; quam ex te domi meæ, ſuperiori anno audivi. *Botan.* Quænam fuit illa relatio. Oblivioni tradidi, niſi mihi in memoriam revoees. *Soc.* Fuit obſer-

obſervatio de conflagratione ſeu incendio, totius Europæ, Emporii celeberrimi Londini ſcilicèt: Majoris, ſaltèm ditioris ejus partis. *Botan.* Jam ſubit animum & eandem iterùm paucis tibi perſtringam: Si benè meminiſti, ſecundo die Septembris, Anno Domini, 1666. incepit incendium illud luctuoſum & ad triduum, aut quatriduum duravit. Nec ope humanâ (divinitùs evenit, quum non eſt malum in civitate, quod non fecit Dominus) extingui poterat: Nam Æolus apperto ventorum carcere (ut ita loquar) regnabat: per triduum aut quatriduum illud. Poſt octomeſtre ſpatium, per rudera ducentorum jugerum, ſolo æquatatorum, mihi perambulanti verſus excambium vetus nunc. Antè illud tempus; Collegium Greſhamianum dictum tendenti, in veſtigiis, ædificiorum & tectorum, mihi tanta ſeſe objecit copia, Eryſimi illius, quod irio lævis Apulus alter Fabio Columnæ dicitur: Et eôdem revertens, menſibus duobus poſt hoc; adeò densè pullulavit, ut falce quaſi Triticum, aut ſecale demeti potuerit. *Soc.* Quid

inde

inde ſequetur, unde proveniſſe tantam copiam iſtius Irionis? putas tu; an à ſemine ſeu ſatione. *Botan.* Quid quæſo, te movet ad talem proponendam quæſtionem, cum ædificia omnia circa ædem Divi *Pauli*, & alibì paſsim in meditullio celeberrimi Emporii Londini, à Mille aut ſaltèm, aliquot centenis annis: Fuêre conſtructa & tectis conſervata. *Soc.* Ergò tanta copia illius ſeminis, latebat in cellis & cavearum fundis, & ſoli & pluviæ expoſita, fructificavit. *Botan.* Unum hoc addam: Ego non ſum Plinius, ut ex aliorum re*lati*one mundo imponam, nec Mattheolus ut appingam ea quæ nunquam extitere: Sed ut vis appertis verbis nec Calamiſtratis: meum tibi dicam animum. *Soc.* Dicas quæſo? *Botan.* Nullum eſt ſemen plantæ, quod producit (conſervatum quàm diligentiſſimè) poſt decennium; perraro poſt quinquennium: multò minùs poſt centenos aliquot, & Mille annos. *Soc.* Ergò aliquis ſemina iſtius plantæ, per rudera ſparſit. *Botan.* Non credo imò, certò ſcio tantam iſtius Irionis, ſeminis copiam non fuiſſe *in* tota inſula Britanniâ, imò nec in Gallia: dubito

dubito an in Germania & Italia ipſa; (cujus Neapolis eſt regnum, ubi frequentèr creſcebat tempore Fab. Columnæ) unquam floruit tanta iſtius plantæ copia, ergò etiamſi ſeminatores fuiſſent (ex tuâ opinione, poſt hæc tibi à me auditâ) non poterat tanta copia iſtius individualis ſpeciei, ſeminis; à tot Regnis ſuppeditari. *Soc.* De hoc non multùm nunc dubito: Sed quid concludis, ſis rationi conſentaneus. Unde provenit tanta copia iſtius Irionis, forte ſponte. Sub idem tempus, ibidem vidiſti & obſervaſti multas alias plantas, pappeſcentes, imo gramineas, aliaſque diverſarum claſſium. *Botan.* Vidi & attentè obſervavi. *Soc.* Undè hæ aliæ venêre. *Botan.* A ſemine volatili pappeſcenti quod poteſt (ut ſupra clarè ſatis docui) ad multa Milliaria, vento transferri, & in altum attolli & ubicunque ceciderit, germinat & fructificat. *Soc.* De pappeſcentibus non dubito quod dicis, inſuper Gramina, densè ſatìs proveniunt; in qualibet terra ſi negligatur: quare non poteſt tuum Eryſimum, ſeu Irio lævis Apulus alter in ruderibus Londinenſibus, ſponte etiam provenire. *Botan.* Non

eſt par ratio inter Gramina & Eryſimum hòc: Quia Graminum ſemina ſpargun-tur paſsìm; eſt omnium vegetantium plan-tarum, in omnibus regionibus, frequentiſſi-ma & facilius ſeſe propagat. *Soc.* Eſt planta tamen perfecta, ex ſuprâ dictis à te: ergò à ſemine, multiplicatur. *Botan.* Hoc ego ſemper credidi, & in hanc horam credo. Unum à te ſciſcitari velim, putaſne hanc plantam, Irionem lævem Apulum Col. à quovis hortulano, aut incola hujus civi-tatis ſatam, in ruderibus fuiſſe. *Soc.* Nemi-nem hujus inſulæ primò tam curioſum, ſe-cundò nec tanta ejus plantæ ſeminis, copia inſtructum fuiſſe, pro certo ratum & ſta-tutum habeo. Quis tam ſtolidus aut malè feriatus homo, ſi ſemina ad manum haberet (quod impoſſibile ſuprâ demonſtratum eſt) ruderibus ducentorum jugerum terræ, ſolo æquatorum, committeret. Ergò cum nec à ſatione, nec à ſemine, ad aliquot cente-nos annos in ruderibus latente, produci poterat; hujus plantæ tanta copia. Unde concludere vis, tantam ipſius multitudi-nem proveniſſe. *Botan.* Certè ut ſupra dixi ex ſale partìm volatili, partim fixo, ſalpetro, ſulphure

ſulphure, & ex terra ſive calcoſa aut ruderoſa & aqua, mixtaque materia, quocunque modo appelles, per me non ſtabit. Neſcio quid mihi perſuaſum habere debeo, adhuc. Probabile certè eſt, hanc plantam tam copiosè proveniſſe ſponté; Ut ſupra dictum fuit. Sed hæc opinio apperit januam ad Philoſophaſtros contemplativos, qui indifferentèr credunt cujuſlibet generis plantas, Arbores, frutices, ſuffruticeſve, ex terra tanquam matrice, ſponte ſine ſemine provenire. Sed hæc opinio (ut mihi videtur) repugnat ſacræ Scripturæ, & rationi. Hæc per dialogum inter nos dixiſſe, impræſentiarum, ſat eſſe puto. Quod reſtat de hac materia; Sociis virtuoſis, Pariſienſibus & Londinenſibus, viris nobiliſſimis, clariſſimis & Doctiſſimis (ex quorum numero, te eſſe ſcio) diſcutiendum relinquo. Vale mi Doctiſſime vir.

FINIS.

Digitized by Google

Errata in præludiis Botanicis contenta.

PAgina 1. *linea* 7. lege Abrotanum. *p.* 5. *l.* 18. l. nonum. *p.* 17. *l.* 13. l. Anthyllis. *ibidem l.* 19. Antirrhinum majus paſſim. *pro* ☿ l. ☉. *p.* 19. *l* 6. *pro* ☿ l. ☉. *p.* 26. *l.* 18. l. Halimoïdes. *p.* 34. *l.* 3. & paſſim l. Buphthalmum. *p.* 39. *l.* 2. l. Galactites. *p.* 46. *l.* 14. l. Cerefolium. *p.* 51. *l.* 16. *pro* ☿ l. ☉. *p.* 53. *l.* 5, & 6. *deleatur* humilis. *p.* 55. *l.* 11. *pro* ☉ l. ♃. *p.* 67. *l.* 4. *pro* ☿ l. ☉. *ibid. l.* 14. l. Doronicum foliis viridibus, ſubtùs lanuginoſis. *p.* 68. *l.* 2. l. Leucoium *p.* 69. *l* 5. & 7. l. Elichryſum. *p.* 77. *l.* 1. l. rubris. *p.* 81. *l* 4. l. Geranium. *p.* 91. *l* 7. l. Peruanus. *p.* 96. *l.* 11. l. Piloſella. *ibid. l.* 17. *pro* ☿ l. ☉. *p.* 97. *l. antepenult. pro* ☿ l. ☉. *p.* 110. *l.* 4. *pro* ☿ l. ☉. *p.* 109 *l.* 18. l. ſemine rotundo *p.* 110. *l.* 6. Lapathũ fidiũ inſtar, l. nobis. Pulchrũ ſinuatũ Bononienſe J. B. *p.* 111. *l.* 11. *pro* ☉ l. ☿. *p.* 123. *l.* 22. l ſyl alba ſimplex. *p.* 128. *l.* 10 *del.* ♋. *p.* 129 *l.* 1. *pro* ☿ l. ☉. *p.* 129, & 130 l. Armeniaca. *p.* 129 *l.* 12. l. Aurantia. *p.* 137. *l.* 14. *pro* ☿ l. ☉. *p.* 145. *l.* 16 *pro* ☉ l. ☿. *p.* 149 *l.* 19, & 20. *pro* ἀρύλ☉. 3. l. 1ª. & 2ª. *p.* 160. *l.* 13. l. Virginiana *p.* 160. *l.* 18. l denſis ſurculis. *ib. l* 12. l. Scabioſæ. *p.* 173. *l.* 16. l. 1ma. Cluſ. & *pro* ☿ l. ☉. *p.* 178. *l.* 2, & 4 l. Moſchata *ibid. l.* 21. l. Pimpinellæ. *p.* 197. *l.* 14. l. Sphodylium. *p.* 270. *l.* 2. *pro* ☉ l. ☿. *p.* 213 *l.* 2. l Hybernum *p.* 218. *l.* 6. l ſem nigro *p.* 231 *l. ult.* l. Androſæmi *p.* 243. *l.* 14. l. Cachryophora *p.* 268. *l.* 8. l. Malacoïdes. *p.* 171. *l.* 12 l. hac duplici nota. *p.* 258. *l.* 17. l innititur *p.* 276. *l.* 10. l. ſolia. *p.* 280. *l.* 1, &

2.

Digitized by Google

2. l. fructuſve. *p.* 281. *l.* 13. l. relinquentes. *p.* 282. *l* 2. l. Dodrantalem. *p.* 295. *l.* 14. l. Piſtacia. *p.* 307. *l. penult.* l. Gallicè & pigra. *p.* 305. *l. antepenult.* l. Germanica. *p.* 318. *l. penult.* l. Britannia. 319. *l.* 19. l. prior vulgaris. *p.* 320. *antepenult.* l. minutæ *p.* 340. *l.* 1. l. radice. *p.* 341. *l. penult.* l. expetitur. *p.* 371. *l. antepenult.* l. propriè dicta. *p.* 372. *l.* 11. l. averruncandum. *p.* 384. *l.* 13. l. fecerit *p.* 392. *l.* 11, & 12. l. denominandæ. *p.* 417. *l.* 17. l. folliculos. *p.* 419. *l.* 23. l. mimoſam. *p.* 440. *l.* 10. l. Cachryophora. *ibid. l.* 18. l. ferularum *p.* 464. *l.* 7. l. præſtantiſſima & certiſſima. *ibid. l. penult.* l edita. *p.* 476. *l.* 3. Dod. *deleatur*. *ibid. l.* 7. *pro* à l. videlicet. *p.* 488. *l.* 12. l. Ceterach. *p.* 496. *l.* 16. l. vis ut apertis.

VIRIS

VIRIS CLARISSIMIS

In re

BOTANICA,

Facilè inter primos Magiſtris.

ABELI BRUNTER Medico & Botanico, *Celſitudini* E. Regiæ.

NICOLAO MARCHANT Apothecario, Botanico *Celſitud.* E. Regiæ.

Collegis ſuis, Horti Regii *Bleſenſis* Collectoribus & coexaminatoribus digniſſimis: Author hæc ſcribit.

Collegæ digniſſimi,

CUM *Hortum Regium*, Bleſenſem, *trecentis ſexaginta plantis, plus minúſve auctum animadverteram: quæ fuêre à nobis recuperatæ & collectæ ab anno* 1655. (*quo anno Hortum Regium* Bleſenſem, *Typis pro ſecundâ vice commiſimus*) *uſque ad annum* 1660, *quo extinctus eſt ſereniſſimus noſter Mœcenas & Patronus, verè Regius, digniſſimus*

Digitized by Google

niſſimus Princeps. Cum hunc catalogum cum auctario à quamplurimis paſſim deſiderari obſervaverim. Eundem auctum, cum durationis notulis, aliiſque notis ſeu characteriſmis, nulli cuiquam Botanicorum *hucuſque uſitatis, in lucem edere decrevi. Partìm ut* Botanicorum *tàm hìc, quàm alibì terrarum degentium, precibus ſatisfacerem; partìm ut nobiſmetipſis ſatisfieret; quippe ſudoribus & laboribus noſtris per decennium exantlatis, collectæ & examinatæ fuerunt quotquot hìc exſtant plantæ. Quum ſub Auſpiciis ſereniſſimi Domini noſtri,* Gaſtonis Franciæ, *&c. hæc à me incepta & hìc elaborata & finita; ne blattis & tineis diutius committerentur, in* Botanicorum *rem eſſe mecum non dubitabitis: Quare publici juris facere animum induxi meum. Porro plantas non ſcriptas in eodem Horto Regio* Blæſenſi *auctо* 260 *contentas, brevitèr & ſuccinctè delineatas exhibeo; Cum obſervationibus generalioribus plantarum ex eodem Horto Regio* Bleſenſi *auctо excerptis, rei herbariæ amantibus neceſſariis & cognitu perutilibus. Hallucinationes* Caſpari Bauhini *tàm in digerendis quam denominandis plantis. Animadverſiones itidem in* Johannis Bauhini *tres To-*

mos

nios historiæ suæ universalis edidi: Unà cum Dialogo inter Philobotanum D. B. popularem meum Societatis Regiæ Socium *& memetipsum. Ubi passim in hac meâ doctrinâ præsterno viam ad methodum novam & inauditam quam brevi (si Deus dederit) lucem videre spero. Vobis hæc scribere (quicquid æquum est) judicavi. Vestrum judicium, vestra existimatio valebit. Sublimiora sub Auspiciis serenissimi* CAROLI *Secundi, magnæ* Britanniæ, Franciæ *&* Hiberniæ *Regis Domini nunc nostri; jam penè in promptu habeo. Videlicet, classes Stirpium, per Tabulas cognationis seu affinitatis dispositas.* Diu Valete

Collegæ dignissimi,

Vester Collega,

Collector & coexaminator

Robertus Morison M. D.
Botan. *Celsitud. R. Regiæ*
Blesis degens.

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

Digitized by Google

BIBLIOTHECA
REGIA
MONACENSIS.